Christoph Schiller

MOTION MOUNTAIN


THE ADVENTURE OF PHYSICS - VOL.I
FALL, FLOW AND HEAT





Christoph Schiller

MOTION MOUNTAIN



The Adventure of Physics Volume I

Fall, Flow and Heat

Edition 25.04, available as free pdf at www.motionmountain.net

Editio vicesima quinta.

Proprietas scriptoris © Chrestophori Schiller primo anno Olympiadis trigesimae.

Omnia proprietatis iura reservantur et vindicantur. Imitatio prohibita sine auctoris permissione. Non licet pecuniam expetere pro aliqua, quae partem horum verborum continet; liber pro omnibus semper gratuitus erat et manet.

Twenty-fifth edition.

Copyright © 2012 by Christoph Schiller, the first year of the 30th Olympiad.



This pdf file is licensed under the Creative Commons
Attribution-Noncommercial-No Derivative Works 3.0 Germany
Licence, whose full text can be found on the website
creativecommons.org/licenses/by-nc-nd/3.0/de,
with the additional restriction that reproduction, distribution and use,
in whole or in part, in *any* product or service, be it
commercial or not, is not allowed without the written consent of
the copyright owner. The pdf file was and remains free for everybody
to read, store and print for personal use, and to distribute

electronically, but only in unmodified form and at no charge.







Primum movere, deinde docere.*

Antiquity

This book is written for anybody who is curious about nature and motion. Curiosity about how people, animals, things, images and space move leads to many adventures. This volume presents the best of them in the domain of everyday life.

Carefully observing everyday motion allows us to deduce six essential statements: everyday motion is continuous, conserved, relative, reversible, mirror-invariant – and lazy. Yes, nature is indeed lazy: in every motion, it minimizes change. This text explores how these six results are deduced and how they fit with all those observations that seem to contradict them.

In the structure of modern physics, shown in Figure 1, the results on everyday motion form the major part of the starting point at the bottom. The present volume is the first of a six-volume overview of physics. It resulted from a threefold aim I have pursued since 1990: to present motion in a way that is simple, up to date and captivating.

In order to be *simple*, the text focuses on concepts, while keeping mathematics to the necessary minimum. Understanding the concepts of physics is given precedence over using formulae in calculations. The whole text is within the reach of an undergraduate.

In order to be *up to date*, the text is enriched by the many gems – both theoretical and empirical – that are scattered throughout the scientific literature.

In order to be *captivating*, the text tries to startle the reader as much as possible. Reading a book on general physics should be like going to a magic show. We watch, we are astonished, we do not believe our eyes, we think, and finally we understand the trick. When we look at nature, we often have the same experience. Indeed, every page presents at least one surprise or provocation for the reader to think about. Numerous interesting challenges are proposed.

The motto of the text, *die Menschen stärken*, *die Sachen klären*, a famous statement by Hartmut von Hentig on pedagogy, translates as: 'To fortify people, to clarify things.' Clarifying things requires courage, as changing habits of thought produces fear, often hidden by anger. But by overcoming our fears we grow in strength. And we experience intense and beautiful emotions. All great adventures in life allow this, and exploring motion is one of them. Enjoy it!

Munich, 22 March 2012.

^{* &#}x27;First move, then teach.' In modern languages, the mentioned type of *moving* (the heart) is called *motivating*; both terms go back to the same Latin root.

8 PREFACE




FIGURE 1 A complete map of physics: the connections are defined by the speed of light c, the gravitational constant G, the Planck constant G

ADVICE FOR LEARNERS

In my experience, there is one learning method that never fails to transform unsuccessful pupils into successful ones: if you read a book for study, summarize every section you read, *in your own words, aloud*. If you are unable to do so, read the section again. Repeat this until you can clearly summarize what you read in your own words, aloud. You can do this alone in a room, or with friends, or while walking. If you do this with everything you read, you will reduce your learning and reading time significantly.

The most inefficient learning method is to use a marker or to underline text: it wastes time, provides false comfort and makes the text unreadable. Nobody marking text is an efficient learner. Instead, by repeating every section in your own words, aloud, you will

PREFACE 9

save time and money, enjoy learning from good texts much more and hate bad texts much less. Masters of the method can use it even while listening to a lecture, in a low voice, thus avoiding to ever take notes.

ADVICE FOR TEACHERS

A teacher likes pupils and enjoys his topic. A teacher likes to lead pupils into exploring the field he chose; his or her enthousiasm for the job is the key to job satisfaction. If you are a teacher, at the start of every lesson, picture, feel and tell yourself your own enthousiasm for the topic of the day, and how you want to lead each of your pupils into enjoying it as much as you do. Do this exercise consciously, before every lesson. You will minimize trouble in your class and maximize your success.

Using this book

Text in green, as found in many marginal notes, marks a link that can be clicked in a pdf reader. Such green links are either bibliographic references, footnotes, cross references to other pages, challenge solutions, or pointers to websites.

Solutions and hints for *challenges* are given in the appendix. Challenges are classified as research level (r), difficult (d), standard student level (s) and easy (e). Challenges of type r, d or s for which no solution has yet been included in the book are marked (ny).

FEEDBACK AND SUPPORT

This text is and will remain free to download from the internet. I would be delighted to receive an email from you at fb@motionmountain.net, especially on the following issues:

Challenge 1 s

- What was unclear and should be improved?
- What story, topic, riddle, picture or movie did you miss?
- What should be corrected?

In order to simplify annotations, the pdf file allows adding yellow sticker notes in Adobe Reader

Alternatively, you can provide feedback on www.motionmountain.net/wiki. Help on the specific points listed on the www.motionmountain.net/help.html web page would be particularly welcome. All feedback will be used to improve the next edition. On behalf of all readers, thank you in advance for your input. For a particularly useful contribution you will be mentioned – if you want – in the acknowledgements, receive a reward, or both.

Your donation to the charitable, tax-exempt non-profit organisation that produces, translates and publishes this book series is welcome! For details, see the web page www. motionmountain.net/donation.html. If you want, your name will be included in the sponsor list. Thank you in advance for your help, on behalf of all readers across the world.

A paper edition of this book, printed on demand and delivered by mail to any address, can be ordered at www.lulu.com/spotlight/motionmountain. But above all, enjoy the reading!

10 PREFACE



CONTENTS

- 15 1 WHY SHOULD WE CARE ABOUT MOTION?
 - Does motion exist? 16 How should we talk about motion? 18 What are the types of motion? 20 Perception, permanence and change 24 Does the world need states? 26 Galilean physics in six interesting statements 28 Curiosities and fun challenges about motion 29 Summary on motion 32
- 33 2 From motion measurement to continuity

What is velocity? 34 • What is time? 39 • Clocks 44 • Why do clocks go clockwise? 47 • Does time flow? 47 • What is space? 48 • Are space and time absolute or relative? 52 • Size – why area exists, but volume does not 53 • What is straight? 57 • A hollow Earth? 57 • Curiosities and fun challenges about everyday space and time 58 • Summary about everyday space and time 70

- 72 3 How to describe motion kinematics
 - Throwing, jumping and shooting 75 Enjoying vectors 77 What is rest? What is velocity? 78 Acceleration 80 Objects and point particles 82 Legs and wheels 85 Curiosities and fun challenges about kinematics 87 Summary of kinematics 89
- 90 4 FROM OBJECTS AND IMAGES TO CONSERVATION

Motion and contact 91 • What is mass? 92 • Momentum and mass 94 • Is motion eternal? – Conservation of momentum 98 • More conservation – energy 101 • The cross product, or vector product 104 • Rotation and angular momentum 107 • Rolling wheels 111 • How do we walk? 112 • Curiosities and fun challenges about conservation and rotation 113 • Summary on conservation 121

- 123 5 FROM THE ROTATION OF THE EARTH TO THE RELATIVITY OF MOTION

 How does the Earth rotate? 131 Does the Earth move? 135 Is velocity absolute?

 The theory of everyday relativity 141 Is rotation relative? 142 Curiosities and fun challenges about relativity 142 Legs or wheels? Again 148 Summary on Galilean relativity 150
- 151 6 MOTION DUE TO GRAVITATION

Properties of gravitation 155 • The gravitational potential 160 • The shape of the Earth 161 • Dynamics – how do things move in various dimensions? 163 • Gravitation in the sky 164 • The Moon 167 • Orbits – and conic sections 168 • Tides 172 • Can light fall? 177 • What is mass? – Again 177 • Curiosities and fun challenges about gravitation 179 • Summary on gravitation 194

- 7 CLASSICAL MECHANICS AND THE PREDICTABILITY OF MOTION

 Should one use force? Power? 195 Forces, surfaces and conservation 198 Friction and motion 198 Friction, sport, machines and predictability 200 Complete states initial conditions 203 Do surprises exist? Is the future determined? 204 Free will 206 Summary on predictability 207 Predictability and global descriptions of motion 208
- 213 8 MEASURING CHANGE WITH ACTION

The principle of least action $_{217}$ • Lagrangians and motion $_{220}$ • Why is motion so often bounded? $_{222}$ • Curiosities and fun challenges about Lagrangians $_{225}$ • Summary on action $_{228}$

12 CONTENTS

229 9 MOTION AND SYMMETRY

Why can we think and talk about the world? 230 • Viewpoints 233 • Symmetries and groups 235 • Representations 235 • Symmetries, motion and Galilean physics 238 • Reproducibility, conservation and Noether's theorem 241 • Curiosities and fun challenges about symmetry 245 • Parity and time invariance 246 • Summary on symmetry 246

248 10 SIMPLE MOTIONS OF EXTENDED BODIES – OSCILLATIONS AND WAVES
Oscillations 248 • Resonance 251 • Waves: general and harmonic 253 • Water
waves 254 • Waves and their motion 258 • Why can we talk to each other? –
Huygens' principle 261 • Why is music so beautiful? 263 • Is ultrasound imaging
safe for babies? 266 • Signals 267 • Solitary waves and solitons 268 • Curiosities
and fun challenges about waves and extended bodies 271 • Summary on waves
and oscillations 281

283 11 DO EXTENDED BODIES EXIST? - LIMITS OF CONTINUITY

Mountains and fractals 283 • Can a chocolate bar last forever? 283 • The case of Galileo Galilei 285 • How high can animals jump? 287 • Felling trees 287 • Little hard balls 288 • The sound of silence 289 • How to count what cannot be seen 289 • Experiencing atoms 291 • Seeing atoms 292 • Curiosities and fun challenges about solids 294 • Summary on atoms 300

302 12 FLUIDS AND THEIR MOTION

The state of a fluid 302 • The state of a fluid 302 • Curiosities and fun challenges about fluids 307 • What can move in nature? – Flows 316 • Summary on fluids 318

319 13 FROM HEAT TO TIME-INVARIANCE

Temperature 319 • Thermal energy 321 • Why do balloons take up space? – The end of continuity 324 • Brownian motion 326 • Why stones can be neither smooth nor fractal, nor made of little hard balls 329 • Entropy 329 • Entropy from particles 332 • The minimum entropy of nature – the quantum of information 333 • Is everything made of particles? 334 • The second principle of thermodynamics 336 • Why can't we remember the future? 337 • Flow of entropy 338 • Do isolated systems exist? 339 • Curiosities and fun challenges about heat and reversibility 339 • Summary on heat and time-invariance 347

- 348 14 SELF-ORGANIZATION AND CHAOS THE SIMPLICITY OF COMPLEXITY
 - Appearance of order 351 Self-organization in sand 352 Self-organization of spheres 355 Appearance of order 355 The mathematics of order appearance 356 Chaos 357 Emergence 358 Curiosities and fun challenges about self-organization 359 Summary on self-organization and chaos 364
- 366 15 FROM THE LIMITATIONS OF PHYSICS TO THE LIMITS OF MOTION

 Research topics in classical dynamics 366 What is contact? 367 Precision and accuracy 368 Can all of nature be described in a book? 368 Something is wrong about our description of motion 369 Why is measurement possible? 370 Is motion unlimited? 370
- 372 A NOTATION AND CONVENTIONS

The Latin alphabet 372 • The Greek alphabet 374 • The Hebrew alphabet and other scripts 376 • Numbers and the Indian digits 376 • The symbols used in the text 377 • Calendars 379 • People Names 381 • Abbreviations and eponyms or concepts? 381

CONTENTS 13

```
383 B UNITS, MEASUREMENTS AND CONSTANTS
SI units 383 • The meaning of measurement 386 • Curiosities and fun challenges about units 386 • Precision and accuracy of measurements 388 • Limits to precision 390 • Physical constants 390 • Useful numbers 397
```

- 398 C Sources of information on motion
- 403 CHALLENGE HINTS AND SOLUTIONS
- 444 BIBLIOGRAPHY
- 474 CREDITS

Acknowledgements 474 • Film credits 475 • Image credits 475





FALL, FLOW AND HEAT

In our quest to learn how things move, the experience of hiking and other motion leads us to introduce the concepts of velocity, time, length, mass and temperature. We learn to use them to *measure change* and find that nature minimizes it. We discover how to float in free space, why we have legs instead of wheels, why disorder can never be eliminated, and why one of the most difficult open issues in science is the flow of water through a tube.

Ref. 1

Ref. 2

WHY SHOULD WE CARE ABOUT MOTION?

All motion is an illusion.

Zeno of Elea*

THAM! The lightning striking the tree nearby violently disrupts our quiet forest alk and causes our hearts to suddenly beat faster. In the top of the tree e see the fire start and fade again. The gentle wind moving the leaves around prestore the calminess of the place. Nearby, the water in a small river follows.

us helps to restore the calmness of the place. Nearby, the water in a small river follows its complicated way down the valley, reflecting on its surface the ever-changing shapes of the clouds.

Motion is everywhere: friendly and threatening, terrible and beautiful. It is fundamental to our human existence. We need motion for growing, for learning, for thinking and for enjoying life. We use motion for walking through a forest, for listening to its noises and for talking about all this. Like all animals, we rely on motion to get food and to survive dangers. Like all living beings, we need motion to reproduce, to breathe and to digest. Like all objects, motion keeps us warm.

Motion is the most fundamental observation about nature at large. It turns out that *everything* that happens in the world is some type of motion. There are no exceptions. Motion is such a basic part of our observations that even the origin of the word is lost in the darkness of Indo-European linguistic history. The fascination of motion has always made it a favourite object of curiosity. By the fifth century BCE in ancient Greece, its study had been given a name: *physics*.

Motion is also important to the human condition. What can we know? Where does the world come from? Who are we? Where do we come from? What will we do? What should we do? What will the future bring? What is death? Where does life lead? All these questions are about motion. The study of motion provides answers that are both deep and surprising.

Motion is mysterious. Though found everywhere – in the stars, in the tides, in our eyelids – neither the ancient thinkers nor myriads of others in the 25 centuries since then have been able to shed light on the central mystery: *what is motion?* We shall discover that the standard reply, 'motion is the change of place in time', is inadequate. Just recently an answer has finally been found. This is the story of the way to find it.

Motion is a part of human experience. If we imagine human experience as an island, then destiny, symbolized by the waves of the sea, carried us to its shore. Near the centre of

Aotion Mountain – The Adventure of Physics 🛮 pdf file available free of charge at www.motionmountain.net 🛮 Copyright © Christoph Schiller November 1997–March

^{*} Zeno of Elea (c. 450 BCE), one of the main exponents of the Eleatic school of philosophy.




FIGURE 2 Experience Island, with Motion Mountain and the trail to be followed.

the island an especially high mountain stands out. From its top we can see over the whole landscape and get an impression of the relationships between all human experiences, and in particular between the various examples of motion. This is a guide to the top of what I have called Motion Mountain (see Figure 2; a less artistic but more exact version is given in Figure 1). The hike is one of the most beautiful adventures of the human mind. The first question to ask is:

Does motion exist?

Das Rätsel gibt es nicht. Wenn sich eine Frage überhaupt stellen läßt, so kann sie beantwortet werden.*

Ludwig Wittgenstein, Tractatus, 6.5

To sharpen the mind for the issue of motion's existence, have a look at Figure 3 or Figure 4 and follow the instructions. In all cases the figures seem to rotate. You can experience similar effects if you walk over cobblestone pavement that is arranged in arched patterns or if you look at the numerous motion illusions collected by Kitaoka Akiyoshi at www.ritsumei.ac.jp/~akitaoka. How can we make sure that real motion is different from these or other similar illusions?


Many scholars simply argued that motion does not exist at all. Their arguments deeply influenced the investigation of motion over many centuries. For example, the Greek

Ref.

Ref. 5

Ref. 4 Challenge 2 s

^{* &#}x27;The riddle does not exist. If a question can be put at all, it can also be answered.'






FIGURE 3 Illusions of motion: look at the figure on the left and slightly move the page, or look at the white dot at the centre of the figure on the right and move your head back and forward.

philosopher Parmenides (born *c.* 515 BCE in Elea, a small town near Naples) argued that since nothing comes from nothing, change cannot exist. He underscored the *permanence* of nature and thus consistently maintained that all change and thus all motion is an illusion.

Ref. 6

Heraclitus (c. 540 to c. 480 BCE) held the opposite view. He expressed it in his famous statement πάντα ῥεῖ 'panta rhei' or 'everything flows'.* He saw change as the essence of nature, in contrast to Parmenides. These two equally famous opinions induced many scholars to investigate in more detail whether in nature there are *conserved* quantities or whether *creation* is possible. We will uncover the answer later on; until then, you might ponder which option you prefer.

Challenge 3 s

Parmenides' collaborator Zeno of Elea (born *c*. 500 BCE) argued so intensely against motion that some people still worry about it today. In one of his arguments he claims – in simple language – that it is impossible to slap somebody, since the hand first has to travel halfway to the face, then travel through half the distance that remains, then again so, and so on; the hand therefore should never reach the face. Zeno's argument focuses on the relation between *infinity* and its opposite, finitude, in the description of motion. In modern quantum theory, a related issue is a subject of research up to this day.

Ref. 7

Zeno also maintained that by looking at a moving object at a *single* instant of time, one cannot maintain that it moves. He argued that at a single instant of time, there is no difference between a moving and a resting body. He then deduced that if there is no difference at a single time, there cannot be a difference for longer times. Zeno therefore questioned whether motion can clearly be distinguished from its opposite, *rest.* Indeed, in the history of physics, thinkers switched back and forward between a positive and a negative answer. It was this very question that led Albert Einstein to the development of general relativity, one of the high points of our journey. In our adventure, we will explore all known differences between motion and rest. Eventually, we will dare to ask whether single instants of time do exist at all. Answering this question is essential for reaching the top of Motion Mountain.

When we explore quantum theory, we will discover that motion is indeed – to a certain extent – an illusion, as Parmenides claimed. More precisely, we will show that motion is observed only due to the limitations of the human condition. We will find that we experience motion only because

^{*} Appendix A explains how to read Greek text.

FIGURE 4 Zoom this image to large size or approach it closely in order to enjoy its apparent motion (© Michael Bach after the discovery of Kitaoka Akiyoshi).

- we have a finite size,
- we are made of a large but finite number of atoms,
- we have a finite but moderate temperature,
- we are electrically neutral,
- we move much more slowly than the speed of light,
- we live in three dimensions,
- we are large compared with a black hole of our same mass,
- we are large compared with our quantum mechanical wavelength,
- we are small compared with the universe,
- we have a limited memory,
- we are forced by our brain to approximate space and time as continuous entities, and
- we are forced by our brain to approximate nature as made of different parts.

If any one of these conditions were not fulfilled, we would not observe motion; motion, then, would not exist! If that were not enough, note that none of the conditions requires human beings; they are equally valid for many animals and machines. Each of these conditions can be uncovered most efficiently if we start with the following question:

How should we talk about motion?

Le hais le mouvement, qui déplace les lignes, Et jamais je ne pleure et jamais je ne ris. Charles Baudelaire, *La Beauté*.*

^{*} Charles Baudelaire (b. 1821 Paris, d. 1867 Paris) *Beauty*: 'I hate movement, which changes shapes, and never do I weep and never do I laugh.'




FIGURE 5 A time line of scientific and *political* personalities in antiquity (the last letter of the name is aligned with the year of death).

Like any science, the approach of physics is twofold: we advance with *precision* and with *curiosity*. Precision makes meaningful communication possible, and curiosity makes it worthwhile. Be it an eclipse, a beautiful piece of music, or a feat at the Olympic games: the world is full of fascinating examples of motion.*

If you ever find yourself talking about motion, whether to understand it more precisely or more deeply, you are taking steps up Motion Mountain. The examples of Figure 6 make the point. When you fill a bucket with a small amount of water, it does not hang vertically. (Why?) If you continue adding water, it starts to hang vertically at a certain moment. How much water is necessary? When you pull a thread from a reel in the way shown, the reel will move either forwards or backwards, depending on the angle at which you pull. What is the limiting angle between the two possibilities?

High precision means going into fine details, and being attuned to details actually *increases* the pleasure of the adventure.** Figure 7 shows an example. The higher we get on Motion Mountain, the further we can see and the more our curiosity is rewarded. The views offered are breathtaking, especially from the very top. The path we will follow – one of the many possible routes – starts from the side of biology and directly enters the forest that lies at the foot of the mountain.

Intense curiosity drives us to go straight to the limits: understanding motion requires exploration of the largest distances, the highest velocities, the smallest particles, the strongest forces and the strangest concepts. Let us begin.

Challenge 4 s

Ref. 9 Challenge 5 s

Ref. 10

^{*} For a collection of interesting examples of motion in everyday life, see the excellent book by Walker.

^{**} Distrust anybody who wants to talk you *out* of investigating details. He is trying to deceive you. Details are important. Be vigilant also during *this* journey.






FIGURE 6 How much water is required to make a bucket hang vertically? At what angle does the pulled reel change direction of motion? (© Luca Gastaldi).



FIGURE 7 An example of how precision of observation can lead to the discovery of new effects: the deformation of a tennis ball during the *c*. 6 ms of a fast bounce (© International Tennis Federation).

WHAT ARE THE TYPES OF MOTION?

Every movement is born of a desire for change.

Antiquity

A good place to obtain a general overview on the types of motion is a large library (see Table 1). The domains in which motion, movements and moves play a role are indeed varied. Already the earliest researchers in ancient Greece – listed in Figure 5 – had the suspicion that all types of motion, as well as many other types of change, are related. Three categories of change are commonly recognized:

- 1. *Transport*. The only type of change we call motion in everyday life is material transport, such as a person walking, a leaf falling from a tree, or a musical instrument playing. Transport is the change of position or orientation of objects, fluids included. To a large extent, the behaviour of people also falls into this category.
- 2. Transformation. Another category of change groups observations such as the dissolution of salt in water, the formation of ice by freezing, the rotting of wood, the cooking of food, the coagulation of blood, and the melting and alloying of metals. These changes of colour, brightness, hardness, temperature and other material properties are all transformations. Transformations are changes not visibly connected with transport. To this category, a few ancient thinkers added the emission and absorption of light. In the twentieth century, these two effects were proven to be special cases of transformations, as were the newly discovered appearance and disappearance of mat-

Ref. 18

TABLE 1 Content of books about motion found in a public library.

MOTION TOPICS	MOTION TOPICS
motion pictures and digital effects	motion as therapy for cancer, diabetes, acne and depression
motion perception Ref. 11	motion sickness
motion for fitness and wellness	motion for meditation
motion control and training in sport and singing	l motion ability as health check
perpetual motion	motion in dance, music and other performing arts
motion as proof of various gods Ref. 12	motion of planets, stars and angels Ref. 13
economic efficiency of motion	the connection between motional and emotional habits
motion as help to overcome trauma	motion in psychotherapy Ref. 14
locomotion of insects, horses, animals and robots	l motion of cells and plants
collisions of atoms, cars, stars and galaxies	growth of multicellular beings, mountains, sunspots and galaxies
motion of springs, joints, mechanisms, liq	-motion of continents, bird flocks, shadows and
uids and gases	empty space
commotion and violence	motion in martial arts
motions in parliament	movements in art, sciences and politics
movements in watches	movements in the stock market
movement teaching and learning	movement development in children Ref. 15
musical movements	troop movements Ref. 16
religious movements	bowel movements
moves in chess	cheating moves in casinos Ref. 17
connection between gross national product a	and citizen mobility

ter, as observed in the Sun and in radioactivity. *Mind change*, such as change of mood, of health, of education and of character, is also (mostly) a type of transformation.

Ref. 19 3. *Growth*. This last and especially important category of change, is observed for animals, plants, bacteria, crystals, mountains, planets, stars and even galaxies. In the nineteenth century, changes in the population of systems, *biological evolution*, and in the twentieth century, changes in the size of the universe, *cosmic evolution*, were added to this category. Traditionally, these phenomena were studied by separate sciences. Independently they all arrived at the conclusion that growth is a combination of transport and transformation. The difference is one of complexity and of time scale.

At the beginnings of modern science during the Renaissance, only the study of transport was seen as the topic of physics. Motion was equated to transport. The other two domains were neglected by physicists. Despite this restriction, the field of enquiry remains large, covering a large part of Experience Island. Early scholars differentiated types of transport by their origin. Movements such as those of the legs when walking were classified



FIGURE 8 An example of transport, at Mount Etna (© Marco Fulle).

as *volitional*, because they are controlled by one's will, whereas movements of external objects, such as the fall of a snowflake, which cannot be influenced by will-power, were classified as *passive*. Young humans, especially young male humans, spend considerable time in learning elaborate volitional movements. An example is shown in Figure 10.

The complete distinction between passive and volitional motion is made by children by the age of six, and this marks a central step in the development of every human towards a precise description of the environment.* From this distinction stems the historical but now outdated definition of physics as the science of the motion of non-living things.

The advent of machines forced scholars to rethink the distinction between volitional and passive motion. Like living beings, machines are self-moving and thus mimic volitional motion. However, careful observation shows that every part in a machine is moved by another, so their motion is in fact passive. Are living beings also machines? Are human actions examples of passive motion as well? The accumulation of observations in the last 100 years made it clear that volitional movement** indeed has the same physical properties as passive motion in non-living systems. (Of course, from the emotional viewpoint, the differences are important; for example, *grace* can only be ascribed to volitional movements.) A distinction between the two types of motion is thus unnecessary. But since passive and volitional motion have the same properties, through the study of motion of non-living objects we can learn something about the human condition. This

Ref. 20

^{*} Failure to pass this stage completely can result in a person having various strange beliefs, such as believing in the ability to influence roulette balls, as found in compulsive players, or in the ability to move other bodies by thought, as found in numerous otherwise healthy-looking people. An entertaining and informative account of all the deception and self-deception involved in creating and maintaining these beliefs is given by James Randi, *The Faith Healers*, Prometheus Books, 1989. A professional magician, he presents many similar topics in several of his other books. See also his www.randi.org website for more details.

^{**} The word 'movement' is rather modern; it was imported into English from the old French and became popular only at the end of the eighteenth century. It is never used by Shakespeare.



FIGURE 9 Transport, growth and transformation (© Philip Plisson).



FIGURE 10 One of the most difficult volitional movements known, performed by Alexander Tsukanov, the first man able to do this: jumping from one ultimate wheel to another (© Moscow State Circus).

is most evident when touching the topics of determinism, causality, probability, infinity, time, love and death, to name but a few of the themes we will encounter during our adventure.

In the nineteenth and twentieth centuries other classically held beliefs about motion fell by the wayside. Extensive observations showed that all transformations and all growth phenomena, including behaviour change and evolution, are also examples of transport. In other words, over 2 000 years of studies have shown that the ancient classification of observations was useless: *all change is transport*.

In the middle of the twentieth century the study of motion culminated in the experimental confirmation of an even more specific idea, previously articulated in ancient Greece:

▶ Every type of change is due to the motion of particles.

It takes time and work to reach this conclusion, which appears only when we relentlessly pursue higher and higher precision in the description of nature. The first five parts of this adventure retrace the path to this result. (Do you agree with it?)

Challenge 6 s

Ref. 21

The last decade of the twentieth century again completely changed the description of motion: the particle idea turns out to be wrong. This new result, reached through a combination of careful observation and deduction, will be explored in the last part of our adventure. But we still have some way to go before we reach that part, just below the summit of our journey.

In summary, history has shown that classifying the various types of motion is not productive. Only by trying to achieve maximum precision can we hope to arrive at the fundamental properties of motion. *Precision, not classification, is the path to follow.* As Ernest Rutherford said: 'All science is either physics or stamp collecting.'

To achieve precision in our description of motion, we need to select specific examples of motion and study them fully in detail. It is intuitively obvious that the most precise description is achievable for the *simplest* possible examples. In everyday life, this is the case for the motion of any non-living, solid and rigid body in our environment, such as a stone thrown through the air. Indeed, like all humans, we learned to throw objects long before we learned to walk. Throwing is one of the first physical experiments we performed by ourselves.* During our early childhood, by throwing stones, toys and other objects until our parents feared for every piece of the household, we explored the perception and the properties of motion. We do the same here.

Die Welt ist unabhängig von meinem Willen.**
Ludwig Wittgenstein, *Tractatus*, 6.373

PERCEPTION, PERMANENCE AND CHANGE

Only wimps study only the general case; real scientists pursue examples.

Beresford Parlett

Human beings enjoy perceiving. Perception starts before birth, and we continue enjoying it for as long as we can. That is why television, even when devoid of content, is so successful. During our walk through the forest at the foot of Motion Mountain we cannot avoid perceiving. Perception is first of all the ability to *distinguish*. We use the basic mental act of distinguishing in almost every instant of life; for example, during childhood we first learned to distinguish familiar from unfamiliar observations. This is possible in combination with another basic ability, namely the capacity to *memorize* experiences. Memory gives us the ability to experience, to talk and thus to explore nature. Perceiving, classifying and memorizing together form *learning*. Without any one of these three abilities, we could not study motion.



^{*} The importance of throwing is also seen from the terms derived from it: in Latin, words like *subject* or 'thrown below', *object* or 'thrown in front', and *interjection* or 'thrown in between'; in Greek, it led to terms like *symbol* or 'thrown together', *problem* or 'thrown forward', *emblem* or 'thrown into', and – last but not least – *devil* or 'thrown through'.

^{** &#}x27;The world is independent of my will.'



FIGURE 11 How do we distinguish a deer from its environment? (© Tony Rodgers).

Children rapidly learn to distinguish *permanence* from *variability*. They learn to *recognize* human faces, even though a face never looks exactly the same each time it is seen. From recognition of faces, children extend recognition to all other observations. Recognition works pretty well in everyday life; it is nice to recognize friends, even at night, and even after many beers (not a challenge). The act of recognition thus always uses a form of *generalization*. When we observe, we always have some general idea in our mind. Let us specify the main ones.

Sitting on the grass in a clearing of the forest at the foot of Motion Mountain, surrounded by the trees and the silence typical of such places, a feeling of calmness and tranquillity envelops us. We are thinking about the essence of perception. Suddenly, something moves in the bushes; immediately our eyes turn and our attention focuses. The nerve cells that detect motion are part of the most ancient part of our brain, shared with birds and reptiles: the brain stem. Then the cortex, or modern brain, takes over to analyse the type of motion and to identify its origin. Watching the motion across our field of vision, we observe two invariant entities: the fixed landscape and the moving animal. After we recognize the animal as a deer, we relax again.

How did we distinguish, in case of Figure 11, between landscape and deer? Perception involves several processes in the eye and in the brain. An essential part for these processes is motion, as is best deduced from the flip film shown in the lower left corners of these pages. Each image shows only a rectangle filled with a mathematically random pattern. But when the pages are scanned in rapid succession, you discern a shape – a square – moving against a fixed background. At any given instant, the square cannot be distinguished from the background; there is no visible object at any given instant of time. Nevertheless it is easy to perceive its motion.* Perception experiments such as this one have been performed in many variations. For example, it was found that detecting a moving square against a random background is nothing special to humans; flies have the same ability, as do, in fact, all animals that have eyes.

^{*} The human eye is rather good at detecting motion. For example, the eye can detect motion of a point of light even if the change of angle is smaller than that which can be distinguished in a fixed image. Details of this and similar topics for the other senses are the domain of perception research.





The flip film in the lower left corner, like many similar experiments, illustrates two central attributes of motion. First, motion is perceived only if an object can be distinguished from a background or environment. Many motion illusions focus on this point.* Second, motion is required to define both the object and the environment, and to distinguish them from each other. In fact, the concept of space is - among others - an abstraction of the idea of background. The background is extended; the moving entity is localized. Does this seem boring? It is not; just wait a second.

We call the set of localized aspects that remain invariant or permanent during motion, such as size, shape, colour etc., taken together, a (physical) object or a (physical) body. We will tighten the definition shortly, since otherwise images would be objects as well. In other words, right from the start we experience motion as a *relative* process; it is perceived in relation and in opposition to the environment. The concept of an object is therefore also a relative concept. But the basic conceptual distinction between localized, isolable objects and the extended environment is not trivial or unimportant. First, it has the appearance of a circular definition. (Do you agree?) This issue will keep us busy later on. Second, we are so used to our ability of isolating local systems from the environment that we take it for granted. However, as we will see in the last part of our walk, this distinction turns out to be logically and experimentally impossible!** Our walk will lead us to discover the reason for this impossibility and its important consequences. Finally, apart from moving entities and the permanent background, we need a third concept, as shown in Table 2.

Challenge 7 s Page 369

Vol. VI, page 78

Ref. 24

Wisdom is one thing: to understand the thought which steers all things through all things. Heraclitus of Ephesus

Does the world need states?

Oas Feste, das Bestehende und der Gegenstand sind Eins. Der Gegenstand ist das Feste, Bestehende; die Konfiguration ist das Wechselnde, Unbeständige.*** Ludwig Wittgenstein, Tractatus, 2.027 - 2.0271

What distinguishes the various patterns in the lower left corners of this text? In everyday life we would say: the situation or configuration of the involved entities. The situation somehow describes all those aspects that can differ from case to case. It is customary to call the list of all variable aspects of a set of objects their (physical) state of motion, or simply their state.

The situations in the lower left corners differ first of all in time. Time is what makes opposites possible: a child is in a house and the same child is outside the house. Time




^{*} The topic of motion perception is full of interesting aspects. An excellent introduction is chapter 6 of the beautiful text by Donald D. Hoffman, Visual Intelligence - How We Create What We See, W.W. Norton & Co., 1998. His collection of basic motion illusions can be experienced and explored on the associated www.cogsci.uci.edu/~ddhoff website.

^{**} Contrary to what is often read in popular literature, the distinction is possible in quantum theory. It becomes impossible only when quantum theory is unified with general relativity.

^{*** &#}x27;The fixed, the existent and the object are one. The object is the fixed, the existent; the configuration is the changing, the variable.'

TABLE 2 Family tree of the basic physical concepts.



world – nature – universe – cosmosthe collection of all parts, relations and backgrounds

describes and resolves this type of contradiction. But the state not only distinguishes situations in time: the state contains *all* those aspects of a system (i.e., of a group of objects) that set it apart from all *similar* systems. Two objects can have the same mass, shape, colour, composition and be indistinguishable in all other intrinsic properties; but at least they will differ in their position, or their velocity, or their orientation. The state pinpoints the *individuality* of a physical system,* and allows us to distinguish it from exact copies of itself. Therefore, the state also describes the relation of an object or a system with respect to its environment. Or in short: *the state describes all aspects of a system that depend on the observer.* These properties are not boring – just ponder this: does the universe have a state?

Challenge 9 s

Describing nature as a collection of permanent entities and changing states is the starting point of the study of motion. The observation of motion requires the distinction of permanent, intrinsic properties – describing the objects that move – and changing states – describing the way the objects move. Without this distinction, there is no motion.

The various aspects of objects and of their states are called *observables*. All these rough, preliminary definitions will be refined step by step in the following. Using the terms just

^{*} A *physical system* is a localized entity of investigation. In the classification of Table 2, the term 'physical system' is (almost) the same as 'object' or 'physical body'. Images are usually not counted as physical systems (though radiation is one). Are holes physical systems?





introduced, we can say that motion is the change of state of objects.*

States are required for the description of motion. In order to proceed and to achieve a *complete* description of motion, we thus need a complete description of objects and a complete description of their possible states. The first approach, called Galilean physics, consists in specifying our *everyday* environment as precisely as possible.

GALILEAN PHYSICS IN SIX INTERESTING STATEMENTS


The study of everyday motion, Galilean physics, is already worthwhile in itself: we will uncover many results that are in contrast with our usual experience. For example, if we recall our own past, we all have experienced how important, delightful or unwelcome *surprises* can be. Nevertheless, the study of everyday motion shows that there are *no* surprises in nature. Motion, and thus the world, is *predictable* or *deterministic*.

The main surprise is thus that there are no surprises in nature. In fact, we will uncover six aspects of the predictability of everyday motion:

- 1. We know that eyes, cameras and measurement apparatus have a finite resolution. All have a smallest distance they can observe. We know that clocks have a smallest time they can measure. Nevertheless, in everyday life all movements, their states, as well as space and time, are *continuous*.
- 2. We all observe that people, music and many other things in motion stop moving after a while. The study of motion yields the opposite result: motion never stops. In fact, several aspects of motion do not change, but are *conserved*: energy with mass, momentum and angular momentum are conserved in all examples of motion. No exception to conservation has ever been found. In addition, we will discover that conservation implies that motion and its properties are the same at all places and all times: motion is *universal*.
- 3. We all know that motion differs from rest. Nevertheless, careful study shows that there is no intrinsic difference between the two. Motion and rest depend on the observer. Motion is *relative*. This is the first step towards understanding the theory of relativity.
- 4. We all observe that many processes happen only in one direction. For example, spilled milk never returns into the container by itself. Nevertheless, the study of motion will show us that all everyday motion is *reversible*. Physicists call this the invariance of everyday motion under motion reversal (or, sloppily, under time reversal).
- 5. Most of us find scissors difficult to handle with the left hand, have difficulties to write with the other hand, and have grown with a heart on the left side. Nevertheless, our exploration will show that everyday motion is *mirror-invariant* or *parity-invariant*. Mirror processes are always possible in everyday life.



^{*} The exact separation between those aspects belonging to the object and those belonging to the state depends on the precision of observation. For example, the length of a piece of wood is not permanent; wood shrinks and bends with time, due to processes at the molecular level. To be precise, the length of a piece of wood is not an aspect of the object, but an aspect of its state. Precise observations thus *shift* the distinction between the object and its state; the distinction itself does not disappear – at least not for quite a while.






FIGURE 12 A block and tackle and a differential pulley (left) and a farmer (right).

6. We all are astonished by the many observations that the world offers: colours, shapes, sounds, growth, disasters, happiness, friendship, love. The variation, beauty and complexity of nature is amazing. But despite all appearance, all motion is simple. Our study will uncover that all observations can be summarized in a simple way: Nature is lazy. All motion happens in a way that *minimizes change*. Change can be measured, and nature keeps it to a minimum. Situations – or states, as physicists like to say – evolve by minimizing change.

These six aspects are essential in understanding motion in sport, in music, in animals, in machines and among the stars. This first volume of our adventure will be an exploration of such movements. In particular, we will confirm the mentioned six key properties of everyday motion: continuity, conservation, reversibility, mirror-invariance, relativity and minimization.

CURIOSITIES AND FUN CHALLENGES ABOUT MOTION*

In contrast to most animals, sedentary creatures, like plants or sea anemones, have no legs and cannot move much; for their self-defence, they developed *poisons*. Examples of such plants are the stinging nettle, the tobacco plant, digitalis, belladonna and poppy; poisons include caffeine, nicotine, and curare. Poisons such as these are at the basis of most medicines. Therefore, most medicines exist essentially because plants have no legs.

* *

A man climbs a mountain from 9 a.m. to 1 p.m. He sleeps on the top and comes down the next day, taking again from 9 a.m. to 1 p.m. for the descent. Is there a place on the path that he passes at the same time on the two days?

Challenge 10 s

* *



^{*} Sections entitled 'curiosities' are collections of topics and problems that allow one to check and to expand the usage of concepts already introduced.

Every time a soap bubble bursts, the motion of the surface during the burst is the same,

even though it is too fast to be seen by the naked eye. Can you imagine the details? Challenge 11 s

Is the motion of a ghost an example of motion? Challenge 12 s

Can something stop moving? How would you show it? Challenge 13 s

Challenge 14 s Does a body moving forever in a straight line show that nature or space is infinite?

What is the length of rope one has to pull in order to lift a mass by a height h with a block Challenge 15 s and tackle with four wheels, as shown on the left of Figure 12? Does the farmer on the right of the figure do something sensible?

Can the universe move? Challenge 16 s

To talk about precision with precision, we need to measure precision itself. How would you do that? Challenge 17 s

Would we observe motion if we had no memory?

What is the lowest speed you have observed? Is there a lowest speed in nature?

According to legend, Sissa ben Dahir, the Indian inventor of the game of chaturanga or chess, demanded from King Shirham the following reward for his invention: he wanted one grain of wheat for the first square, two for the second, four for the third, eight for the fourth, and so on. How much time would all the wheat fields of the world take to


produce the necessary grains? Challenge 20 s

When a burning candle is moved, the flame lags behind the candle. How does the flame behave if the candle is inside a glass, still burning, and the glass is accelerated? Challenge 21 s

A good way to make money is to build motion detectors. A motion detector is a small box with a few wires. The box produces an electrical signal whenever the box moves. What types of motion detectors can you imagine? How cheap can you make such a box?

How precise? Challenge 22 d








FIGURE 13 What happens?

FIGURE 14 What is the speed of the rollers? Are other roller shapes possible?

* *

A perfectly frictionless and spherical ball lies near the edge of a perfectly flat and hori-Challenge 23 d zontal table, as shown in Figure 13. What happens? In what time scale?

* *

You step into a closed box without windows. The box is moved by outside forces un-Challenge 24 s known to you. Can you determine *how* you are moving from inside the box?

* *

When a block is rolled over the floor over a set of cylinders, as shown in Figure 14, how Challenge 25 s are the speed of the block and that of the cylinders related?

* *

Challenge 26 s

Do you dislike formulae? If you do, use the following three-minute method to change the situation. It is worth trying it, as it will make you enjoy this book much more. Life is short; as much of it as possible, like reading this text, should be a pleasure.

- 1. Close your eyes and recall an experience that was *absolutely marvellous*, a situation when you felt excited, curious and positive.
- 2. Open your eyes for a second or two and look at page 241 or any other page that contains many formulae.
- 3. Then close your eyes again and return to your marvellous experience.
- 4. Repeat the observation of the formulae and the visualization of your memory steps 2 and 3 three more times.

Then leave the memory, look around yourself to get back into the here and now, and test yourself. Look again at page 241. How do you feel about formulae now?

* *

In the sixteenth century, Niccolò Tartaglia* proposed the following problem. Three young couples want to cross a river. Only a small boat that can carry two people is available. The men are extremely jealous, and would never leave their brides with another man. How many journeys across the river are necessary?

Challenge 27 s



^{*} Niccolò Fontana Tartaglia (1499–1557), important Renaissance mathematician.

* *

Cylinders can be used to roll a flat object over the floor, as shown in Figure 14. The cylinders keep the object plane always at the same distance from the floor. What cross-sections *other* than circular, so-called *curves of constant width*, can a cylinder have to realize the same feat? How many examples can you find? Are objects different than cylinders possible?

SUMMARY ON MOTION

Motion is the most fundamental observation in nature. Everyday motion is predictable and deterministic. Predictability is reflected in six aspects of motion: continuity, conservation, reversibility, mirror-invariance, relativity and minimization. Some of these aspects are valid for all motion, and some are valid only for everyday motion. Which ones, and why? We explore this now.

Challenge 29 d

Challenge 28 s







Challenge 30 s

CHAPTER 2

FROM MOTION MEASUREMENT TO CONTINUITY

Physic ist wahrlich das eigentliche Studium des Menschen.*

Georg Christoph Lichtenberg

hroughout our everyday life: only one thing can be at a given spot at a given time. his general description can be separated into three assumptions: matter is *impenetrable* and *moves*, time is made of *instants*, and space is made of *points*. Without these three assumptions (do you agree with them?) it is not possible to define velocity in everyday life. This description of nature is called *Galilean physics*, or sometimes *Newtonian physics*.

Galileo Galilei (1564–1642), Tuscan professor of mathematics, was a founder of modern physics and is famous for advocating the importance of observations as checks of statements about nature. By requiring and performing these checks throughout his life, he was led to continuously increase the accuracy in the description of motion. For example, Galileo studied motion by measuring change of position with a self-constructed stopwatch. His approach changed the speculative description of ancient Greece into the experimental physics of Renaissance Italy.**

The English alchemist, occultist, theologian, physicist and politician Isaac Newton (1643–1727) was one of the first to pursue with vigour the idea that different types of motion have the same properties, and he made important steps in constructing the concepts necessary to demonstrate this idea.***

The explorations by Galileo and his predecessors provided the first clear statements on the properties of speed, space and time.

Ref. 25

Ref. 26



^{* &#}x27;Physics truly is the proper study of man.' Georg Christoph Lichtenberg (1742–1799) was an important physicist and essayist.

^{**} The best and most informative book on the life of Galileo and his times is by Pietro Redondi (see the section on page 285). Galileo was born in the year the pencil was invented. Before his time, it was impossible to do paper and pencil calculations. For the curious, the www.mpiwg-berlin.mpg.de website allows you to read an original manuscript by Galileo.

^{***} Newton was born a year after Galileo died. Newton's other hobby, as master of the Mint, was to supervise personally the hanging of counterfeiters. About Newton's infatuation with alchemy, see the books by Dobbs. Among others, Newton believed himself to be chosen by god; he took his Latin name, *Isaacus Neuutonus*, and formed the anagram *Jeova sanctus unus*. About Newton and his importance for classical mechanics, see the text by Clifford Truesdell.



FIGURE 15 Galileo Galilei (1564-1642).









FIGURE 16 Some speed measurement devices: an anemometer, a tachymeter for inline skates, a sport radar gun and a Pitot–Prandtl tube in an aeroplane (© Fachhochschule Koblenz, Silva, Tracer, Wikimedia).

WHAT IS VELOCITY?

There is nothing else like it.

Jochen Rindt*

Velocity fascinates. To physicists, not only car races are interesting, but any moving entity is. Therefore they first measure as many examples as possible. A selection of measured speed values is given in Table 3. The units and prefixes used are explained in detail in Appendix B. Some speed measurement devices are shown in Figure 16.

Page 383

Page 77

Everyday life teaches us a lot about motion: objects can overtake each other, and they can move in different directions. We also observe that velocities can be added or changed smoothly. The precise list of these properties, as given in Table 4, is summarized by mathematicians in a special term; they say that velocities form a *Euclidean vector space*.** More details about this strange term will be given shortly. For now we just note that in describing nature, mathematical concepts offer the most accurate vehicle.

When velocity is assumed to be an Euclidean vector, it is called *Galilean* velocity. Velocity is a profound concept. For example, velocity does not need space and time measurements to be defined. Are you able to find a means of measuring velocities without



^{*} Jochen Rindt (1942–1970), famous Austrian Formula One racing car driver, speaking about speed.

^{**} It is named after Euclid, or Eukleides, the great Greek mathematician who lived in Alexandria around 300 BCE. Euclid wrote a monumental treatise of geometry, the Στοιχεῖα or *Elements*, which is one of the milestones of human thought. The text presents the whole knowledge on geometry of that time. For the first time, Euclid introduces two approaches that are now in common use: all statements are deduced from a small number of basic *axioms* and for every statement a *proof* is given. The book, still in print today, has been the reference geometry text for over 2000 years. On the web, it can be found at aleph0.clarku.edu/~djoyce/java/elements/elements.html.

 $2 \cdot 10^9 \,\mathrm{m/s}$

 $7 \cdot 10^{13} \, \text{m/s}$

no limit

OBSERVATION	
Growth of deep sea manganese crust	
Can you find something slower?	

TABLE 3 Some measured velocity values.

OBSERVATION	VELOCITY
Growth of deep sea manganese crust	80 am/s
Can you find something slower?	Challenge 31 s
Stalagmite growth	0.3 pm/s
Lichen growth	down to 7 pm/s
cm Typical motion of continents	$10\mathrm{mm/a}=0.3\mathrm{nm/s}$
Human growth during childhood, hair growth	4 nm/s
Tree growth	up to 30 nm/s
Electron drift in metal wire	1 μm/s
Sperm motion	60 to 160 µm/s
Speed of light at Sun's centre	0.1 mm/s
Ketchup motion	1 mm/s
Slowest speed of light measured in matter on Earth Ref. 27	$0.3 \mathrm{m/s}$
Speed of snowflakes	0.5 m/s to 1.5 m/s
Signal speed in human nerve cells Ref. 28	0.5 m/s to 120 m/s
Wind speed at 1 Beaufort (light air)	below 1.5 m/s
Speed of rain drops, depending on radius	2 m/s to 8 m/s
Fastest swimming fish, sailfish (Istiophorus platypterus)	22 m/s
2006 Speed sailing record over 500 m (by windsurfer Finian Maynard)	25.1 m/s
2008 Speed sailing record over 500 m (by kitesurfer Alex Caizergues)	26.0 m/s
2009 Speed sailing record over 500 m (by trimaran Hydroptère)	26.4 m/s
Fastest running animal, cheetah (Acinonyx jubatus)	30 m/s
Wind speed at 12 Beaufort (hurricane)	above 33 m/s
Speed of air in throat when sneezing	42 m/s
Fastest throw: a cricket ball thrown with baseball technique while running	g 50 m/s
Freely falling human, depending on clothing	50 to 90 m/s
Fastest bird, diving Falco peregrinus	60 m/s
Fastest badminton smash	70 m/s
Average speed of oxygen molecule in air at room temperature	280 m/s
Speed of sound in dry air at sea level and standard temperature	330 m/s
Cracking whip's end	750 m/s
Speed of a rifle bullet	1 km/s
Speed of crack propagation in breaking silicon	5 km/s
Highest macroscopic speed achieved by man – the Voyager satellite	14 km/s
Speed of Earth through universe	370 km/s
Average speed (and peak speed) of lightning tip	600 km/s (50 Mm/s)
Highest macroscopic speed measured in our galaxy Ref. 29	$0.97 \cdot 10^8 \text{m/s}$
Speed of electrons inside a colour TV	$1 \cdot 10^8 \mathrm{m/s}$
Speed of radio messages in space	299 792 458 m/s
Highest ever measured group velocity of light	$10 \cdot 10^8 \mathrm{m/s}$
	2 109 /

Speed of light spot from a light tower when passing over the Moon

Highest proper velocity ever achieved for electrons by man

Highest possible velocity for a light spot or shadow



VELOCITIES PHYSICAL MATHEMATICAL DEFINITION PROPERTY NAME CAN Be distinguished element of set distinguishability Vol. III, page 223 Change gradually continuum real vector space Page 77, Vol. V, page 288 Page 77 Point somewhere direction vector space, dimensionality Be compared measurability metricity Vol. V, page 280 Be added additivity vector space Page 77 Have defined angles direction Euclidean vector space Page 77 Exceed any limit infinity unboundedness Vol. III, page 224

TABLE 4 Properties of everyday – or Galilean – velocity.

Challenge 32 d

measuring space and time? If so, you probably want to skip to the next volume, jumping 2000 years of enquiries. If you cannot do so, consider this: whenever we measure a quantity we assume that everybody is able to do so, and that everybody will get the same result. In other words, we define *measurement* as a comparison with a standard. We thus implicitly assume that such a standard exists, i.e., that an example of a 'perfect' velocity can be found. Historically, the study of motion did not investigate this question first, because for many centuries nobody could find such a standard velocity. You are thus in good company.

Ref. 30

Some researchers have specialized in the study of the lowest velocities found in nature: they are called geologists. Do not miss the opportunity to walk across a landscape while listening to one of them.

How is velocity measured in everyday life? Animals and people estimate their velocity in two ways: by estimating the frequency of their own movements, such as their steps, or by using their eyes, ears, sense of touch or sense of vibration to deduce how their own position changes with respect to the environment. But several animals have additional capabilities: certain snakes can determine speeds with their infrared-sensing organs, others with their magnetic field sensing organs. Still other animals emit sounds that create echoes in order to measure speeds to high precision. The same range of solutions is used by technical devices. Table 5 gives an overview.

Velocity is not always an easy subject. Physicists like to say, provokingly, that what cannot be measured does not exist. Can you measure your own velocity in empty interstellar space?

Challenge 33 s

Velocity is of interest to both engineers and evolution. In general, self-propelled systems are faster the larger they are. As an example, Figure 17 shows how this applies to the cruise speed of flying things. In general, cruise speed scales with the sixth root of the weight, as shown by the trend line drawn in the graph. (Can you find out why?) By the way, similar *allometric scaling* relations hold for many other properties of moving systems, as we will see later on.

Challenge 34 d

Velocity is a profound subject for an additional reason: we will discover that all seven properties of Table 4 are only approximate; *none* is actually correct. Improved experiments will uncover exceptions for every property of Galilean velocity. The failure of the last three properties of Table 4 will lead us to special and general relativity, the failure of








FIGURE 17 How wing load and sea-level cruise speed scales with weight in flying objects, compared with the general trend line (after a graph © Henk Tennekes).



TABLE 5 Speed measurement devices in biological and engineered systems.

M E A S U R E M E N T	Device	Range
Own running speed in insects, mammals and humans	leg beat frequency measured with internal clock	0 to 33 m/s
Own car speed	tachymeter attached to wheels	0 to 150 m/s
Predators and hunters measuring prey speed	vision system	0 to 30 m/s
Police measuring car speed	radar or laser gun	0 to 90 m/s
Bat measuring own and prey speed at night	doppler sonar	0 to 20 m/s
Sliding door measuring speed of approaching people	doppler radar	0 to 3 m/s
Own swimming speed in fish and humans	friction and deformation of skin	0 to 30 m/s
Own swimming speed in dolphins and ships	sonar to sea floor	0 to 20 m/s
Diving speed in fish, animals, divers and submarines	pressure change	0 to 5 m/s
Water predators and fishing boats measuring prey speed	sonar	0 to 20 m/s
Own speed relative to Earth in insects	often none (grasshoppers)	n.a.
Own speed relative to Earth in birds	visual system	0 to 60 m/s
Own speed relative to Earth in aeroplanes or rockets	radio goniometry, radar	0 to 8000 m/s
Own speed relative to air in insects and birds	filiform hair deflection, feather deflection	0 to 60 m/s
Own speed relative to air in aeroplanes	Pitot-Prandtl tube	0 to 340 m/s
Wind speed measurement in meteorological stations	thermal, rotating or ultrasound anemometers	0 to 80 m/s
Swallows measuring prey speed	visual system	0 to 20 m/s
Bats measuring prey speed	sonar	0 to 20 m/s
Macroscopic motion on Earth	Global Positioning System, Galileo, Glonass	0 to 100 m/s
Pilots measuring target speed	radar	0 to 1000 m/s
Motion of stars	optical Doppler effect	0 to 1000 km/s
Motion of star jets	optical Doppler effect	0 to 200 Mm/s

the middle two to quantum theory and the failure of the first two properties to the unified description of nature. But for now, we'll stick with Galilean velocity, and continue with another Galilean concept derived from it: time.






FIGURE 18 A typical path followed by a stone thrown through the air – a parabola – with photographs (blurred and stroboscopic) of a table tennis ball rebounding on a table (centre) and a stroboscopic photograph of a water droplet rebounding on a strongly hydrophobic surface (right, © Andrew Davidhazy, Max Groenendijk).

Without the concepts *place*, *void* and *time*, change cannot be. [...] It is therefore clear [...] that their investigation has to be carried out, by studying each of them separately.

Aristotle* *Physics*, Book III, part 1.

WHAT IS TIME?

Time does not exist in itself, but only through the perceived objects, from which the concepts of past, of present and of future ensue. Lucretius,** De rerum natura, lib. 1, v. 460 ss.

In their first years of life, children spend a lot of time throwing objects around. The term 'object' is a Latin word meaning 'that which has been thrown in front'. Developmental psychology has shown experimentally that from this very experience children extract the concepts of time and space. Adult physicists do the same when studying motion at university.

When we throw a stone through the air, we can define a *sequence* of observations. Figure 18 illustrates how. Our memory and our senses give us this ability. The sense of hearing registers the various sounds during the rise, the fall and the landing of the stone. Our eyes track the location of the stone from one point to the next. All observations have their place in a sequence, with some observations preceding them, some observations



^{*} Aristotle (b. 384/3 Stageira, d. 322 BCE Euboea), important Greek philosopher and scientist, founder of the *Peripatetic school* located at the Lyceum, a gymnasium dedicated to Apollo Lyceus.

^{**} Titus Lucretius Carus (c. 95 to c. 55 BCE), Roman scholar and poet.

TABLE 6 Selected time measurements.

OBSERVATION	Тіме
Shortest measurable time	$10^{-44} \mathrm{s}$
Shortest time ever measured	10 ys
Time for light to cross a typical atom	0.1 to 10 as
Shortest laser light pulse produced so far	200 as
Period of caesium ground state hyperfine transition	108.782 775 707 78 ps
Beat of wings of fruit fly	1 ms
Period of pulsar (rotating neutron star) PSR 1913+16	0.059 029 995 271(2) s
Human 'instant'	20 ms
Shortest lifetime of living being	0.3 d
Average length of day 400 million years ago	79 200 s
Average length of day today	86 400.002(1) s
From birth to your 1000 million seconds anniversary	31.7 a
Age of oldest living tree	4600 a
Use of human language	0.2 Ma
Age of Himalayas	35 to 55 Ma
Age of oldest rocks, found in Isua Belt, Greenland and in Porpoise Cove, Hudson Bay	3.8 Ga
Age of Earth	4.6 Ga
Age of oldest stars	13.7 Ga
Age of most protons in your body	13.7 Ga
Lifetime of tantalum nucleus ^{180m} Ta	10 ¹⁵ a
Lifetime of bismuth ²⁰⁹ Bi nucleus	$1.9(2) \cdot 10^{19}$ a

simultaneous to them, and still others succeeding them. We say that observations are perceived to happen at various *instants* and we call the sequence of all instants *time*.

An observation that is considered the smallest part of a sequence, i.e., not itself a sequence, is called an *event*. Events are central to the definition of time; in particular, starting or stopping a stopwatch are events. (But do events really exist? Keep this question in the back of your head as we move on.)

Sequential phenomena have an additional property known as stretch, extension or duration. Some measured values are given in Table 6.* *Duration* expresses the idea that sequences *take* time. We say that a sequence takes time to express that other sequences can take place in parallel with it.

How exactly is the concept of time, including sequence and duration, deduced from observations? Many people have looked into this question: astronomers, physicists, watchmakers, psychologists and philosophers. All find:

▶ Time is deduced by comparing motions.



Challenge 35 s

^{*} A year is abbreviated a (Latin 'annus').

Aotion Mountain – The Adventure of Physics 🏻 pdf file available free of charge at www.motionmountain.net 🔻 Copyright © Christoph Schiller November 1997–March 20

Ref. 21

This is even the case for children and animals. Beginning at a very young age, they develop the concept of 'time' from the comparison of motions in their surroundings. Grown-ups take as a standard the motion of the Sun and call the resulting type of time *local time*. From the Moon they deduce a *lunar calendar*. If they take a particular village clock on a European island they call it the *universal time coordinate* (UTC), once known as 'Greenwich mean time.'* Astronomers use the movements of the stars and call the result *ephemeris time* (or one of its successors). An observer who uses his personal watch calls the reading his *proper time*; it is often used in the theory of relativity.

Page 386

Challenge 37 s

Not every movement is a good standard for time. In the year 2000, an Earth rotation did not take 86 400 seconds any more, as it did in the year 1900, but 86 400.002 seconds. Can you deduce in which year your birthday will have shifted by a whole day from the time predicted with 86 400 seconds?

All methods for the definition of time are thus based on comparisons of motions. In order to make the concept as precise and as useful as possible, a *standard* reference motion is chosen, and with it a standard sequence and a standard duration is defined. The device that performs this task is called a *clock*. We can thus answer the question of the section title:

▶ Time is what we read from a clock.

Note that all definitions of time used in the various branches of physics are equivalent to this one; no 'deeper' or more fundamental definition is possible.** Note that the word 'moment' is indeed derived from the word 'movement'. Language follows physics in this case. Astonishingly, the definition of time just given is final; it will never be changed, not even at the top of Motion Mountain. This is surprising at first sight, because many books have been written on the nature of time. Instead, they should investigate the nature of motion! But this is the aim of our walk anyhow. We are thus set to discover all the secrets of time as a side result of our adventure. Every clock reminds us that in order to understand time, we need to understand motion.

A *clock* is thus a moving system whose position can be read. Of course, a *precise* clock is a system moving as regularly as possible, with as little outside disturbance as possible. Is there a perfect clock in nature? Do clocks exist at all? We will continue to study these questions throughout this work and eventually reach a surprising conclusion. At this point, however, we state a simple intermediate result: since clocks do exist, somehow there is in nature an intrinsic, natural and *ideal* way to measure time. Can you see it?

Challenge 38 s

Time is not only an aspect of observations, it is also a facet of personal experience. Even in our innermost private life, in our thoughts, feelings and dreams, we experience sequences and durations. Children learn to relate this internal experience of time with external observations, and to make use of the sequential property of events in their actions. Studies of the origin of psychological time show that it coincides – apart from its lack

Challenge 36 s

31 ** The oldest clocks are sundials. The science of making them is called *gnomonics*.



^{*} Official UTC is used to determine the phase of the power grid, phone companies' bit streams and the signal to the GPS system. The latter is used by many navigation systems around the world, especially in ships, aeroplanes and lorries. For more information, see the www.gpsworld.com website. The time-keeping infrastructure is also important for other parts of the modern economy. Can you spot the most important ones?

TABLE 7 Properties of Galilean time.

INSTANTS OF TIME	PHYSICAL	MATHEMATICAL	DEFINITION
	PROPERTY	N A M E	
Can be distinguished	distinguishability	element of set	Vol. III, page 223
Can be put in order	sequence	order	Vol. V, page 288
Define duration	measurability	metricity	Vol. V, page 280
Can have vanishing duration	continuity	denseness, completeness	Vol. V, page 288
Allow durations to be added	additivity	metricity	Vol. V, page 280
Don't harbour surprises	translation invariance	homogeneity	Page 204
Don't end	infinity	unboundedness	Vol. III, page 224
Are equal for all observers	absoluteness	uniqueness	

of accuracy – with clock time.* Every living human necessarily uses in his daily life the concept of time as a combination of sequence and duration; this fact has been checked in numerous investigations. For example, the term 'when' exists in all human languages.

Time is a concept *necessary* to distinguish between observations. In any sequence, we observe that events succeed each other smoothly, apparently without end. In this context, 'smoothly' means that observations that are not too distant tend to be not too different. Yet between two instants, as close as we can observe them, there is always room for other events. Durations, or *time intervals*, measured by different people with different clocks agree in everyday life; moreover, all observers agree on the order of a sequence of events. Time is thus unique in everyday life.

The properties of everyday time, also listed in Table 7, correspond to the precise version of our everyday experience of time. This concept is called *Galilean time*; all its properties can be expressed simultaneously by describing time with the help of real numbers. In fact, real numbers have been constructed by mathematicians to have exactly the same properties as Galilean time, as explained in the chapter on the brain. Every instant of time can be described by a real number, often abbreviated t, and the duration of a sequence of events is given by the difference between the values for the final and the starting event.

When Galileo studied motion in the seventeenth century, there were as yet no stopwatches. He thus had to build one himself, in order to measure times in the range between a fraction and a few seconds. Can you imagine how he did it?

We will have quite some fun with Galilean time in this part of our adventure. However, hundreds of years of close scrutiny have shown that *every single* property of time just listed is approximate, and none is strictly correct. This story is told in the rest of our adventure.

Vol. V, page 31

Ref. 32

* The brain contains numerous clocks. The most precise clock for short time intervals, the internal interval timer of the brain, is more accurate than often imagined, especially when trained. For time periods between a few tenths of a second, as necessary for music, and a few minutes, humans can achieve timing accuracies of a few per cent.

Vol. III, page 232

Challenge 39 s












FIGURE 19 Different types of clocks: a high-tech sundial (size c. 30 cm), a naval pocket chronometer (size c. 6 cm), a caesium atomic clock (size c. 4 m), a group of cyanobacteria and the Galilean satellites of Jupiter (© Carlo Heller at www.heliosuhren.de, Anonymous, INMS, Wikimedia, NASA).



CLOCKS

A *clock* is a moving system whose position can be read. There are many types of clocks: stopwatches, twelve-hour clocks, sundials, lunar clocks, seasonal clocks, etc. A few are shown in Figure 19. Almost all clock types are also found in plants and animals, as shown in Table 8.

Ref. 35 Interestingly, there is a strict rule in the animal kingdom: large clocks go slow. How this happens, is shown in Figure 20, another example of an *allometric scaling 'law'*.

Clock makers are experts in producing motion that is as regular as possible. We will discover some of their tricks below. We will also explore, later on, the fundamental limits for the precision of clocks.

Ref. 34



TABLE 8 Examples of biological rhythms and clocks.

LIVING BEING	OSCILLATING SYSTEM	Period
Sand hopper (Talitrus saltator)	knows in which direction to flee from the position of the Sun or Moon	circadian
Human (Homo sapiens)	gamma waves in the brain	0.023 to 0.03 s
	alpha waves in the brain	0.08 to 0.13 s
	heart beat	0.3 to 1.5 s
	delta waves in the brain	0.3 to 10 s
	blood circulation	30 s
	cellular circhoral rhythms	1 to 2 ks
	rapid-eye-movement sleep period	5.4 ks
	nasal cycle	4 to 9 ks
	growth hormone cycle	11 ks
	suprachiasmatic nuclei (SCN), circadian hormone concentration, temperature, etc.; leads to jet lag	90 ks
	monthly period	2.4(4) Ms
	built-in aging	3.2(3) Gs
Common fly (Musca domestica)	wing beat	30 ms
Fruit fly (Drosophila melanogaster)	wing beat for courting	34 ms
Most insects (e.g. wasps, fruit flies)	winter approach detection (diapause) by length of day measurement; triggers metabolism changes	yearly
Algae (Acetabularia)	Adenosinetriphosphate (ATP) concentration	
Moulds (e.g. Neurospora crassa)	conidia formation	circadian
Many flowering plants	flower opening and closing	circadian
Tobacco plant	flower opening clock; triggered by length of days, discovered in 1920 by Garner and Allard	annual
Arabidopsis	circumnutation	circadian
	growth	a few hours
Telegraph plant (<i>Desmodium gyrans</i>)	side leaf rotation	200 s
Forsythia europaea, F. suspensa, F. viridissima, F. spectabilis	Flower petal oscillation, discovered by Van Gooch in 2002	5.1 ks






FIGURE 20 How biological rhythms scale with size in mammals: all scale more or less with a quater poweer of the mass (after data from the EMBO and Enrique Morgado).



WHY DO CLOCKS GO CLOCKWISE?

Challenge 40 s

What time is it at the North Pole now?

))

Most rotational motions in our society, such as athletic races, horse, bicycle or ice skating races, turn anticlockwise.* Mathematicians call this the positive rotation sense. Every supermarket leads its guests anticlockwise through the hall. Why? Most people are right-handed, and the right hand has more freedom at the outside of a circle. Therefore thousands of years ago chariot races in stadia went anticlockwise. As a result, all stadium races still do so to this day, and that is why runners move anticlockwise. For the same reason, helical stairs in castles are built in such a way that defending right-handers, usually from above, have that hand on the outside.

On the other hand, the clock imitates the shadow of sundials; obviously, this is true on the northern hemisphere only, and only for sundials on the ground, which were the most common ones. (The old trick to determine south by pointing the hour hand of a horizontal watch to the Sun and halving the angle between it and the direction of 12 o'clock does not work on the southern hemisphere.) So every clock implicitly continues to state on which hemisphere it was invented. In addition, it also tells us that sundials on walls came in use much later than those on the floor.

Does time flow?

Wir können keinen Vorgang mit dem 'Ablauf der Zeit' vergleichen – diesen gibt es nicht –, sondern nur mit einem anderen Vorgang (etwa dem Gang des Chronometers).**

Ludwig Wittgenstein, Tractatus, 6.3611

Si le temps est un fleuve, quel est son lit?***

))

The expression 'the flow of time' is often used to convey that in nature change follows after change, in a steady and continuous manner. But though the hands of a clock 'flow', time itself does not. Time is a concept introduced specially to describe the flow of events around us; it does not itself flow, it *describes* flow. Time does not advance. Time is neither linear nor cyclic. The idea that time flows is as hindering to understanding nature as is the idea that mirrors exchange right and left.

Vol. III, page 81

Ref. 3

Challenge 41 e

The misleading use of the expression 'flow of time', propagated first by some flawed Greek thinkers and then again by Newton, continues. Aristotle, careful to think logically, pointed out its misconception, and many did so after him. Nevertheless, expressions such as 'time reversal', the 'irreversibility of time', and the much-abused 'time's arrow' are still common. Just read a popular science magazine chosen at random. The fact is: time cannot be reversed, only motion can, or more precisely, only velocities of objects; time has



^{*} Notable exceptions are most, but not all, Formula 1 races.

^{** &#}x27;We cannot compare any process with 'the passage of time' – there is no such thing – but only with another process (say, with the working of a chronometer).'

^{*** &#}x27;If time is a river, what is his bed?'

Ref. 37

no arrow, only motion has; it is not the flow of time that humans are unable to stop, but the motion of all the objects in nature. Incredibly, there are even books written by respected physicists that study different types of 'time's arrows' and compare them with each other. Predictably, no tangible or new result is extracted. Time does *not* flow.

In the same manner, colloquial expressions such as 'the start (or end) of time' should be avoided. A motion expert translates them straight away into 'the start (or end) of motion'.

WHAT IS SPACE?

The introduction of numbers as coordinates [...] is an act of violence [...].

Hermann Weyl, Philosophie der Mathematik

und Naturwissenschaft.*

Whenever we distinguish two objects from each other, such as two stars, we first of all distinguish their positions. We distinguish positions with our senses of sight, touch, hearing and proprioperception. Position is therefore an important aspect of the physical state of an object. A position is taken by only one object at a time. Positions are limited. The set of all available positions, called *(physical) space*, acts as both a container and a background.

Closely related to space and position is *size*, the set of positions an object occupies. Small objects occupy only subsets of the positions occupied by large ones. We will discuss size in more detail shortly.

Page 53

How do we deduce space from observations? During childhood, humans (and most higher animals) learn to bring together the various *perceptions* of space, namely the visual, the tactile, the auditory, the kinaesthetic, the vestibular etc., into one coherent set of experiences and description. The result of this learning process is a certain concept of space in the brain. Indeed, the question 'where?' can be asked and answered in all languages of the world. Being more precise, adults derive space from distance measurements. The concepts of length, area, volume, angle and solid angle are all deduced with their help. Geometers, surveyors, architects, astronomers, carpet salesmen and producers of metre sticks base their trade on distance measurements. Space is a concept formed to summarize all the distance relations between objects for a precise description of observations.

Challenge 42 s

Metre sticks work well only if they are straight. But when humans lived in the jungle, there were no straight objects around them. No straight rulers, no straight tools, nothing. Today, a cityscape is essentially a collection of straight lines. Can you describe how humans achieved this?


Once humans came out of the jungle with their newly built metre sticks, they collected a wealth of results. The main ones are listed in Table 9; they are easily confirmed by personal experience. Objects can take positions in an apparently *continuous* manner: there indeed are more positions than can be counted.** Size is captured by defining the



Motion Mountain – The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christoph Schiller November 199

^{*} Hermann Weyl (1885–1955) was one of the most important mathematicians of his time, as well as an important theoretical physicist. He was one of the last universalists in both fields, a contributor to quantum theory and relativity, father of the term 'gauge' theory, and author of many popular texts.

^{**} For a definition of uncountability, see page 226 in Volume III.






FIGURE 21 Two proofs that space has three dimensions: the vestibular labyrinth in the inner ear of mammals (here a human) with three canals and a knot (© Northwestern University).

distance between various positions, called *length*, or by using the field of view an object takes when touched, called its *surface*. Length and surface can be measured with the help of a metre stick. (Selected measurement results are given in Table 10; some length measurement devices are shon in Figure 23.) The length of objects is independent of the person measuring it, of the position of the objects and of their orientation. In daily life the sum of angles in any triangle is equal to two right angles. There are no limits to distances, lengths and thus to space.

Experience shows us that space has three dimensions; we can define sequences of positions in precisely three independent ways. Indeed, the inner ear of (practically) all vertebrates has three semicircular canals that sense the body's acceleration in the three dimensions of space, as shown in Figure 21.* Similarly, each human eye is moved by three pairs of muscles. (Why three?) Another proof that space has three dimensions is provided by shoelaces: if space had more than three dimensions, shoelaces would not be useful, because knots exist only in three-dimensional space. But why does space have three dimensions? This is one of the most difficult question of physics; it will be answered only in the very last part of our walk.

It is often said that thinking in four dimensions is impossible. That is wrong. Just try. For example, can you confirm that in four dimensions knots are impossible?

Like time intervals, length intervals can be described most precisely with the help of *real numbers*. In order to simplify communication, standard *units* are used, so that everybody uses the same numbers for the same length. Units allow us to explore the general properties of *Galilean space* experimentally: space, the container of objects, is continuous, three-dimensional, isotropic, homogeneous, infinite, Euclidean and unique or 'absolute'. In mathematics, a structure or mathematical concept with all the properties just mentioned is called a three-dimensional *Euclidean space*. Its elements, (*mathematical*) *points*, are described by three real parameters. They are usually written as

$$(x, y, z) \tag{1}$$







^{*} Note that saying that space has three dimensions *implies* that space is continuous; the Dutch mathematician and philosopher Luitzen Brouwer (b. 1881 Overschie, d. 1966 Blaricum) showed that dimensionality is only a useful concept for continuous sets.

TABLE 9 Properties of Galilean space.

POINTS	PHYSICAL PROPERTY	M ATHEMATICAL NAME	DEFINI-
Can be distinguished	distinguishability	element of set	Vol. III, page 223
Can be lined up if on one line	sequence	order	Vol. V, page 288
Can form shapes	shape	topology	Vol. V, page 287
Lie along three independent directions	possibility of knots	3-dimensionality	Page 77 , Vol. V, page 280
Can have vanishing distance	continuity	denseness, completeness	Vol. V, page 288
Define distances	measurability	metricity	Vol. V, page 280
Allow adding translations	additivity	metricity	Vol. V, page 280
Define angles	scalar product	Euclidean space	Page 77
Don't harbour surprises	translation invariance	homogeneity	
Can beat any limit	infinity	unboundedness	Vol. III, page 224
Defined for all observers	absoluteness	uniqueness	Page 52



FIGURE 22 René Descartes (1596–1650).

and are called *coordinates*. They specify and order the location of a point in space. (For the precise definition of Euclidean spaces, see below..)

What is described here in just half a page actually took 2000 years to be worked out, mainly because the concepts of 'real number' and 'coordinate' had to be discovered first. The first person to describe points of space in this way was the famous mathematician and philosopher René Descartes*, after whom the coordinates of expression (1) are named *Cartesian*.

Like time, space is a *necessary* concept to describe the world. Indeed, space is automatically introduced when we describe situations with many objects. For example, when many spheres lie on a billiard table, we cannot avoid using space to describe the relations between them. There is no way to avoid using spatial concepts when talking about nature.

Even though we need space to talk about nature, it is still interesting to ask why this is possible. For example, since many length measurement methods do exist – some are



^{*} René Descartes or Cartesius (b. 1596 La Haye, d. 1650 Stockholm), French mathematician and philosopher, author of the famous statement 'je pense, donc je suis', which he translated into 'cogito ergo sum' – I think therefore I am. In his view this is the only statement one can be sure of.

TABLE 10 Some measured distance values.

OBSERVATION	DISTANCE
Galaxy Compton wavelength	10 ⁻⁸⁵ m (calculated only)
Planck length, the shortest measurable length	$10^{-35} \mathrm{m}$
Proton diameter	1 fm
Electron Compton wavelength	2.426 310 215(18) pm
Smallest air oscillation detectable by human ear	11 pm
Hydrogen atom size	30 pm
Size of small bacterium	0.2 μm
Wavelength of visible light	0.4 to 0.8 μm
Point: diameter of smallest object visible with naked eye	20 μm
Diameter of human hair (thin to thick)	30 to 80 μm
Total length of DNA in each human cell	2 m
Largest living thing, the fungus Armillaria ostoyae	3 km
Longest human throw with any object, using a boomerang	427 m
Highest human-built structure, Burj Khalifa	828 m
Largest spider webs in Mexico	<i>c</i> . 5 km
Length of Earth's Equator	40 075 014.8(6) m
Total length of human blood vessels (rough estimate)	$4to16 \cdot 10^4 \mathrm{km}$
Total length of human nerve cells (rough estimate)	$1.5to8 \cdot 10^5 \mathrm{km}$
Average distance to Sun	149 597 870 691(30) m
Light year	9.5 Pm
Distance to typical star at night	10 Em
Size of galaxy	1 Zm
Distance to Andromeda galaxy	28 Zm
Most distant visible object	125 Ym

Challenge 45 s

listed in Table 11 – and since they all yield consistent results, there must be a *natural* or *ideal* way to measure distances, sizes and straightness. Can you find it?

As in the case of time, each of the properties of space just listed has to be checked. And again, careful observations will show that each property is an approximation. In simpler and more drastic words, *all* of them are wrong. This confirms Weyl's statement at the beginning of this section. In fact, his statement about the violence connected with the introduction of numbers is told by every forest in the world, and of course also by the one at the foot of Motion Mountain. The rest of our adventure will show this.

Μέτρον ἄριστον.*

Cleobulus



^{* &#}x27;Measure is the best (thing).' Cleobulus (Κλεοβουλος) of Lindos, (c. 620–550 BCE) was another of the proverbial seven sages.





ARE SPACE AND TIME ABSOLUTE OR RELATIVE?

In everyday life, the concepts of Galilean space and time include two opposing aspects; the contrast has coloured every discussion for several centuries. On the one hand, space and time express something invariant and permanent; they both act like big containers for all the objects and events found in nature. Seen this way, space and time have an existence of their own. In this sense one can say that they are fundamental or *absolute*. On the other hand, space and time are tools of description that allow us to talk about rela-



TABLE 11 Length measurement devices in biological and engineered systems.

M e a s u r e m e n t	DEVICE	RANGE
Humans		
Measurement of body shape, e.g. finger distance, eye position, teeth distance	muscle sensors	0.3 mm to 2 m
Measurement of object distance	stereoscopic vision	1 to 100 m
Measurement of object distance	sound echo effect	0.1 to 1000 m
Animals		
Measurement of hole size	moustache	up to 0.5 m
Measurement of walking distance by desert ants	step counter	up to 100 m
Measurement of flight distance by honey bees	eye	up to 3 km
Measurement of swimming distance by sharks	magnetic field map	up to 1000 km
Measurement of prey distance by snakes	infrared sensor	up to 2 m
Measurement of prey distance by bats, dolphins, and hump whales	sonar	up to 100 m
Measurement of prey distance by raptors	vision	0.1 to 1000 m
Machines		
Measurement of object distance by laser	light reflection	0.1 m to 400 Mm
Measurement of object distance by radar	radio echo	0.1 to 50 km
Measurement of object length	interferometer	$0.5\mu m$ to $50km$
Measurement of star, galaxy or quasar distance	intensity decay	up to 125 Ym
Measurement of particle size	accelerator	down to 10^{-18} m

Challenge 46 e

Ref. 38

tions between objects. In this view, they do not have any meaning when separated from objects, and only result from the relations between objects; they are derived, relational or *relative*. Which of these viewpoints do you prefer? The results of physics have alternately favoured one viewpoint or the other. We will repeat this alternation throughout our adventure, until we find the solution. And obviously, it will turn out to be a third option.

SIZE - WHY AREA EXISTS, BUT VOLUME DOES NOT

A central aspect of objects is their size. As a small child, under school age, every human learns how to use the properties of size and space in their actions. As adults seeking precision, with the definition of *distance* as the difference between coordinates allows us to define *length* in a reliable way. It took hundreds of years to discover that this is *not* the case. Several investigations in physics and mathematics led to complications.

The physical issues started with an astonishingly simple question asked by Lewis Richardson:* How long is the western coastline of Britain?

Following the coastline on a map using an odometer, a device shown in Figure 24, Richardson found that the length l of the coastline depends on the scale s (say 1:10 000



^{*} Lewis Fray Richardson (1881–1953), English physicist and psychologist.





FIGURE 24 A curvimeter or odometer (photograph © Frank Müller).




FIGURE 25 A fractal: a self-similar curve of infinite length (far right), and its construction.

$$l = l_0 \ s^{0.25} \tag{2}$$

(Richardson found other exponentials for other coasts.) The number l_0 is the length at scale 1:1. The main result is that the larger the map, the longer the coastline. What would happen if the scale of the map were increased even beyond the size of the original? The length would increase beyond all bounds. Can a coastline really have *infinite* length? Yes, it can. In fact, mathematicians have described many such curves; they are called *fractals*. An infinite number of them exist, and Figure 25 shows one example.* Can you construct another?

Challenge 47 e

Length has other strange properties. The Italian mathematician Giuseppe Vitali was the first to discover that it is possible to cut a line segment of length 1 into pieces that can be reassembled – merely by shifting them in the direction of the segment – into a line segment of length 2. Are you able to find such a division using the hint that it is only possible using infinitely many pieces?

Challenge 48 d

To sum up, length is well defined for lines that are straight or nicely curved, but not for intricate lines, or for lines made of infinitely many pieces. We therefore avoid fractals and other strangely shaped curves in the following, and we take special care when we talk about infinitely small segments. These are the central assumptions in the first five volumes of this adventure, and we should never forget them. We will come back to these assumptions in the last volume of our adventure.

In fact, all these problems pale when compared with the following problem. Commonly, area and volume are defined using length. You think that it is easy? You're wrong, as well as being a victim of prejudices spread by schools around the world. To define area

^{*} Most of these curves are *self-similar*, i.e., they follow scaling 'laws' similar to the above-mentioned. The term 'fractal' is due to the Polish mathematician Benoît Mandelbrot and refers to a strange property: in a certain sense, they have a non-integral number D of dimensions, despite being one-dimensional by construction. Mandelbrot saw that the non-integer dimension was related to the exponent e of Richardson by D = 1 + e, thus giving D = 1.25 in the example above.






FIGURE 26 A polyhedron with one of its dihedral angles (© Luca Gastaldi).

and volume with precision, their definitions must have two properties: the values must be *additive*, i.e., for finite and infinite sets of objects, the total area and volume have to be the sum of the areas and volumes of each element of the set; and they must be *rigid*, i.e., if one cuts an area or a volume into pieces and then rearranges the pieces, the value remains the same. Are such definitions possible? In other words, do such concepts of volume and area exist?

For areas in a plane, one proceeds in the following standard way: one defines the area A of a rectangle of sides a and b as A = ab; since any polygon can be rearranged into a rectangle with a finite number of straight cuts, one can then define an area value for all polygons. Subsequently, one can define area for nicely curved shapes as the limit of the sum of infinitely many polygons. This method is called *integration*; it is introduced in detail in the section on physical action.

Page 214

Challenge 49 s

Challenge 50 s

However, integration does not allow us to define area for arbitrarily bounded regions. (Can you imagine such a region?) For a complete definition, more sophisticated tools are needed. They were discovered in 1923 by the famous mathematician Stefan Banach.* He proved that one can indeed define an area for any set of points whatsoever, even if the border is not nicely curved but extremely complicated, such as the fractal curve previously mentioned. Today this generalized concept of area, technically a 'finitely additive isometrically invariant measure,' is called a *Banach measure* in his honour. Mathematicians sum up this discussion by saying that since in two dimensions there is a Banach measure, there is a way to define the concept of area – an additive and rigid measure – for any set of points whatsoever.**



^{*} Stefan Banach (b. 1892 Krakow, d. 1945 Lvov), important Polish mathematician.

^{**} Actually, this is true only for sets on the plane. For curved surfaces, such as the surface of a sphere, there are complications that will not be discussed here. In addition, the problems mentioned in the definition of length of fractals also reappear for area if the surface to be measured is not flat. A typical example is the area of the human lung: depending on the level of details examined, one finds area values from a few up to over a hundred square metres.

What is the situation in *three* dimensions, i.e., for volume? We can start in the same way as for area, by defining the volume V of a rectangular polyhedron with sides a, b, c as V=abc. But then we encounter a first problem: a general polyhedron cannot be cut into a cube by straight cuts! The limitation was discovered in 1900 and 1902 by Max Dehn.* He found that the possibility depends on the values of the edge angles, or dihedral angles, as the mathematicians call them. (They are defined in Figure 26.) If one ascribes to every edge of a general polyhedron a number given by its length l times a special function $g(\alpha)$ of its dihedral angle α , then Dehn found that the sum of all the numbers for all the edges of a solid does not change under dissection, provided that the function fulfils $g(\alpha + \beta) = g(\alpha) + g(\beta)$ and $g(\pi) = 0$. An example of such a strange function g is the one assigning the value 0 to any rational multiple of π and the value 1 to a basis set of irrational multiples of π . The values for all other dihedral angles of the polyhedron can then be constructed by combination of rational multiples of these basis angles. Using this function, you may then deduce for yourself that a cube cannot be dissected into a regular tetrahedron because their respective Dehn invariants are different.**

Challenge 51 s

Ref. 40

Despite the problems with Dehn invariants, one *can* define a rigid and additive concept of volume for polyhedra, since for all polyhedra and, in general, for all 'nicely curved' shapes, one can again use integration for the definition of their volume.

Now let us consider general shapes and general cuts in three dimensions, not just the 'nice' ones mentioned so far. We then stumble on the famous *Banach–Tarski theorem* (or paradox). In 1924, Stefan Banach and Alfred Tarski*** proved that it is possible to cut one sphere into five pieces that can be recombined to give *two* spheres, each the size of the original. This counter-intuitive result is the Banach–Tarski theorem. Even worse, another version of the theorem states: take any two sets not extending to infinity and containing a solid sphere each; then it is always possible to dissect one into the other with a *finite* number of cuts. In particular it is possible to dissect a pea into the Earth, or vice versa. Size does not count!**** Volume is thus not a useful concept at all.

Challenge 52 s

The Banach–Tarski theorem raises two questions: first, can the result be applied to gold or bread? That would solve many problems. Second, can it be applied to empty space? In other words, are matter and empty space continuous? Both topics will be explored later in our walk; each issue will have its own, special consequences. For the moment, we eliminate this troubling issue by restricting our interest to smoothly curved shapes (and cutting knives). With this restriction, volumes of matter and of empty space do behave nicely: they are additive and rigid, and show no paradoxes.***** Indeed, the cuts required for the Banach–Tarski paradox are not smooth; it is not possible to perform

***** Mathematicians say that a so-called *Lebesgue measure* is sufficient in physics. This countably additive isometrically invariant measure provides the most general way to define a volume.



Vol. III, page 224 Ref. 41

^{*} Max Dehn (b. 1878 Hamburg, d. 1952 Black Mountain), German mathematician, student of David Hilbert.
** This is also told in the beautiful book by M. AIGLER & G. M. ZIEGLER, *Proofs from the Book*, Springer Verlag, 1999. The title is due to the famous habit of the great mathematician Paul Erdős to imagine that all beautiful mathematical proofs can be assembled in the 'book of proofs'.

^{***} Alfred Tarski (b. 1902 Warsaw, d. 1983 Berkeley), Polish mathematician.

^{****} The proof of the result does not need much mathematics; it is explained beautifully by Ian Stewart in *Paradox of the spheres*, New Scientist, 14 January 1995, pp. 28–31. The proof is based on the axiom of choice, which is presented later on. The Banach–Tarski paradox also exists in four dimensions, as it does in any higher dimension. More mathematical detail can be found in the beautiful book by Stan Wagon.





FIGURE 27 Straight lines found in nature: cerussite (picture width approx. 3 mm, © Stephan Wolfsried) and selenite (picture width approx. 15 m, © Arch. Speleoresearch & Films/La Venta at www.laventa.it and www.naica.com.mx).

them with an everyday knife, as they require (infinitely many) infinitely sharp bends performed with an infinitely sharp knife. Such a knife does not exist. Nevertheless, we keep in the back of our mind that the size of an object or of a piece of empty space is a tricky quantity - and that we need to be careful whenever we talk about it.

WHAT IS STRAIGHT?

When you see a solid object with a straight edge, it is a 99 %-safe bet that it is manmade. Of course, there are exceptions, as shown in Figure 27.* The largest crystals ever found are 18 m in length. But in general, the contrast between the objects seen in a city – buildings, furniture, cars, electricity poles, boxes, books - and the objects seen in a forest - trees, plants, stones, clouds - is evident: in the forest nothing is straight or flat, in the city most objects are. How is it possible for humans to produce straight objects while there are almost none to be found in nature?

Any forest teaches us the origin of straightness; it presents tall tree trunks and rays of daylight entering from above through the leaves. For this reason we call a line straight if it touches either a plumb-line or a light ray along its whole length. In fact, the two definitions are equivalent. Can you confirm this? Can you find another definition? Obviously, we call a surface flat if for any chosen orientation and position the surface touches a plumb-line or a light ray along its whole extension.

In summary, the concept of straightness – and thus also of flatness – is defined with the help of bodies or radiation. In fact, all spatial concepts, like all temporal concepts, require motion for their definition.

A HOLLOW EARTH?

Space and straightness pose subtle challenges. Some strange people maintain that all humans live on the *inside* of a sphere; they (usually) call this the *hollow Earth theory*. They claim that the Moon, the Sun and the stars are all near the centre of the hollow sphere, as illustrated in Figure 29. They also explain that light follows curved paths in the sky and

* Another famous exception, unrelated to atomic structures, is the well-known Irish geological formation called the Giant's Causeway. Other candidates that might come to mind, such as certain bacteria which have (almost) square or (almost) triangular shapes are not counter-examples, as the shapes are only approximate.

Ref 44

Ref. 43

Page 407

Challenge 53 s



Page 362

Ref. 42



FIGURE 28 A photograph of the Earth – seen from the direction of the Sun (NASA).

Challenge 54 s

Challenge 55 e

Vol. II, page 263

that when conventional physicists talk about a distance r from the centre of the Earth, the real hollow Earth distance is $r_{\rm he} = R_{\rm Earth}^2/r$. Can you show that this model is wrong? Roman Sexl* used to ask this question to his students and fellow physicists. The answer is simple: if you think you have an argument to show that this view is wrong, you are mistaken! There is *no way* of showing that such a view is wrong. It is possible to explain the horizon, the appearance of day and night, as well as the satellite photographs of the round Earth, such as Figure 28. To explain what happened during a flight to the Moon is also fun. A coherent hollow Earth view is fully *equivalent* to the usual picture of an infinitely extended space. We will come back to this problem in the section on general relativity.

CURIOSITIES AND FUN CHALLENGES ABOUT EVERYDAY SPACE AND TIME

Challenge 56 s

How does one measure the speed of a gun bullet with a stop watch, in a space of 1 m³, without electronics? Hint: the same method can also be used to measure the speed of light.

* *

Challenge 57 s What is faster: an arrow or a motorbike?



^{*} Roman Sexl, (1939–1986), important Austrian physicist, author of several influential textbooks on gravitation and relativity.



FIGURE 29 A model illustrating the hollow Earth theory, showing how day and night appear (© Helmut Diehl).

Time measurements require periodic phenomena. Tree rings are traces of the seasons. Glaciers also have such traces, the *ogives*. Similar traces are found in teeth. Do you know more examples?

* *

A man wants to know how many stairs he would have to climb if the escalator in front of him, which is running upwards, were standing still. He walks up the escalator and counts 60 stairs; walking down the same escalator with the same speed he counts 90 stairs. What is the answer?

Challenge 58 s

* *

You have two hourglasses: one needs 4 minutes and one needs 3 minutes. How can you use them to determine when 5 minutes are over?

* *

You have two water containers: one of 3 litres, another of 5 litres. How can you use them Challenge 60 e to bring 4 litres of water?

* *

Challenge 61 s How can you make a hole in a postcard that allows you to step through it?





FIGURE 30 At what height is a conical glass half full?




FIGURE 31 Can the snail reach the horse once it starts galloping away?

What fraction of the height of a conical glass, shown in Figure 30, must be filled to make the glass half full?

* *

Challenge 63 s How many pencils are needed to draw a line as long as the equator of the Earth?

* *

Everybody knows the puzzle about the bear: A hunter leaves his home, walks 10 km to the South and 10 km to the West, shoots a bear, walks 10 km to the North, and is back home. What colour is the bear? You probably know the answer straight away. Now comes the harder question, useful for winning money in bets. The house could be on several *additional* spots on the Earth; where are these less obvious spots from which a man can have *exactly* the same trip (forget the bear now) that was just described and be at home again?

Challenge 64 s

Challenge 62 s

* *

Imagine a rubber band that is attached to a wall on one end and is attached to a horse at the other end, as shown in Figure 31. On the rubber band, near the wall, there is a snail. Both the snail and the horse start moving, with typical speeds – with the rubber being infinitely stretchable. Can the snail reach the horse?

Challenge 65 s



For a mathematician, 1 km is the same as $1000 \, \text{m}$. For a physicist the two are different! Indeed, for a physicist, 1 km is a measurement lying between $0.5 \, \text{km}$ and $1.5 \, \text{km}$, whereas $1000 \, \text{m}$ is a measurement between $999.5 \, \text{m}$ and $1000.5 \, \text{m}$. So be careful when you write down measurement values. The professional way is to write, for example, $1000(8) \, \text{m}$ to mean $1000 \pm 8 \, \text{m}$, i.e., a value that lies between $992 \, \text{and} \, 1008 \, \text{m}$ with a probability of 68.3%.

Page 388

* *

Imagine a black spot on a white surface. What is the colour of the line separating the spot from the background? This question is often called Peirce's puzzle.

* *

Also bread is an (approximate) fractal, though an irregular one. The fractal dimension Challenge 67 s of bread is around 2.7. Try to measure it!

* *

Challenge 68 e How do you find the centre of a beer mat using paper and pencil?

* *

How often in 24 hours do the hour and minute hands of a clock lie on top of each other?

Challenge 69 s

For clocks that also have a second hand, how often do all three hands lie on top of each other?

* *

How many times in twelve hours can the two hands of a clock be *exchanged* with the result that the new situation shows a *valid* time? What happens for clocks that also have a third hand for seconds?

* *

Challenge 71 s How many minutes does the Earth rotate in one minute?

* *

What is the highest speed achieved by throwing (with and without a racket)? What was Challenge 72 s the projectile used?

* *

A rope is put around the Earth, on the Equator, as tightly as possible. The rope is then lengthened by 1 m. Can a mouse slip under the rope? The original, tight rope is lengthened by 1 mm. Can a child slip under the rope?

* *

Jack was rowing his boat on a river. When he was under a bridge, he dropped a ball into the river. Jack continued to row in the same direction for 10 minutes after he dropped the ball. He then turned around and rowed back. When he reached the ball, the ball had

Challenge 74 s floated 600 m from the bridge. How fast was the river flowing?






FIGURE 32 A 9-to-10 vernier/nonius/clavius and a 19-to-20 version (in fact, a 38-to-40 version) in a caliper (© www.medien-werkstatt.de).

Adam and Bert are brothers. Adam is 18 years old. Bert is twice as old as at the time when Challenge 75 e Adam was the age that Bert is now. How old is Bert?

* *

'Where am I?' is a common question; 'When am I?' is never asked, not even in other Challenge 76 s languages. Why?

* *

Challenge 77 s Is there a smallest time interval in nature? A smallest distance?

* *

Given that you know what straightness is, how would you characterize or define the cur-Challenge 78 s vature of a curved line using numbers? And that of a surface?

* *

Challenge 79 s What is the speed of your eyelid?

Challenge 80 s

* *

The surface area of the human body is about 400 m². Can you say where this large number comes from?

* *

How does a *vernier* work? It is called *nonius* in other languages. The first name is derived from a French military engineer* who did not invent it, the second is a play of words on the Latinized name of the Portuguese inventor of a more elaborate device** and the Latin word for 'nine'. In fact, the device as we know it today – shown in Figure 32 – was designed around 1600 by Christophonius Clavius,*** the same astronomer whose



^{*} Pierre Vernier (1580–1637), French military officer interested in cartography.

 $^{^{**}}$ Pedro Nuñes or Peter Nonnius (1502–1578), Portuguese mathematician and cartographer.

^{***} Christophonius Clavius or Schlüssel (1537–1612), Bavarian astronomer, one of the main astronomers of his time.




FIGURE 33 Leaving a parking space.

studies were the basis of the Gregorian calendar reform of 1582. Are you able to design a vernier/nonius/clavius that, instead of increasing the precision tenfold, does so by an arbitrary factor? Is there a limit to the attainable precision?

Challenge 81 s

Fractals in three dimensions bear many surprises. Let us generalize Figure 25 to three dimensions. Take a regular tetrahedron; then glue on every one of its triangular faces a smaller regular tetrahedron, so that the surface of the body is again made up of many equal regular triangles. Repeat the process, gluing still smaller tetrahedrons to these new (more numerous) triangular surfaces. What is the shape of the final fractal, after an infinite number of steps?

Challenge 82 s

Challenge 83 s

Challenge 84 s

Challenge 85 s

Motoring poses many mathematical problems. A central one is the following parking issue: what is the shortest distance d from the car in front necessary to leave a parking spot without using reverse gear? (Assume that you know the geometry of your car, as shown in Figure 33, and its smallest outer turning radius R, which is known for every car.) Next question: what is the smallest gap required when you are allowed to manoeuvre back and forward as often as you like? Now a problem to which no solution seems to be available in the literature: How does the gap depend on the number, n, of times you use reverse gear? (The author had offered 50 euro for the first well-explained solution; the winning solution by Daniel Hawkins is now found in the appendix.)

Scientists use a special way to write large and small numbers, explained in Table 12.

In 1996 the smallest experimentally probed distance was 10⁻¹⁹ m, achieved between quarks at Fermilab. (To savour the distance value, write it down without the exponent.) What does this measurement mean for the continuity of space? Challenge 86 s

Zeno, the Greek philosopher, discussed in detail what happens to a moving object at a given instant of time. To discuss with him, you decide to build the fastest possible shutter



TABLE 12 The exponential notation: how to write small and large numbers.

Nимвек	EXPONENTIAL NOTATION	Number	Exponential notation
1	10 ⁰		
0.1	10^{-1}	10	10^1
0.2	$2\cdot 10^{-1}$	20	$2\cdot 10^1$
0.0324	$3.24 \cdot 10^{-2}$	32.4	$3.24\cdot 10^1$
0.01	10^{-2}	100	10^2
0.001	10^{-3}	1000	10^{3}
0.0001	10^{-4}	10 000	10^4
0.000 056	$5.6 \cdot 10^{-5}$	56 000	$5.6 \cdot 10^4$
0.000 01	10^{-5} etc.	100 000	10^5 etc.




FIGURE 34 The definition of plane and solid angles.

for a photographic camera that you can imagine. You have all the money you want. What Challenge 87 s is the shortest shutter time you would achieve?

* *

Can you prove Pythagoras' theorem by geometrical means alone, without using coordinates? (There are more than 30 possibilities.)

* *

Page 58 Why are most planets and moons, including ours, (almost) spherical (see, for example, Challenge 89 s Figure 28)?

* *

A rubber band connects the tips of the two hands of a clock. What is the path followed Challenge 90 s by the mid-point of the band?

* *

There are two important quantities connected to angles. As shown in Figure 34, what is



Challenge 88 s




FIGURE 35 Two equivalent definitions of the sine, cosine, tangent, cotangent, secant and cosecant of an angle.

usually called a *(plane)* angle is defined as the ratio between the lengths of the arc and the radius. A right angle is $\pi/2$ radian (or $\pi/2$ rad) or 90°.

The *solid angle* is the ratio between area and the square of the radius. An eighth of a sphere is $\pi/2$ *steradian* or $\pi/2$ sr. (Mathematicians, of course, would simply leave out the steradian unit.) As a result, a small solid angle shaped like a cone and the angle of the cone tip are *different*. Can you find the relationship?

Challenge 91 s

* *

The definition of angle helps to determine the size of a firework display. Measure the time T, in seconds, between the moment that you see the rocket explode in the sky and the moment you hear the explosion, measure the (plane) angle α – pronounced 'alpha' – of the ball formed by the firework with your hand. The diameter D is

$$D \approx \frac{6 \,\mathrm{m}}{s^{\,\circ}} T \,\alpha \ . \tag{3}$$

Challenge 92 e Why? For more information about fireworks, see the cc.oulu.fi/~kempmp website. By the way, the angular distance between the knuckles of an extended fist are about 3°, 2° and 3°, the size of an extended hand 20°. Can you determine the other angles related to your hand?

* *

Using angles, the *sine*, *cosine*, *tangent*, *cotangent*, *secant* and *cosecant* can be defined, as shown in Figure 35. You should remember this from secondary school. Can you confirm

Challenge 94 e that
$$\sin 15^\circ = (\sqrt{6} - \sqrt{2})/4$$
, $\sin 18^\circ = (-1 + \sqrt{5})/4$, $\sin 36^\circ = \sqrt{10 - 2\sqrt{5}}/4$, $\sin 54^\circ = (-1 + \sqrt{5})/4$



FIGURE 36 How the apparent size of the Moon and the Sun changes during a day.

 $(1+\sqrt{5})/4$ and that $\sin 72^\circ = \sqrt{10+2\sqrt{5}}/4$? Can you show also that

$$\frac{\sin x}{x} = \cos \frac{x}{2} \cos \frac{x}{4} \cos \frac{x}{8} \dots \tag{4}$$

Challenge 95 e is correct?

Challenge 96 e

Challenge 97 s

Challenge 98 d

Measuring angular size with the eye only is tricky. For example, can you say whether the Moon is larger or smaller than the nail of your thumb at the end of your extended arm? Angular size is not an intuitive quantity; it requires measurement instruments.

A famous example, shown in Figure 36, illustrates the difficulty of estimating angles. Both the Sun and the Moon seem larger when they are on the horizon. In ancient times, Ptolemy explained this so-called Moon illusion by an unconscious apparent distance change induced by the human brain. Indeed, the Moon illusion disappears when you look at the Moon through your legs. In fact, the Moon is even further away from the observer when it is just above the horizon, and thus its image is smaller than it was a few hours earlier, when it was high in the sky. Can you confirm this?

The Moon's angular size changes even more due to another effect: the orbit of the Moon round the Earth is elliptical. An example of the consequence is shown in Figure 37.

Galileo also made mistakes. In his famous book, the Dialogues, he says that the curve formed by a thin chain hanging between two nails is a parabola, i.e., the curve defined by $y = x^2$. That is not correct. What is the correct curve? You can observe the shape (approximately) in the shape of suspension bridges.







FIGURE 37 How the size of the Moon actually changes during its orbit (© Anthony Ayiomamitis).




FIGURE 38 A famous puzzle: how are the four radii related?




FIGURE 39 What is the area ABC, given the other three areas and three right angles at O?

Draw three circles, of different sizes, that touch each other, as shown in Figure 38. Now draw a fourth circle in the space between, touching the outer three. What simple relation do the inverse radii of the four circles obey?

* *

Take a tetrahedron OABC whose triangular sides OAB, OBC and OAC are rectangular in O, as shown in Figure 39. In other words, the edges OA, OB and OC are all perpendicular to each other. In the tetrahedron, the areas of the triangles OAB, OBC and OAC are respectively 8, 4 and 1. What is the area of triangle ABC?

Ref. 46 There are many puzzles about ladders. Two are illustrated in Figure 40. If a 5 m ladder



Challenge 99 s

Challenge 100 s




FIGURE 40 Two ladder puzzles: a moderately difficult (left) and a difficult one (right).



FIGURE 41 Anticrepuscular rays - where is the Sun in this situation? (© Peggy Peterson)

is put against a wall in such a way that it just touches a box with 1 m height and depth, how high does the ladder reach? If two ladders are put against two facing walls, and if the lengths of the ladders and the height of the crossing point are known, how distant are the walls?

* *

With two rulers, you can add and subtract numbers by lying them side by side. Are you able to design rulers that allow you to multiply and divide in the same manner?

* *

 $\label{thm:continuous} How many days would a year have if the Earth turned the other way with the same rotation frequency?$

* *

Challenge 105 s The Sun is hidden in the spectacular situation shown in Figure 41 Where is it?



Challenge 101 s

Challenge 102 d







FIGURE 42 Ideal configurations of ropes made of two, three and four strands. In the ideal configuration, the specific pitch angle relative to the equatorial plane – 39.4°, 42.8° and 43.8°, respectively – leads to zero-twist structures. In these ideal configurations, the rope will neither rotate in one nor in the other direction under vertical strain (© Jakob Bohr).

A slightly different, but equally fascinating situation – and useful for getting used to perspective drawing – appears when you have a lighthouse in your back. Can you draw the rays you see in the sky up to the horizon?

Challenge 106 e

Challenge 107 s

* *

Two cylinders of equal radius intersect at a right angle. What is the value of the intersection volume? (First make a drawing.)

* *

Two sides of a hollow cube with side length 1 dm are removed, to yield a tunnel with square opening. Is it true that a cube with edge length 1.06 dm can be made to pass through the hollow cube with side length 1 dm?

Challenge 108 s

* *

Could a two-dimensional universe exist? Alexander Dewdney imagined such a universe in great detail and wrote a well-known book about it. He describes houses, the transportation system, digestion, reproduction and much more. Can you explain why a two-dimensional universe is impossible?

Challenge 109 d

* *

Ropes are wonderful structures. They are flexible, they are helically woven, but despite this, they do not unwind or twist, they are almost inextensible, and their geometry depends little on the material used in making them. What is the origin of all these properties?

Laying rope is an old craft; it is based on a purely geometric result: among all possible helices of *n* strands of given length laid around a central structure of fixed radius, there is one helix for which the number of turns is *maximal*. For purely geometric reasons,





FIGURE 43 An open research problem: What is the ropelength of a tight knot? (© Piotr Pieranski, from Ref. 50)

ropes with that specific number of turns and the corresponding inner radius have the mentioned properties that make ropes so useful. The geometries of ideal ropes made of two, three and four strands are shown in Figure 42.

* *

Some researchers are investigating whether time could be two-dimensional. Can this be?

Challenge 110 s

Challenge 111 s

* *

Other researchers are investigating whether space could have more than three dimensions. Can this be?

* *

One way to compare speeds of animals and machines is to measure them in 'body lengths per second'. The click beetle achieves a value of around 2000 during its jump phase, certain Archaea (bacteria-like) cells a value of 500, and certain hummingbirds 380. These are the record-holders so far. Cars, aiplanes, cheetahs, falcons, crabs, and all other motorical protections are the classes.

Ref. 49 torized systems are much slower.

* *

Draw a square consisting of four equally long connecting line segments hinged at the vertices. Such a structure may be freely deformed into a rhombus if some force is applied. How many additional line interlinks of the same length must be supplemented to avoid this freedom and to prevent the square from being deformed? The extra line interlinks must be in the same plane as the square and each one may only be pegged to others at the endpoints.

Challenge 112 s

* *

Here is a simple challenge on length that nobody has solved yet. Take a piece of ideal rope: of constant radius, ideally flexible, and completely slippery. Tie a tight knot into it, as shown in Figure 43. By how much did the two ends of the rope come closer together?

Challenge 113 r

SUMMARY ABOUT EVERYDAY SPACE AND TIME

Motion defines speed, time and length. Observations of everyday life and precision experiments are conveniently and precisely described by modelling velocity as a Euclidean vector, space as a three-dimensional Euclidean space, and time as a one-dimensional real line. These three definitions form the everyday, or *Galilean*, description of our environ-



ment.

Modelling velocity, time and space as *continuous* quantities is precise and convenient. The modelling works during most of the adventures that follows. However, this common model of space and time *cannot* be confirmed by experiment. For example, no experiments can check distances larger than 10^{25} m or smaller than 10^{-25} m; the continuuum model is likely to be incorrect there. We will find out in the last part of our mountain ascent that this is indeed the case.





CHAPTER 3

HOW TO DESCRIBE MOTION – KINEMATICS

La filosofia è scritta in questo grandissimo libro che continuamente ci sta aperto innanzi agli occhi (io dico l'universo) ... Egli è scritto in lingua matematica.*

Galileo Galilei, Il saggiatore VI.

EXPERIMENTS show that the properties of Galilean time and space are xtracted from the environment both by children and animals. This xtraction has been confirmed for cats, dogs, rats, mice, ants and fish, among others. They all find the same results.

First of all, *motion is change of position with time*. This description is illustrated by rapidly flipping the lower left corners of this book, starting at page 208. Each page simulates an instant of time, and the only change that takes place during motion is in the position of the object, say a stone, represented by the dark spot. The other variations from one picture to the next, which are due to the imperfections of printing techniques, can be taken to simulate the inevitable measurement errors.

Stating that 'motion' the change of position with time is *neither* an explanation *nor* a definition, since both the concepts of time and position are deduced from motion itself. It is only a *description* of motion. Still, the statement is useful, because it allows for high precision, as we will find out by exploring gravitation and electrodynamics. After all, precision is our guiding principle during this promenade. Therefore the detailed description of changes in position has a special name: it is called *kinematics*.

The idea of change of positions implies that the object can be *followed* during its motion. This is not obvious; in the section on quantum theory we will find examples where this is impossible. But in everyday life, objects can always be tracked. The set of all positions taken by an object over time forms its *path* or *trajectory*. The origin of this concept is evident when one watches fireworks or again the flip film in the lower left corners starting at page 208.

In everyday life, animals and humans agree on the Euclidean properties of velocity, space and time. In particular, this implies that a trajectory can be described by specifying three numbers, three *coordinates* (x, y, z) – one for each dimension – as continuous



Ref. 5

^{*} Science is written in this huge book that is continuously open before our eyes (I mean the universe) ... It is written in mathematical language.




FIGURE 44 Two ways to test that the time of free fall does not depend on horizontal velocity.

Vol. III, page 227 functions of time t. (Functions are defined in detail later on.) This is usually written as

$$x = x(t) = (x(t), y(t), z(t))$$
 (5)

For example, already Galileo found, using stopwatch and ruler, that the height z of any thrown or falling stone changes as

$$z(t) = z_0 + v_{z0} (t - t_0) - \frac{1}{2} g (t - t_0)^2$$
(6)

where t_0 is the time the fall starts, z_0 is the initial height, v_{z0} is the initial velocity in the vertical direction and $g=9.8\,\mathrm{m/s^2}$ is a constant that is found to be the same, within about one part in 300, for all falling bodies on all points of the surface of the Earth. Where do the value $9.8\,\mathrm{m/s^2}$ and its slight variations come from? A preliminary answer will be given shortly, but the complete elucidation will occupy us during the larger part of this hike.

The special case with no initial velocity is of great interest. Like a few people before him, Galileo made it clear that *g* is the same for all bodies, if air resistance can be neglected. He had many arguments for this conclusion; can you find one? And of course, his famous experiment at the leaning tower in Pisa confirmed the statement. (It is a *false* urban legend that Galileo never performed the experiment.)

Equation (6) therefore allows us to determine the depth of a well, given the time a stone takes to reach its bottom. The equation also gives the speed v with which one hits the ground after jumping from a tree, namely

$$v = \sqrt{2gh} \tag{7}$$

A height of 3 m yields a velocity of 27 km/h. The velocity is thus proportional only to the square root of the height. Does this mean that one's strong fear of falling results from an overestimation of its actual effects?

Galileo was the first to state an important result about free fall: the motions in the horizontal and vertical directions are *independent*. He showed that the time it takes for



Page 178

Ref. 52

Ref. 53

Challenge 114 s

Challenge 115 s




FIGURE 45 Various types of graphs describing the same path of a thrown stone.

Ref. 54

Challenge 116 s

a cannon ball that is shot exactly horizontally to fall is *independent* of the strength of the gunpowder, as shown in Figure 44. Many great thinkers did not agree with this statement even after his death: in 1658 the Academia del Cimento even organized an experiment to check this assertion, by comparing the flying cannon ball with one that simply fell vertically. Can you imagine how they checked the simultaneity? Figure 44 also shows how you can check this at home. In this experiment, whatever the spring load of the cannon, the two bodies will always collide in mid-air (if the table is high enough), thus proving the assertion.

In other words, a flying canon ball is not accelerated in the horizontal direction. Its horizontal motion is simply unchanging – as long as air resistance is negligible. By extending the description of equation (6) with the two expressions for the horizontal coordinates x and y, namely

$$x(t) = x_0 + v_{x0}(t - t_0)$$

$$y(t) = y_0 + v_{y0}(t - t_0),$$
(8)

Page 39 Challenge 117 s Ref. 55 a *complete* description for the path followed by thrown stones results. A path of this shape is called a *parabola*; it is shown in Figures 18, 44 and 45. (A parabolic shape is also used for light reflectors inside pocket lamps or car headlights. Can you show why?)

Physicists enjoy generalizing the idea of a path. As Figure 45 shows, a path is a trace left in a diagram by a moving object. Depending on what diagram is used, these paths have different names. Space-time diagrams are useful to make the theory of relativity accessible. The *configuration space* is spanned by the coordinates of all particles of a system. For many particles, it has a high number of dimensions. It plays an important role in self-organization. The difference between chaos and order can be described as a difference in the properties of paths in configuration space. *Hodographs*, the paths in 'velocity



space', are used in weather forecasting. The phase space diagram is also called *state space diagram*. It plays an essential role in thermodynamics.

THROWING, JUMPING AND SHOOTING

The kinematic description of motion is useful for answering a whole range of questions.

* *

What is the upper limit for the long jump? The running peak speed world record in 2008 was over $12.5\,\text{m/s}\approx45\,\text{km/h}$ by Usain Bolt, and the 1997 women's record was Ref. 57 $11\,\text{m/s}\approx40\,\text{km/h}$. However, male long jumpers never run much faster than about 9.5 m/s. How much extra jump distance could they achieve if they could run full speed? How could they achieve that? In addition, long jumpers take off at angles of about 20°, as they are not able to achieve a higher angle at the speed they are running. How much would they gain if they could achieve 45°? Is 45° the optimal angle?

Challenge 118 s

* *

What do the athletes Usain Bolt and Michael Johnson, the last two world record holders on the 200 m race at time of this writing, have in common? They were tall, athletic, and had many fast twitch fibres in the muscles. These properties made them good sprinters. A last difference made them world class sprinters: they had a flattened spine, with almost no S-shape. This abnormal condition saves them a little bit of time at every step, because their spine is not as flexible as in usual people. This allows them to excel at short distance races.

* *

Athletes continuously improve speed records. Racing horses do not. Why? For racing horses, breathing rhythm is related to gait; for human, it is not. As a result, racing horses cannot change or improve their technique, and the speed of racing horses is essentially the same since it is measured.

* *

Challenge 119 s

How can the speed of falling rain be measured using an umbrella? The answer is important: the same method can also be used to measure the speed of light, as we will find out later. (Can you guess how?)

Vol. II, page 16

* *

Challenge 120 ny

When a dancer jumps in the air, how many times can he or she rotate around its vertical axis before arriving back on earth?

* *

Ref. 59

Numerous species of moth and butterfly caterpillars shoot away their frass – to put it more crudely: their shit – so that its smell does not help predators to locate them. Stanley Caveney and his team took photographs of this process. Figure 46 shows a caterpillar (yellow) of the skipper *Calpodes ethlius* inside a rolled up green leaf caught in the act. Given that the record distance observed is 1.5 m (though by another species, *Epargyreus*






FIGURE 46 Three superimposed images of a frass pellet shot away by a caterpillar inside a rolled-up leaf (© Stanley Caveney).

Challenge 121 s *clarus*), what is the ejection speed? How do caterpillars achieve it?

* *

What is the horizontal distance one can reach by throwing a stone, given the speed and Challenge 122 s the angle from the horizontal at which it is thrown?

* *

Challenge 123 s What is the maximum numbers of balls that could be juggled at the same time? This is unclear. The human record is ten balls. The internet is full of material on the topic. For robots, the present record is three balls, as shown by the Sarcoman robot on www.physio. northwestern.edu/Secondlevel/Miller/FirstLevel/histresearch.html. It is a challenge for robotics to reach the maximum number of balls in the future.

* *

Is it true that rain drops would kill if it weren't for the air resistance of the atmosphere? What about hail?

* *

Challenge 125 s Are bullets, fired into the air from a gun, dangerous when they fall back down?

* *

Police finds a dead human body at the bottom of cliff with a height of 30 m, at a distance of 12 m from the cliff. Was it suicide or murder?

* *

Ref. 60 All land animals, regardless of their size, achieve jumping heights of at most 2.2 m, as shown in Figure 47. The explanation of this fact takes only two lines. Can you find it? The last two issues arise because the equation (6) describing free fall does not hold in all cases. For example, leaves or potato crisps do not follow it. As Galileo already knew, this






FIGURE 47 The height achieved by jumping land animals.

is a consequence of air resistance; we will discuss it shortly. Because of air resistance, the path of a stone is not a parabola.

In fact, there are other situations where the path of a falling stone is not a parabola, even without air resistance. Can you find one?

Challenge 128 s

ENJOYING VECTORS

Physical quantities with a defined direction, such as speed, are described with three numbers, or three components, and are called *vectors*. Learning to calculate with such multicomponent quantities is an important ability for many sciences. Here is a summary.

Vectors can be pictured by small arrows. Note that vectors do not have specified points at which they start: two arrows with same direction and the same length are the *same* vector, even if they start at different points in space. Since vectors behave like arrows, they can be added and they can be multiplied by numbers. For example, stretching an arrow $\mathbf{a} = (a_x, a_y, a_z)$ by a number c corresponds, in component notation, to the vector $c\mathbf{a} = (ca_x, ca_y, ca_z)$.

In precise, mathematical language, a vector is an element of a set, called *vector space*, in which the following properties hold for all vectors a and b and for all numbers c and d:

$$c(\mathbf{a} + \mathbf{b}) = c\mathbf{a} + c\mathbf{b}$$
, $(c + d)\mathbf{a} = c\mathbf{a} + d\mathbf{a}$, $(cd)\mathbf{a} = c(d\mathbf{a})$ and $1\mathbf{a} = \mathbf{a}$. (9)

Challenge 129 s

Examples of vector spaces are the set of all *positions* of an object, or the set of all its possible velocities. Does the set of all rotations form a vector space?

Challenge 130 e

All vector spaces allow defining a unique *null vector* and a unique *negative vector* for each vector.

In most vector spaces of importance in science the concept of *length* (specifying the 'magnitude') can be introduced. This is done via an intermediate step, namely the intro-



duction of the scalar product of two vectors. The product is called 'scalar' because its result is a scalar; a *scalar* is a number that is the same for all observers; for example, it is the same for observers with different orientations.* The *scalar product* between two vectors **a** and **b** is a number that satisfies

$$aa \ge 0$$
,
 $ab = ba$,
 $(a + a')b = ab + a'b$,
 $a(b + b') = ab + ab'$ and
 $(ca)b = a(cb) = c(ab)$.

This definition of a scalar product is not unique; however, there is a *standard* scalar product. In Cartesian coordinate notation, the standard scalar product is given by

$$ab = a_x b_x + a_y b_y + a_z b_z . ag{11}$$

Challenge 131 e

If the scalar product of two vectors vanishes the two vectors are *orthogonal*, at a right angle to each other. (Show it!)

The *length* or *magnitude* or *norm* of a vector can then be defined as the square root of the scalar product of a vector with itself: $a = \sqrt{aa}$. Often, and also in this text, lengths are written in italic letters, whereas vectors are written in bold letters. A vector space with a scalar product is called an *Euclidean* vector space.

The scalar product is also useful for specifying directions. Indeed, the scalar product between two vectors encodes the angle between them. Can you deduce this important relation?

Challenge 132 s

WHAT IS REST? WHAT IS VELOCITY?

In the Galilean description of nature, motion and rest are opposites. In other words, a body is at rest when its position, i.e., its coordinates, do not change with time. In other words, (Galilean) *rest* is defined as

$$x(t) = \text{const} . (12)$$

We recall that x(t) is the abbreviation for the three coordinates (x(t), y(t), z(t)). Later we will see that this definition of rest, contrary to first impressions, is not much use and will have to be expanded. Nevertheless, any definition of rest implies that non-resting objects can be distinguished by comparing the rapidity of their displacement. Thus we can define the *velocity* v of an object as the change of its position x with time t. This is usually written as

$$v = \frac{\mathrm{d}x}{\mathrm{d}t} \ . \tag{13}$$

^{*} We mention that in mathematics, a scalar is a *number*; in physics, a scalar is an *invariant* number, i.e., a number that is the same for all observers. Likewise, in mathematics, a vector is an element of a vector space; in physics, a vector is an invariant element of a vector space, i.e., a quantity whose coordinates, when observed by different observers, change like the components of velocity.




FIGURE 48 The derivative in a point as the limit of secants.

In this expression, valid for each coordinate separately, d/dt means 'change with time'. We can thus say that velocity is the *derivative* of position with respect to time. The *speed v* is the name given to the magnitude of the velocity v. Thus we have $v = \sqrt{vv}$. Derivatives are written as fractions in order to remind the reader that they are derived from the idea of slope. The expression

$$\frac{\mathrm{d}s}{\mathrm{d}t}$$
 is meant as an abbreviation of $\lim_{\Delta t \to 0} \frac{\Delta s}{\Delta t}$, (14)

a shorthand for saying that the *derivative at a point* is the limit of the secant slopes in the neighbourhood of the point, as shown in Figure 48. This definition implies the working rules

Challenge 133 e

$$\frac{\mathrm{d}(s+r)}{\mathrm{d}t} = \frac{\mathrm{d}s}{\mathrm{d}t} + \frac{\mathrm{d}r}{\mathrm{d}t} \quad , \quad \frac{\mathrm{d}(cs)}{\mathrm{d}t} = c\frac{\mathrm{d}s}{\mathrm{d}t} \quad , \quad \frac{\mathrm{d}}{\mathrm{d}t}\frac{\mathrm{d}s}{\mathrm{d}t} = \frac{\mathrm{d}^2s}{\mathrm{d}t^2} \quad , \quad \frac{\mathrm{d}(sr)}{\mathrm{d}t} = \frac{\mathrm{d}s}{\mathrm{d}t}r + s\frac{\mathrm{d}r}{\mathrm{d}t} \quad , \quad (15)$$

c being any number. This is all one ever needs to know about derivatives. Quantities such as d*t* and d*s*, sometimes useful by themselves, are called *differentials*. These concepts are due to Gottfried Wilhelm Leibniz.* Derivatives lie at the basis of all calculations based on the continuity of space and time. Leibniz was the person who made it possible to describe and use velocity in physical formulae and, in particular, to use the idea of velocity at a given point in time or space for calculations.

The definition of velocity assumes that it makes sense to take the limit $\Delta t \to 0$. In other words, it is assumed that *infinitely small* time intervals do exist in nature. The definition of velocity with derivatives is possible only because both space and time are described by sets which are *continuous*, or in mathematical language, *connected and complete*. In the rest of our walk we shall not forget that from the beginning of classical

^{*} Gottfried Wilhelm Leibniz (b. 1646 Leipzig, d. 1716 Hannover), Saxon lawyer, physicist, mathematician, philosopher, diplomat and historian. He was one of the great minds of mankind; he invented the differential calculus (before Newton) and published many influential and successful books in the various fields he explored, among them *De arte combinatoria*, *Hypothesis physica nova*, *Discours de métaphysique*, *Nouveaux essais sur l'entendement humain*, the *Théodicée* and the *Monadologia*.



FIGURE 49 Gottfried Wilhelm Leibniz (1646-1716).

physics, *infinities* are present in its description of nature. The infinitely small is part of our definition of velocity. Indeed, differential calculus can be defined as the study of infinity and its uses. We thus discover that the appearance of infinity does not automatically render a description impossible or imprecise. In order to remain precise, physicists use only the smallest two of the various possible types of infinities. Their precise definition and an overview of other types are introduced later on.

Vol. III, page 225

Ref. 61

Challenge 134 e

Vol. VI, page 59

The appearance of infinity in the usual description of motion was first criticized in his famous ironical arguments by Zeno of Elea (around 445 BCE), a disciple of Parmenides. In his so-called third argument, Zeno explains that since at every instant a given object occupies a part of space corresponding to its size, the notion of velocity at a given instant makes no sense; he provokingly concludes that therefore motion does not exist. Nowadays we would not call this an argument against the *existence* of motion, but against its usual *description*, in particular against the use of infinitely divisible space and time. (Do you agree?) Nevertheless, the description criticized by Zeno actually works quite well in everyday life. The reason is simple but deep: in daily life, changes are indeed continuous.

Large changes in nature are made up of many small changes. This property of nature is not obvious. For example, we note that we have tacitly assumed that the path of an object is not a fractal or some other badly behaved entity. In everyday life this is correct; in other domains of nature it is not. The doubts of Zeno will be partly rehabilitated later in our walk, and increasingly so the more we proceed. The rehabilitation is only partial, as the final solution will be different from that which he envisaged; on the other hand, the doubts about the idea of 'velocity at a point' will turn out to be well-founded. For the moment though, we have no choice: we continue with the basic assumption that in nature changes happen smoothly.

Why is velocity necessary as a concept? Aiming for precision in the description of motion, we need to find the complete list of aspects necessary to specify the state of an object. The concept of velocity is obviously on this list.

ACCELERATION

Continuing along the same line, we call *acceleration* \boldsymbol{a} of a body the change of velocity \boldsymbol{v} with time, or

$$a = \frac{\mathrm{d}v}{\mathrm{d}t} = \frac{\mathrm{d}^2x}{\mathrm{d}t^2} \,. \tag{16}$$

TABLE 13 Some measured acceleration values.

OBSERVATION	ACCELERATION
What is the lowest you can find?	Challenge 135 s
Back-acceleration of the galaxy M82 by its ejected jet	$10\mathrm{fm/s^2}$
Acceleration of a young star by an ejected jet	$10 \mathrm{pm/s^2}$
Fathoumi Acceleration of the Sun in its orbit around the Milky Way	$0.2\mathrm{nm/s}^2$
Deceleration of the Pioneer satellites, due to heat radiation imbalance	$0.8\mathrm{nm/s^2}$
Centrifugal acceleration at Equator due to Earth's rotation	$33 \mathrm{mm/s}^2$
Electron acceleration in household electricity wire due to alternating current	$50 \mathrm{mm/s}^2$
Acceleration of fast underground train	$1.3\mathrm{m/s^2}$
Gravitational acceleration on the Moon	$1.6\mathrm{m/s^2}$
Minimum deceleration of a car, by law, on modern dry asfalt	$5.5\mathrm{m/s^2}$
Gravitational acceleration on the Earth's surface, depending on location	$9.8 \pm 0.3 \mathrm{m/s^2}$
Standard gravitational acceleration	$9.80665 \mathrm{m/s^2}$
Highest acceleration for a car or motorbike with engine-driven wheels	$15 \mathrm{m/s}^2$
Space rockets at take-off	$20 \text{ to } 90 \text{ m/s}^2$
Acceleration of cheetah	32 m/s^2
Gravitational acceleration on Jupiter's surface	$25 \mathrm{m/s}^2$
Flying fly (Musca domestica)	$c. 100 \mathrm{m/s^2}$
Acceleration of thrown stone	$c. 120 \mathrm{m/s}^2$
Acceleration that triggers air bags in cars	360m/s^2
Fastest leg-powered acceleration (by the froghopper, <i>Philaenus spumarius</i> , an insect)	$4 \mathrm{km/s^2}$
Tennis ball against wall	$0.1 \mathrm{Mm/s^2}$
Bullet acceleration in rifle	$2 \mathrm{Mm/s}^2$
Fastest centrifuges	$0.1 \mathrm{Gm/s}^2$
Acceleration of protons in large accelerator	$90 \mathrm{Tm/s^2}$
Acceleration of protons inside nucleus	10^{31}m/s^2
Highest possible acceleration in nature	10^{52}m/s^2

Acceleration is what we feel when the Earth trembles, an aeroplane takes off, or a bicycle goes round a corner. More examples are given in Table 13. Like velocity, acceleration has both a magnitude and a direction, properties indicated by the use of **bold** letters for their abbreviations. In short, acceleration, like velocity, is a vector quantity.

Acceleration is felt. Our body is deformed and the sensors in our semicircular canals in the ear feel it. Higher accelerations can have stronger effects. For example, when accelerating a sitting person in the direction of the head at two or three times the value of usual gravitational acceleration, eyes stop working and the sight is greyed out, because the blood cannot reach the eye any more. Between 3 and 5g of continuous acceleration, or 7 to 9g of short time acceleration, consciousness is lost, because the brain does not re-

TABLE 14 Some acceleration sensors.

-		
MEASUREMENT	SENSOR	RANGE
Direction of gravity in plants (roots, trunk, branches, leaves)	statoliths in cells	$0 \text{ to } 10 \text{ m/s}^2$
Direction and value of accelerations in mammals	the membranes in each semicircular canal, and the utricule and saccule in the inner ear	0 to 20 m/s^2
Direction and value of acceleration in modern step counters for hikers	piezoelectric sensors	$0 \text{ to } 20 \text{ m/s}^2$
Direction and value of acceleration in car crashes	airbag sensor using piezoelectric ceramics	0 to 2000 m/s ²

ceive enough blood, and blood may leak out of the feet or lower legs. High acceleration in the direction of the feet of a sitting person can lead to haemorrhagic strokes in the brain. The people most at risk are jet pilots; they have special clothes that send compressed air onto the pilot's bodies to avoid blood accumulating in the wrong places.

In a usual car, or on a motorbike, we can *feel* being accelerated. (These accelerations are below 1g and are therefore harmless.) Acceleration sensors, such are those listed in Table 14 or those shown in Figure 50, all work in this way. Can you think of a situation where one is accelerated but does *not* feel it?

Higher derivatives than acceleration can also be defined in the same manner. They add little to the description of nature, because – as we will show shortly – neither these higher derivatives nor even acceleration itself are useful for the description of the state of motion of a system.

OBJECTS AND POINT PARTICLES

Wenn ich den Gegenstand kenne, so kenne ich auch sämtliche Möglichkeiten seines Vorkommens in Sachverhalten.*
Ludwig Wittgenstein, *Tractatus*, 2.0123

One aim of the study of motion is to find a complete and precise description of both states and objects. With the help of the concept of space, the description of objects can be refined considerably. In particular, we know from experience that all objects seen in daily life have an important property: they can be divided into *parts*. Often this observation is expressed by saying that all objects, or bodies, have two properties. First, they are made out of *matter*,** defined as that aspect of an object responsible for its impenetrability, i.e., the property preventing two objects from being in the same place. Secondly, bodies have a certain form or *shape*, defined as the precise way in which this impenetrability is distributed in space.

In order to describe motion as accurately as possible, it is convenient to start with

Challenge 136 s


Challenge 137 s

Challenge 138 e


Ref. 63


^{* &#}x27;If I know an object, then I also know all the possibilities of its occurrence in atomic facts.'

^{**} Matter is a word derived from the Latin 'materia', which originally meant 'wood' and was derived via intermediate steps from 'mater', meaning 'mother'.












FIGURE 50 Three accelerometers: a one-axis piezoelectric airbag sensor, a three-axis capacitive accelerometer, and the utricule and saccule near the three semicircular canals inside the human ear (© Bosch, Rieker Electronics, Northwestern University).

those bodies that are as simple as possible. In general, the smaller a body, the simpler it is. A body that is so small that its parts no longer need to be taken into account is called a *particle*. (The older term *corpuscle* has fallen out of fashion.) Particles are thus idealized small stones. The extreme case, a particle whose size is *negligible* compared with the dimensions of its motion, so that its position is described completely by a *single* triplet of coordinates, is called a *point particle* or a *point mass* or a *mass point*. In equation (6), the stone was assumed to be such a point particle.

Do point-like objects, i.e., objects smaller than anything one can measure, exist in daily life? Yes and no. The most notable examples are the stars. At present, angular sizes as small as $2 \mu rad$ can be measured, a limit given by the fluctuations of the air in the atmosphere. In space, such as for the Hubble telescope orbiting the Earth, the angular limit is due to the diameter of the telescope and is of the order of $10 \, nrad$. Practically all stars seen from Earth are smaller than that, and are thus effectively 'point-like', even when seen with the most powerful telescopes.

As an exception to the general rule, the size of a few large and nearby stars, of red giant type, can be measured with special instruments.* Betelgeuse, the higher of the two shoulders of Orion shown in Figure 51, Mira in Cetus, Antares in Scorpio, Aldebaran in

^{*} The website www.astro.uiuc.edu/~kaler/sow/sowlist.html gives an introduction to the different types of stars. The www.astro.wisc.edu/~dolan/constellations website provides detailed and interesting information about constellations.






FIGURE 51 Orion in natural colours (© Matthew Spinelli) and Betelgeuse (ESA, NASA).




FIGURE 52 Regulus and Mars, photographed with 10 second exposure time on 4 June 2010 with a wobbling camera, show the difference between a point-like star that twinkles and an extended planet that does not (© Jürgen Michelberger).

Ref. 64

Challenge 139 s

Challenge 140 e

Taurus and Sirius in Canis Major are examples of stars whose size has been measured; they are all only a few light years from Earth. Of course, like the Sun, also all other stars have a finite size, but one cannot prove this by measuring dimensions in photographs. (True?)

The difference between 'point-like' and finite-size sources can be seen with the naked eye: at night, stars twinkle, but planets do not. (Check it!) A beautiful visualization is shown in Figure 52. This effect is due to the turbulence of air. Turbulence has an effect on the almost point-like stars because it deflects light rays by small amounts. On the other hand, air turbulence is too weak to lead to twinkling of sources of larger angular size, such as planets or artificial satellites,* because the deflection is averaged out in this

For an overview of the planets, see the beautiful book by K. R. Lang & C. A. Whitney, Vagabonds de l'espace – Exploration et découverte dans le système solaire, Springer Verlag, 1993. The most beautiful pictures of the stars can be found in D. Malin, A View of the Universe, Sky Publishing and Cambridge University Press, 1993.

^{*} A *satellite* is an object circling a planet, like the Moon; an *artificial satellite* is a system put into orbit by humans, like the Sputniks.

case.

Challenge 141 s

An object is *point-like for the naked eye* if its angular size is smaller than about 2' = 0.6 mrad. Can you estimate the size of a 'point-like' dust particle? By the way, an object is *invisible* to the naked eye if it is point-like *and* if its luminosity, i.e., the intensity of the light from the object reaching the eye, is below some critical value. Can you estimate whether there are any man-made objects visible from the Moon, or from the space shuttle?

Challenge 142 s

The above definition of 'point-like' in everyday life is obviously misleading. Do proper, real point particles exist? In fact, is it at all possible to show that a particle has vanishing size? This question will be central in the last part of our walk. In the same way, we need to ask and check whether points in space do exist. Our walk will lead us to the astonishing result that all the answers to these questions are negative. Can you imagine why? Do not be disappointed if you find this issue difficult; many brilliant minds have had the same problem.

Challenge 143 s

However, many particles, such as electrons, quarks or photons are point-like for all practical purposes. Once one knows how to describe the motion of point particles, one can also describe the motion of extended bodies, rigid or deformable, by assuming that they are made of parts. This is the same approach as describing the motion of an animal as a whole by combining the motion of its various body parts. The simplest description, the *continuum approximation*, describes extended bodies as an infinite collection of point particles. It allows us to understand and to predict the motion of milk and honey, the motion of the air in hurricanes and of perfume in rooms. The motion of fire and all other gaseous bodies, the bending of bamboo in the wind, the shape changes of chewing gum, and the growth of plants and animals can also be described in this way.

Ref. 65

Vol. IV, page 13

A more precise description than the continuum approximation is given below. Nevertheless, all observations so far have confirmed that the motion of large bodies can be described to high precision as the result of the motion of their parts. This approach will guide us through the first five volumes of our mountain ascent. Only in the final volume will we discover that, at a fundamental scale, this decomposition is impossible.

LEGS AND WHEELS

The parts of a body determine its shape. Shape is an important aspect of bodies: among other things, it tells us how to count them. In particular, living beings are always made of a single body. This is not an empty statement: from this fact we can deduce that animals cannot have wheels or propellers, but only legs, fins, or wings. Why?

Vol. V, page 289 Ref. 66 Living beings have only one surface; simply put, they have only one piece of skin. Mathematically speaking, animals are *connected*. This is often assumed to be obvious, and it is often mentioned that the blood supply, the nerves and the lymphatic connections to a rotating part would get tangled up. However, this argument is not so simple, as Figure 53 shows. It shows that it is indeed possible to rotate a body continuously against a second one, without tangling up the connections. Can you find an example for this kind of motion in your own body? Are you able to see how many cables may be attached to the rotating body of the figure without hindering the rotation?

Challenge 144 s

Challenge 145 s

Despite the possibility of animals having rotating parts, the method of Figure 53 still cannot be used to make a practical wheel or propeller. Can you see why? Evolution had

Challenge 146 s

FIGURE 53 How an object can rotate continuously without tangling up the connection to a second object.




FIGURE 54 Legs and 'wheels' in living beings: the red millipede *Aphistogoniulus erythrocephalus* (15 cm body length), a gekko on a glass pane (15 cm body length), an amoeba *Amoeba proteus* (1 mm size), the rolling shrimp *Nannosquilla decemspinosa* (2 cm body length, 1.5 rotations per second, up to 2 m, can even roll slightly uphill slopes) and the rolling caterpillar *Pleurotya ruralis* (can only roll downhill, to escape predators), (© David Parks, Marcel Berendsen, Antonio Guillén, Robert Full, John Brackenbury / Science Photo Library).

no choice: it had to avoid animals with parts rotating around axles. That is the reason that propellers and wheels do not exist in nature. Of course, this limitation does not rule out that living bodies move by rotation as a whole: tumbleweed, seeds from various trees, some insects, several spiders,* certain other animals, children and dancers occasionally move by rolling or rotating as a whole.

^{*} Rolling is known for desert spiders of the *Cebrennus* and the *Carparachne* genus; films can be found on www.youtube.com/watch?v=5XwIXFFVOSA and www.youtube.com/watch?v=ozn31QBOHtk. Cebrennus seems even to be able to accelerate its rolling motion with its legs.

Ref. 68 Ref. 69

Vol. V, page 208 Vol. V, page 210 Single bodies, and thus all living beings, can only move through deformation of their shape: therefore they are limited to walking, running, rolling, crawling or flapping wings or fins. Extreme examples of leg use in nature are shown in Figure 54. The most extreme example are rolling spiders living in the sand in Morocco; they use their legs to accelerate and steer the rolling direction. Walking on water is shown in Figure 106 on page 149; examples of wings are given later on, as are the various types of deformations that allow swimming in water. In contrast, systems of several bodies, such as bicycles, pedal boats or other machines, can move without any change of shape of their components, thus enabling the use of axles with wheels, propellers or other rotating devices.*

In summary, whenever we observe a construction in which some part is turning continuously (and without the 'wiring' of Figure 53) we know immediately that it is an artefact: it is a machine, not a living being (but built by one). However, like so many statements about living creatures, this one also has exceptions. The distinction between one and two bodies is poorly defined if the whole system is made of only a few molecules. This happens most clearly inside bacteria. Organisms such as *Escherichia coli*, the well-known bacterium found in the human gut, or bacteria from the *Salmonella* family, all swim using flagella. *Flagella* are thin filaments, similar to tiny hairs that stick out of the cell membrane. In the 1970s it was shown that each flagellum, made of one or a few long molecules with a diameter of a few tens of nanometres, does in fact turn about its axis. A bacterium is able to turn its flagella in both clockwise and anticlockwise directions, can achieve more than 1000 turns per second, and can turn all its flagella in perfect synchronization. (These wheels are so tiny that they do not need a mechanical connection.) Therefore wheels actually do exist in living beings, albeit only tiny ones. But let us now continue with our study of simple objects.

Vol. V, page 210

Ref. 70

CURIOSITIES AND FUN CHALLENGES ABOUT KINEMATICS

Challenge 148 s What is the biggest wheel ever made?

* *

A soccer ball is shot, by a goalkeeper, with around 30 m/s. Calculate the distance it should fly and compare it with the distances found in a soccer match. Where does the difference come from?

Challenge 149 s

* *

A train starts to travel at a constant speed of 10 m/s between two cities A and B, 36 km apart. The train will take one hour for the journey. At the same time as the train, a fast dove starts to fly from A to B, at 20 m/s. Being faster than the train, the dove arrives at B first. The dove then flies back towards A; when it meets the train, it turns back again,

^{*} Despite the disadvantage of not being able to use rotating parts and of being restricted to one piece only, nature's moving constructions, usually called animals, often outperform human built machines. As an example, compare the size of the smallest flying systems built by evolution with those built by humans. (See, e.g., pixelito.reference.be.) There are two reasons for this discrepancy. First, nature's systems have integrated repair and maintenance systems. Second, nature can build large structures inside containers with small openings. In fact, nature is very good at what people do when they build sailing ships inside glass bottles. The human body is full of such examples; can you name a few?

Challenge 150 e

Challenge 151 e

Challenge 152 e



FIGURE 55 Are comets, such as the beautiful comet McNaught seen in 2007, images or bodies? How can one settle the issue? (© Robert McNaught)

to city B. It goes on flying back and forward until the train reaches B. What distance did the dove cover?

* *

Balance a pencil vertically (tip upwards!) on a piece of paper near the edge of a table. How can you pull out the paper without letting the pencil fall?

* *

Is a return flight by plane – from a point A to B and back to A – faster if the wind blows or if it does not?

* *

The level of acceleration a human can survive depends on the duration over which one is subjected to it. For a tenth of a second, $30 g = 300 \,\text{m/s}^2$, as generated by an ejector seat in an aeroplane, is acceptable. (It seems that the record acceleration a human has survived is about $80 g = 800 \,\text{m/s}^2$.) But as a rule of thumb it is said that accelerations of $15 g = 150 \,\text{m/s}^2$ or more are fatal.

* *

The highest *microscopic* accelerations are observed in particle collisions, where one gets values up to 10^{35} m/s². The highest *macroscopic* accelerations are probably found in the collapsing interiors of *supernovae*, exploding stars which can be so bright as to be visible



FIGURE 56 Observation of sonoluminescence (© Detlev Lohse).

in the sky even during the daytime. A candidate on Earth is the interior of collapsing bubbles in liquids, a process called *cavitation*. Cavitation often produces light, an effect discovered by Frenzel and Schultes in 1934 and called *sonoluminescence*. (See Figure 56.) It appears most prominently when air bubbles in water are expanded and contracted by underwater loudspeakers at around 30 kHz and allows precise measurements of bubble motion. At a certain threshold intensity, the bubble radius changes at 1500 m/s in as little as a few μ m, giving an acceleration of several 10^{11} m/s².

* *

Legs are easy to build. Nature has even produced a millipede, *Illacme plenipes*, that has 750 legs. The animal is 3 to 4 cm long and about 0.5 mm wide. This seems to be the record so far. In contrast to its name, no millipede actually has a thousand legs.

SUMMARY OF KINEMATICS

The description of everyday motion of mass points with three coordinates as (x(t), y(t), z(t)) is simple, precise and complete. It assumes that objects can be followed along their paths. Therefore, the description often does not work for an important case: the motion of images.





Ref. 73

Ref. 75

CHAPTER 4

FROM OBJECTS AND IMAGES TO CONSERVATION

ALKING through a forest we observe two rather different types of motion: e see the breeze move the leaves, and at the same time, on the ground, e see their shadows move. Shadows are a simple type of image. Both objects and images are able to move; both change position over time. Running tigers, falling snowflakes, and material ejected by volcanoes, but also the shadow following our body, the beam of light circling the tower of a lighthouse on a misty night, and the rainbow that constantly keeps the same apparent distance from us are examples of motion.

Both objects and images differ from their environment in that they have *boundaries* defining their size and shape. But everybody who has ever seen an animated cartoon knows that images can move in more surprising ways than objects. Images can change their size and shape, they can even change colour, a feat only few objects are able to perform.* Images can appear and disappear without trace, multiply, interpenetrate, go backwards in time and defy gravity or any other force. Images, even ordinary shadows, can move faster than light. Images can float in space and keep the same distance from approaching objects. Objects can do almost none of this. In general, the 'laws of cartoon physics' are rather different from those in nature. In fact, the motion of images does not seem to follow any rules, in contrast to the motion of objects. We feel the need for precise criteria allowing the two cases to be distinguished.

Making a clear distinction between images and objects is performed using the same method that children or animals use when they stand in front of a mirror for the first time: they try to *touch* what they see. Indeed, if we are able to touch what we see – or more precisely, if we are able to move it – we call it an *object*, otherwise an *image*.** Images cannot be touched, but objects can. Images cannot hit each other, but objects can. And as everybody knows, touching something means feeling that it resists movement. Certain bodies, such as butterflies, pose little resistance and are moved with ease, others,

* Excluding very slow changes such as the change of colour of leaves in the Fall, in nature only certain crystals, the octopus, the chameleon and a few other animals achieve this. Of man-made objects, television,

Vol. IV, page 119

Challenge 153 s

e 153 s computer displays, heated objects and certain lasers can do it. Do you know more examples? An excellent source of information on the topic of colour is the book by K. NASSAU, *The Physics and Chemistry of Colour – the fifteen causes of colour*, J. Wiley & Sons, 1983. In the popular science domain, the most beautiful book is the classic work by the Flemish astronomer MARCEL G. J. MINNAERT, *Light and Colour in the Outdoors*, Springer, 1993, an updated version based on his wonderful book series, *De natuurkunde van 't vrije veld*, Ref. 74 Thieme & Cie, 1937. Reading it is a must for all natural scientists. On the web, there is also the – simpler, but excellent – webexhibits.org/causesofcolour website.

^{**} One could propose including the requirement that objects may be rotated; however, this requirement, surprisingly, gives difficulties in the case of atoms, as explained on page 72 in Volume IV.




FIGURE 57 In which direction does the bicycle turn?

such as ships, resist more, and are moved with more difficulty. This resistance to motion – more precisely, to change of motion – is called *inertia*, and the difficulty with which a body can be moved is called its *(inertial) mass*. Images have neither inertia nor mass.

Summing up, for the description of motion we must distinguish bodies, which can be touched and are impenetrable, from images, which cannot and are not. Everything visible is either an object or an image; there is no third possibility. (Do you agree?) If the object is so far away that it cannot be touched, such as a star or a comet, it can be difficult to decide whether one is dealing with an image or an object; we will encounter this difficulty repeatedly. For example, how would you show that comets – such as the beautiful example of Figure 55 – are objects and not images, as Galileo (falsely) claimed?

Challenge 155 s

Ref. 76

Challenge 154 s

In the same way that objects are made of *matter*, images are made of *radiation*. Images are the domain of shadow theatre, cinema, television, computer graphics, belief systems and drug experts. Photographs, motion pictures, ghosts, angels, dreams and many hallucinations are images (sometimes coupled with brain malfunction). To understand images, we need to study radiation (plus the eye and the brain). However, due to the importance of objects – after all we are objects ourselves – we study the latter first.

MOTION AND CONTACT

Openocritus affirms that there is only one type of movement: That resulting from collision.

Aetius, Opinions.

Ref. 77

When a child rides a unicycle, she or he makes use of a general rule in our world: one body acting on another puts it in motion. Indeed, in about six hours, anybody can learn to ride and enjoy a unicycle. As in all of life's pleasures, such as toys, animals, women, machines, children, men, the sea, wind, cinema, juggling, rambling and loving, something pushes something else. Thus our first challenge is to describe this transfer of motion in more precise terms.

Contact is not the only way to put something into motion; a counter-example is an apple falling from a tree or one magnet pulling another. Non-contact influences are more fascinating: nothing is hidden, but nevertheless something mysterious happens. Contact motion seems easier to grasp, and that is why one usually starts with it. However, despite this choice, non-contact forces are not easily avoided. Taking this choice we will make a similar experience to that of cyclists. (See Figure 57.) If you ride a bicycle at a sustained speed and try to turn left by pushing the right-hand steering bar, you will turn *right*. By



FIGURE 58 Collisions define mass.

FIGURE 59 The standard kilogram (© BIPM).

Challenge 156 s

the way, this surprising effect, also known to motor bike riders, obviously works only above a certain minimal speed. Can you determine what this speed is? Be careful! Too strong a push will make you fall.

Something similar will happen to us as well; despite our choice for contact motion, the rest of our walk will rapidly force us to study non-contact interactions.

WHAT IS MASS?

Δός μοί (φησι) ποῦ στῷ καὶ κινῷ τὴν γῆν.
Da ubi consistam, et terram movebo.*

Archimedes

When we push something we are unfamiliar with, such as when we kick an object on the street, we automatically pay attention to the same aspect that children explore when they stand in front of a mirror for the first time, or when they see a red laser spot for the first time. They check whether the unknown entity can be pushed or caught, and they pay attention to how the unknown object moves under their influence. The high precision version of the experiment is shown in Figure 58. Repeating the experiment with various pairs of objects, we find – as in everyday life – that a *fixed* quantity m_i can be ascribed to

^{* &#}x27;Give me a place to stand, and I'll move the Earth.' Archimedes (c. 283–212), Greek scientist and engineer.

Ref. 78 This phrase is attributed to him by Pappus. Already Archimedes knew that the distinction used by lawyers between movable and immovable property made no sense.



FIGURE 60 Antoine Lavoisier (1743–1794) and his wife.

every object i, determined by the relation

$$\frac{m_2}{m_1} = -\frac{\Delta v_1}{\Delta v_2} \tag{17}$$

where Δv is the velocity change produced by the collision. The more difficult it is to move an object, the higher the number. The number m_i is called the *mass* of the object i.

In order to have mass values that are common to everybody, the mass value for one particular, selected object has to be fixed in advance. This special object, shown in Figure 59, is called the *standard kilogram* and is kept with great care in a glass container in Sèvres near Paris. The standard kilogram is touched only once every few years because otherwise dust, humidity, or scratches would change its mass. By the way, the standard kilogram is *not* kept under vacuum, because this would lead to outgassing and thus to changes in its mass. The standard kilogram determines the value of the mass of every other object in the world.

The *mass* thus measures the difficulty of getting something moving. High masses are harder to move than low masses. Obviously, only objects have mass; images don't. (By the way, the word 'mass' is derived, via Latin, from the Greek $\mu\alpha\zeta\alpha$ – bread – or the Hebrew 'mazza' – unleavened bread. That is quite a change in meaning.)

Experiments with everyday life objects also show that throughout any collision, the sum of all masses is *conserved*:

$$\sum_{i} m_i = \text{const} . ag{18}$$

The principle of conservation of mass was first stated by Antoine-Laurent Lavoisier.*

^{*} Antoine-Laurent Lavoisier (1743–1794), French chemist and a genius. Lavoisier was the first to understand that combustion is a reaction with oxygen; he discovered the components of water and introduced mass measurements into chemistry. There is a good, but most probably false story about him: When he was (unjustly) sentenced to the guillotine during the French revolution, he decided to use the situations for a scientific experiment. He would try to blink his eyes as frequently as possible after his head was cut off, in



FIGURE 61 Christiaan Huygens (1629–1695).

Conservation of mass also implies that the mass of a composite system is the sum of the mass of the components. In short, *Galilean mass is a measure for the quantity of matter.*

In a famous experiment in the sixteenth century, for several weeks Santorio Santorio (Sanctorius) (1561–1636), friend of Galileo, lived with all his food and drink supply, and also his toilet, on a large balance. He wanted to test mass conservation. How did the measured weight change with time?

Challenge 157 s

MOMENTUM AND MASS

The definition of mass can also be given in another way. We can ascribe a number m_i to every object i such that for collisions free of outside interference the following sum is unchanged *throughout* the collision:

$$\sum_{i} m_{i} v_{i} = \text{const} . \tag{19}$$

The product of the velocity v_i and the mass m_i is called the *momentum* of the body. The sum, or *total momentum* of the system, is the same before and after the collision; momentum is a *conserved* quantity.

▶ Momentum conservation defines mass.

The two conservation principles (18) and (19) were first stated in this way by the important Dutch physicist Christiaan Huygens.* Momentum and mass are conserved in everyday motion of objects. Neither quantity can be defined for the motion of images. Some typical momentum values are given in Table 15.

Momentum conservation implies that when a moving sphere hits a resting one of the same mass and without loss of energy, a simple rule determines the angle between the

order to show others how long it takes to lose consciousness. Lavoisier managed to blink eleven times. It is unclear whether the story is true or not. It is known, however, that it could be true. Indeed, if a decapitated has no pain or shock, he can remain conscious for up to half a minute.

^{*} Christiaan Huygens (b. 1629 's Gravenhage, d. 1695 Hofwyck) was one of the main physicists and mathematicians of his time. Huygens clarified the concepts of mechanics; he also was one of the first to show that light is a wave. He wrote influential books on probability theory, clock mechanisms, optics and astronomy. Among other achievements, Huygens showed that the Orion Nebula consists of stars, discovered Titan, the moon of Saturn, and showed that the rings of Saturn consist of rock. (This is in contrast to Saturn itself, whose density is lower than that of water.)




FIGURE 62 Is this dangerous?

TABLE 15 Some measured momentum values.

0)
O B S E R V A T I O N	Моментим
Images	0
Momentum of a green photon	$1.2 \cdot 10^{-27} \text{ Ns}$
Average momentum of oxygen molecule in air	$10^{-26} \mathrm{Ns}$
X-ray photon momentum	10^{-23} Ns
γ photon momentum	10^{-17} Ns
Highest particle momentum in accelerators	1 fNs
Highest possible momentum of a single elementary	6.5 Ns
particle - the Planck momentum	
Fast billiard ball	3 Ns
Flying rifle bullet	10 Ns
Box punch	15 to 50 Ns
Comfortably walking human	80 Ns
Lion paw strike	kNs
Whale tail blow	kNs
Car on highway	40 kNs
Impact of meteorite with 2 km diameter	100 TNs
Momentum of a galaxy in galaxy collision	up to 10 ⁴⁶ Ns

Challenge 158 s

directions the two spheres take after the collision. Can you find this rule? It is particularly useful when playing billiards. We will find out later that it is *not* valid in special relativity.

Another consequence of momentum conservation is shown in Figure 62: a man is lying on a bed of nails with a large block of concrete on his stomach. Another man is hitting the concrete with a heavy sledgehammer. As the impact is mostly absorbed by the concrete, there is no pain and no danger – unless the concrete is missed. Why?

Challenge 159 s

The above definition of mass has been generalized by the physicist and philosopher Ernst Mach* in such a way that it is valid even if the two objects interact without contact,

^{*} Ernst Mach (1838 Chrlice–1916 Vaterstetten), Austrian physicist and philosopher. The *mach* unit for aeroplane speed as a multiple of the speed of sound in air (about 0.3 km/s) is named after him. He developed the so-called Mach–Zehnder interferometer; he also studied the basis of mechanics. His thoughts about mass

as long as they do so along the line connecting their positions. The mass ratio between two bodies is defined as a negative inverse acceleration ratio, thus as

$$\frac{m_2}{m_1} = -\frac{a_1}{a_2} \,, \tag{20}$$

where *a* is the acceleration of each body during the interaction. This definition has been studied in much detail in the physics community, mainly in the nineteenth century. A few points sum up the results:

- The definition of mass *implies* the conservation of total momentum $\sum mv$. Momentum conservation is *not* a separate principle. Conservation of momentum cannot be checked experimentally, because mass is defined in such a way that the principle holds.
- The definition of mass *implies* the equality of the products m_1a_1 and $-m_2a_2$. Such products are called *forces*. The equality of acting and reacting forces is not a separate principle; mass is defined in such a way that the principle holds.
- The definition of mass is *independent* of whether contact is involved or not, and whether the origin of the accelerations is due to electricity, gravitation, or other interactions.* Since the interaction does not enter the definition of mass, mass values defined with the help of the electric, nuclear or gravitational interaction all agree, as long as momentum is conserved. All known interactions conserve momentum. For some unfortunate historical reasons, the mass value measured with the electric or nuclear interactions is called the 'inertial' mass and the mass measured using gravity is called the 'gravitational' mass. As it turns out, this artificial distinction has no real meaning; this becomes especially clear when one takes an observation point that is far away from all the bodies concerned.
- The definition of mass requires observers at rest or in inertial motion.

By measuring the masses of bodies around us we can explore the science and art of experiments. An overview of devices is given in Table 18 and Figure 63. Some measurement results are listed in Table 16. We also discover the main properties of mass. It is *additive* in everyday life, as the mass of two bodies combined is equal to the sum of the two separate masses. Furthermore, mass is *continuous*; it can seemingly take any positive value. Finally, mass is *conserved*; the mass of a system, defined as the sum of the mass of all constituents, does not change over time if the system is kept isolated from the rest of the world. Mass is not only conserved in collisions but also during melting, evaporation, digestion and all other processes.

and inertia influenced the development of general relativity, and led to Mach's principle, which will appear later on. He was also proud to be the last scientist denying – humorously, and against all evidence – the existence of atoms.

^{*} As mentioned above, only *central* forces obey the relation (20) used to define mass. Central forces act between the centre of mass of bodies. We give a precise definition later. However, since all fundamental forces are central, this is not a restriction. There seems to be one notable exception: magnetism. Is the definition of mass valid in this case?

Motion Mountain – The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christoph Schiller November 1997–Mai

TABLE 16 Some measured mass values.

O B S E R V A T I O N	M a s s
Probably lightest known object: neutrino	$c. 2 \cdot 10^{-36} \mathrm{kg}$
Mass increase due to absorption of one green photon	$4.1 \cdot 10^{-36} \mathrm{kg}$
Lightest known charged object: electron	$9.10938188(72)\cdot 10^{-31}\mathrm{kg}$
Atom of argon	39.962 383 123(3) u = 66.359 1(1) yg
Lightest object ever weighed (a gold particle)	0.39 ag
Human at early age (fertilized egg)	$10^{-8} \mathrm{g}$
Water adsorbed on to a kilogram metal weight	$10^{-5} \mathrm{g}$
Planck mass	$2.2 \cdot 10^{-5} \mathrm{g}$
Fingerprint	$10^{-4} \mathrm{g}$
Typical ant	$10^{-4} \mathrm{g}$
Water droplet	1 mg
Honey bee, Apis mellifera	0.1 g
Euro coin	7.5 g
Blue whale, Balaenoptera musculus	180 Mg
Heaviest living things, such as the fungus <i>Armillaria</i> ostoyae or a large Sequoia Sequoiadendron giganteum	$10^6 \mathrm{kg}$
Heaviest train ever	99.7 · 10 ⁶ kg
Largest ocean-going ship	$400 \cdot 10^6 \text{ kg}$
Largest object moved by man (Troll gas rig)	$687.5 \cdot 10^6 \mathrm{kg}$
Large antarctic iceberg	10 ¹⁵ kg
Water on Earth	$10^{21} \mathrm{kg}$
Earth's mass	$5.98 \cdot 10^{24} \mathrm{kg}$
Solar mass	$2.0 \cdot 10^{30} \mathrm{kg}$
Our galaxy's visible mass	$3\cdot 10^{41} \text{ kg}$
Our galaxy's estimated total mass	$2 \cdot 10^{42} \mathrm{kg}$
virgo supercluster	$2 \cdot 10^{46} \mathrm{kg}$
Total mass visible in the universe	$10^{54} \mathrm{kg}$

Later we will find that in the case of mass all these properties, summarized in Table 17, are only approximate. Precise experiments show that none of them are correct.* For the moment we continue with the present, Galilean concept of mass, as we have not yet a better one at our disposal.

The definition of mass through momentum conservation implies that when an object falls, the Earth is accelerated upwards by a tiny amount. If one could measure this tiny amount, one could determine the mass of the Earth. Unfortunately, this measurement is impossible. Can you find a better way to determine the mass of the Earth?

Summarizing Table 17, the mass of a body is thus most precisely described by a *positive* real number, often abbreviated *m* or *M*. This is a direct consequence of the impenetrabil-

Challenge 161 s

^{*} In particular, in order to define mass we must be able to *distinguish* bodies. This seems a trivial requirement, but we discover that this is not always possible in nature.

Motion Mountain - The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christoph Schiller November 1997-Marc

TABLE 17 Properties of Galilean mass.

Masses	P H Y S I C A L P R O P E R T Y	M athematical name	DEFINI - TION
Can be distinguished	distinguishability	element of set	Vol. III, page 223
Can be ordered	sequence	order	Vol. IV, page 196
Can be compared	measurability	metricity	Vol. V, page 280
Can change gradually	continuity	completeness	Vol. V, page 288
Can be added	quantity of matter	additivity	
			Page 77
Beat any limit	infinity	unboundedness, openness	Vol. III, page 224
Do not change	conservation	invariance	m = const
Do not disappear	impenetrability	positivity	$m \geqslant 0$

Challenge 162 e

Challenge 163 e Vol. II, page 66 Vol. IV, page 166 ity of matter. Indeed, a *negative* (inertial) mass would mean that such a body would move in the opposite direction of any applied force or acceleration. Such a body could not be kept in a box; it would break through any wall trying to stop it. Strangely enough, negative mass bodies would still fall downwards in the field of a large positive mass (though more slowly than an equivalent positive mass). Are you able to confirm this? However, a small positive mass object would float away from a large negative-mass body, as you can easily deduce by comparing the various accelerations involved. A positive and a negative mass of the same value would stay at constant distance and spontaneously accelerate away along the line connecting the two masses. Note that both energy and momentum are conserved in all these situations.* Negative-mass bodies have never been observed. Antimatter, which will be discussed later, also has positive mass.

Is motion eternal? - Conservation of momentum

Every body continues in the state of rest or of uniform motion in a straight line except in so far as it doesn't.

Arthur Eddington**

The product p = mv of mass and velocity is called the *momentum* of a particle; it describes the tendency of an object to keep moving during collisions. The larger it is, the harder it is to stop the object. Like velocity, momentum has a direction and a magnitude: it is a vector. In French, momentum is called 'quantity of motion', a more appropriate term. In the old days, the term 'motion' was used instead of 'momentum', for example by

Page 101

Challenge 164 e

Challenge 165 s

* For more curiosities, see R. H. PRICE, *Negative mass can be positively amusing*, American Journal of Physics 61, pp. 216–217, 1993. Negative mass particles in a box would heat up a box made of positive mass while traversing its walls, and accelerating, i.e., losing energy, at the same time. They would allow one to build a *perpetuum mobile* of the second kind, i.e., a device circumventing the second principle of thermodynamics. Moreover, such a system would have no thermodynamic equilibrium, because its energy could decrease forever. The more one thinks about negative mass, the more one finds strange properties contradicting observations. By the way, what is the range of possible mass values for tachyons?

** Arthur Eddington (1882–1944), British astrophysicist.

TABLE 18 Some mass sensors.

MEASUREMENT	SENSOR	RANGE
Particle collision	speed	below 1 mg
Sense of touch	pressure sensitive cells	1 mg to 500 kg
Doppler effect on light reflected off the object	interferometer	1 mg to 100 g
Precision scales	balance, pendulum, or spring	1 pg to 10 ³ kg
Cosmonaut body mass measurement device	spring frequency	around 70 kg
Truck scales	hydraulic balance	$1 \text{ to } 60 \cdot 10^3 \text{ kg}$
Ship weight	water volume measurement	up to $500 \cdot 10^6$ kg










FIGURE 63 Mass measurement devices: a vacuum balance used in 1890 by Dmitriy Ivanovich Mendeleyev, a modern laboratory balance, a device to measure the mass of a cosmonaut in space and a truck scales (© Thinktank Trust, Mettler-Toledo, NASA Anonymous).

Newton. The conservation of momentum, relation (19), therefore expresses the conservation of motion during interactions.

Momentum is an *extensive quantity*. That means that it can be said that it *flows* from one body to the other, and that it can be *accumulated* in bodies, in the same way that



Challenge 166 s FIGURE 64 What happens in these four situations?

water flows and can be accumulated in containers. Imagining momentum as something that can be *exchanged* between bodies in collisions is always useful when thinking about the description of moving objects.

Momentum is conserved. That explains the limitations you might experience when being on a perfectly frictionless surface, such as ice or a polished, oil covered marble: you cannot propel yourself forward by patting your own back. (Have you ever tried to put a cat on such a marble surface? It is not even able to stand on its four legs. Neither are humans. Can you imagine why?) Momentum conservation also answers the puzzles of Figure 64.

Challenge 167 s

Challenge 168 s

The conservation of momentum and mass also means that teleportation ('beam me up') is impossible in nature. Can you explain this to a non-physicist?

Momentum conservation implies that momentum can be imagined to be like an invisible fluid. In an interaction, the invisible fluid is transferred from one object to another. In such transfers, the sum of fluid is always constant.

Momentum conservation implies that motion never stops; it is only *exchanged*. On the other hand, motion often 'disappears' in our environment, as in the case of a stone dropped to the ground, or of a ball left rolling on grass. Moreover, in daily life we often observe the creation of motion, such as every time we open a hand. How do these examples fit with the conservation of momentum?

It turns out that the answer lies in the microscopic aspects of these systems. A muscle only *transforms* one type of motion, namely that of the electrons in certain chemical compounds* into another, the motion of the fingers. The working of muscles is similar to that of a car engine transforming the motion of electrons in the fuel into motion of the wheels. Both systems need fuel and get warm in the process.

We must also study the microscopic behaviour when a ball rolls on grass until it stops. The disappearance of motion is called *friction*. Studying the situation carefully, we find that the grass and the ball heat up a little during this process. During friction, visible motion is transformed into heat. A striking observation of this effect for a bicycle is shown below, in Figure 214. Later, when we discover the structure of matter, it will become clear that heat is the disorganized motion of the microscopic constituents of every material. When these constituents all move in the same direction, the object as a whole moves;

Page 320

^{*} Usually adenosine triphosphate (ATP), the fuel of most processes in animals.

when they oscillate randomly, the object is at rest, but is warm. Heat is a form of motion. Friction thus only seems to be disappearance of motion; in fact it is a transformation of ordered into unordered motion.

Page 330

Despite momentum conservation, *macroscopic* perpetual motion does not exist, since friction cannot be completely eliminated.* Motion is eternal only at the microscopic scale. In other words, the disappearance and also the spontaneous appearance of motion in everyday life is an illusion due to the limitations of our senses. For example, the motion proper of every living being exists before its birth, and stays after its death. The same happens with its energy. This result is probably the closest one can get to the idea of everlasting life from evidence collected by observation. It is perhaps less than a coincidence that energy used to be called *vis viva*, or 'living force', by Leibniz and many others.

Since motion is conserved, it has no origin. Therefore, at this stage of our walk we cannot answer the fundamental questions: Why does motion exist? What is its origin? The end of our adventure is nowhere near.

More conservation - energy

When collisions are studied in detail, a second conserved quantity turns up. Experiments show that in the case of perfect, or elastic collisions – collisions without friction – the following quantity, called the *kinetic energy T* of the system, is also conserved:

$$T = \sum_{i} \frac{1}{2} m_i v_i^2 = \sum_{i} \frac{1}{2} m_i v_i^2 = \text{const}.$$
 (21)

Kinetic energy is the ability that a body has to induce change in bodies it hits. Kinetic energy thus depends on the mass and on the square of the speed v of a body. The full name 'kinetic energy' was introduced by Gustave-Gaspard Coriolis.** Some measured energy values are given in Table 19.

If the machine is well constructed, i.e., with little friction, it can take the little energy it needs for the sustenance of its motion from very subtle environmental effects. For example, in the Victoria and Albert Museum in London one can admire a beautiful clock powered by the variations of air pressure over time.

Ref. 81

Challenge 169 s Bu

Low friction means that motion takes a long time to stop. One immediately thinks of the motion of the planets. In fact, there *is* friction between the Earth and the Sun. (Can you guess one of the mechanisms?) But the value is so small that the Earth has already circled around the Sun for thousands of millions of years, and will do so for quite some time more.

Page 124

** Gustave-Gaspard Coriolis (b. 1792 Paris, d. 1843 Paris), French engineer and mathematician. He introduced the modern concepts of 'work' and of 'kinetic energy', and discovered the Coriolis effect. Coriolis also introduced the factor 1/2 in the kinetic energy T, in order that the relation $\mathrm{d}T/\mathrm{d}v = p$ would be obeyed. (Why?)

Challenge 170 s

^{*} Some funny examples of past attempts to built a *perpetual motion machine* are described in Stanislav Michel, *Perpetuum mobile*, VDI Verlag, 1976. Interestingly, the idea of eternal motion came to Europe from India, via the Islamic world, around the year 1200, and became popular as it opposed the then standard view that all motion on Earth disappears over time. See also the web.archive.org/web/20040812085618/http://www.geocities.com/mercutio78_99/pmm.html and the www.lhup.edu/~dsimanek/museum/unwork.htm websites. The conceptual mistake made by eccentrics and used by crooks is always the same: the hope of overcoming friction. (In fact, this applied only to the perpetual motion machines of the second kind; those of the first kind – which are even more in contrast with observation – even try to generate energy from nothing.)



FIGURE 65 Robert Mayer (1814–1878).

▶ (Physical) *energy* is the measure of the ability to generate motion.

A body has a lot of energy if it has the ability to move many other bodies. Energy is a number; energy, in contrast to momentum, has no direction. The total momentum of two equal masses moving with opposite velocities is zero; but their total energy is not, and it increases with velocity. Energy thus also measures motion, but in a different way than momentum. Energy measures motion in a more global way. An equivalent definition is the following:

Energy is the ability to perform work.

Energy is the ability to perform work. Here, the physical concept of work is just the precise version of what is meant by work in everyday life.*

Another, equivalent definition of energy will become clear later:

▶ Energy is what can be transformed into heat.

Energy is a word taken from ancient Greek; originally it was used to describe character, and meant 'intellectual or moral vigour'. It was taken into physics by Thomas Young (1773–1829) in 1807 because its literal meaning is 'force within'. (The letters E, W, A and several others are also used to denote energy.)

Do not be surprised if you do not grasp the difference between momentum and energy straight away: physicists took about two centuries to figure it out! For some time they even insisted on using the same word for both, and often they didn't know which situation required which concept. So you are allowed to take some time to get used to it.

Both energy and momentum measure how systems change. Momentum tells how systems change *over distance*, energy measures how systems change *over time*. Momentum is needed to compare motion here and there. Energy is needed to compare motion now and later.

One way to explore the difference between energy and momentum is to think about the following challenges. Is it more difficult to stop a running man with mass m and speed v, or one with mass m/2 and speed v? You may want to ask a rugby-playing friend for confirmation.

Challenge 171 e

^{* (}Physical) *work* is the product of force and distance in direction of the force. In other words, work is the *scalar product* of force and distance.

TABLE 19 Some measured energy values.

OBSERVATION	ENERGY
Average kinetic energy of oxygen molecule in air	6 zJ
Green photon energy	0.37 aJ
X-ray photon energy	1 fJ
γ photon energy	1 pJ
Highest particle energy in accelerators	0.1 μJ
Kinetic energy of a flying mosquito	0.2 μJ
Comfortably walking human	20 J
Flying arrow	50 J
Right hook in boxing	50 J
Energy in torch battery	1 kJ
Energy in explosion of 1 g TNT	4.1 kJ
Energy of 1 kcal	4.18 kJ
Flying rifle bullet	10 kJ
One gram of fat	38 kJ
One gram of gasoline	44 kJ
Apple digestion	0.2 MJ
Car on highway	0.3 to 1 MJ
Highest laser pulse energy	1.8 MJ
Lightning flash	up to 1 GJ
Planck energy	2.0 GJ
Small nuclear bomb (20 ktonne)	84 TJ
Earthquake of magnitude 7	2 PJ
Largest nuclear bomb (50 Mtonne)	210 PJ
Impact of meteorite with 2 km diameter	1 EJ
Yearly machine energy use	420 EJ
Rotation energy of Earth	$2 \cdot 10^{29} \text{ J}$
Supernova explosion	$10^{44} \mathrm{J}$
Gamma ray burst	up to 10 ⁴⁷ J
Energy content $E = mc^2$ of Sun's mass	$1.8 \cdot 10^{47} \text{ J}$
Energy content of Galaxy's central black hole	$4\cdot 10^{53}$ J

Another distinction is illustrated by athletics: the *real* long jump world record, almost 10 m, is still kept by an athlete who in the early twentieth century ran with two weights in his hands, and then threw the weights behind him at the moment he took off. Can you explain the feat?

Challenge 172 s

When a car travelling at 100 m/s runs head-on into a parked car of the same kind and Challenge 173 s make, which car receives the greatest damage? What changes if the parked car has its brakes on?

> To get a better feeling for energy, here is an additional approach. The world consumption of energy by human machines (coming from solar, geothermal, biomass, wind, nu-

Ref. 82

clear, hydro, gas, oil, coal, or animal sources) in the year 2000 was about 420 EJ,* for a world population of about 6000 million people. To see what this energy consumption means, we translate it into a personal power consumption; we get about 2.2 kW. The watt W is the unit of power, and is simply defined as 1 W = 1 J/s, reflecting the definition of (physical) power as energy used per unit time. (The precise wording is: power is energy flowing per time through a defined closed surface.) As a working person can produce mechanical work of about 100 W, the average human energy consumption corresponds to about 22 humans working 24 hours a day. (See Table 20 for some power values found in nature, and Table 21 for some measurement devices.) In particular, if we look at the energy consumption in First World countries, the average inhabitant there has machines working for them equivalent to several hundred 'servants'. Can you point out some of these machines?

Challenge 174 s

Kinetic energy is thus not conserved in everyday life. For example, in non-elastic collisions, such as that of a piece of chewing gum hitting a wall, kinetic energy is lost. *Friction* destroys kinetic energy. At the same time, friction produces heat. It was one of the important conceptual discoveries of physics that *total* energy is conserved if one includes the discovery that heat is a form of energy. Friction is thus in fact a process transforming kinetic energy, i.e., the energy connected with the motion of a body, into heat. On a microscopic scale, energy is conserved.** Indeed, without energy conservation, the concept of time would not be definable. We will show this connection shortly.

In summary, in addition to mass and momentum, everyday linear motion also conserves energy. To discover the last conserved quantity, we explore another type of motion: rotation.

THE CROSS PRODUCT, OR VECTOR PRODUCT

The discussion of rotation is easiest if we introduce an additional way to multiply vectors. This new product between two vectors a and b is called the *cross product* or *vector product* $a \times b$.

The result of the vector product is another vector; thus it differs from the *scalar* product, whose result is a scalar, i.e., a number. The result of the vector product is that vector that is orthogonal to both vectors to be multiplied, whose orientation is given by the *right-hand rule*, and whose length is given by $ab \sin \triangleleft (a, b)$, i.e., by the surface area of the parallelogram spanned by the two vectors.

The definition implies that the cross product vanishes if and only if the vectors are parallel. From the definition you can also show that the vector product has the properties

Page 383

^{*} For the explanation of the abbreviation E, see Appendix B.

^{**} In fact, the conservation of energy was stated in its full generality in public only in 1842, by Julius Robert Mayer. He was a medical doctor by training, and the journal *Annalen der Physik* refused to publish his paper, as it supposedly contained 'fundamental errors'. What the editors called errors were in fact mostly – but not only – contradictions of their prejudices. Later on, Helmholtz, Thomson-Kelvin, Joule and many others acknowledged Mayer's genius. However, the first to have stated energy conservation in its modern form was the French physicist Sadi Carnot (1796–1832) in 1820. To him the issue was so clear that he did not publish the result. In fact he went on and discovered the second 'law' of thermodynamics. Today, energy conservation, also called the first 'law' of thermodynamics, is one of the pillars of physics, as it is valid in all its domains.

TABLE 20 Some measured power values.

O B S E R V A T I O N	Power
Radio signal from the Galileo space probe sending from Jupiter	10 zW
Power of flagellar motor in bacterium	0.1 pW
Power consumption of a typical cell	1 pW
sound power at the ear at hearing threshold	2.5 pW
CR-R laser, at 780 nm	40-80 mW
Sound output from a piano playing fortissimo	$0.4\mathrm{W}$
Dove (0.16 kg) basal metabolic rate	0.97 W
Rat (0.26 kg) basal metabolic rate	1.45 W
Pigeon (0.30 kg) basal metabolic rate	1.55 W
Hen (2.0 kg) basal metabolic rate	4.8 W
Incandescent light bulb light output	1 to 5 W
Dog (16 kg) basal metabolic rate	20 W
Sheep (45 kg) basal metabolic rate	50 W
Woman (60 kg) basal metabolic rate	68 W
Man (70 kg) basal metabolic rate	87 W
Incandescent light bulb electricity consumption	25 to 100 W
A human, during one work shift of eight hours	100 W
Cow (400 kg) basal metabolic rate	266 W
One horse, for one shift of eight hours	300 W
Steer (680 kg) basal metabolic rate	411 W
Eddy Merckx, the great bicycle athlete, during one hour	500 W
Metric horse power power unit $(75 \text{ kg} \cdot 9.81 \text{ m/s}^2 \cdot 1 \text{ m/s})$	735.5 W
British horse power power unit	745.7 W
Large motorbike	100 kW
Electrical power station output	0.1 to 6 GW
World's electrical power production in 2000 Ref. 82	450 GW
Power used by the geodynamo	200 to 500 GW
Input on Earth surface: Sun's irradiation of Earth Ref. 83	0.17 EW
Input on Earth surface: thermal energy from inside of the Earth	32 TW
Input on Earth surface: power from tides (i.e., from Earth's rotation)	3 TW
Input on Earth surface: power generated by man from fossil fuels	8 to 11 TW
Lost from Earth surface: power stored by plants' photosynthesis	40 TW
World's record laser power	1 PW
Output of Earth surface: sunlight reflected into space	0.06 EW
Output of Earth surface: power radiated into space at 287 K	0.11 EW
Peak power of the largest nuclear bomb	5 YW
Sun's output	384.6 YW
Maximum power in nature, $c^5/4G$	$9.1 \cdot 10^{51} \mathrm{W}$

75 to 2000 W 75 to 2000 W 20 to 10 000 W up to 1 MW up to 10 GW up to 1 MW up to a few μJ/ns

RANGE

TABLE 21 Some power sensors.

Heart beat as power meter

Electricity meter at home

Power meter for car engine

Calorimeter for chemical reactions

MEASUREMENT

Fitness power meter

Laser power meter

Calorimeter for particles light detector

Challenge 175 e

$$a \times b = -b \times a , \quad a \times (b+c) = a \times b + a \times c ,$$

$$\lambda a \times b = \lambda(a \times b) = a \times \lambda b , \quad a \times a = 0 ,$$

$$a(b \times c) = b(c \times a) = c(a \times b) , a \times (b \times c) = b(ac) - c(ab) ,$$

$$(a \times b)(c \times d) = a(b \times (c \times d)) = (ac)(bd) - (bc)(ad) ,$$

$$(a \times b) \times (c \times d) = c((a \times b)d) - d((a \times b)c) ,$$

$$a \times (b \times c) + b \times (c \times a) + c \times (a \times b) = 0 .$$
(22)

Vol. IV, page 205

The vector product exists only in vector spaces with three dimensions. We will explore more details on this connection later on.

SENSOR

deformation sensor and clock

piezoelectric sensor

rotating aluminium disc

electromagnetic brake

photoelectric effect in

temperature sensor

semiconductor

The vector product is useful to describe systems that *rotate* – and (thus) also systems with magnetic forces. The main reason for the usefulness is that the motion of an orbiting body is always perpendicular both to the axis and to the shortest line that connects the body with the axis.

Challenge 176 e

Confirm that the best way to calculate the vector product $\mathbf{a} \times \mathbf{b}$ component by component is given by the determinant

$$\mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{e}_{x} & a_{x} & b_{x} \\ \mathbf{e}_{y} & a_{y} & b_{y} \\ \mathbf{e}_{z} & a_{z} & b_{z} \end{vmatrix} \quad \text{or, sloppily} \quad \mathbf{a} \times \mathbf{b} = \begin{vmatrix} + & - & + \\ a_{x} & a_{y} & a_{z} \\ b_{x} & b_{y} & b_{z} \end{vmatrix} . \tag{23}$$

This is easy to remember and easy to perform, both with letters and with numerical values. (Here, e_x is the unit basis vector in the x direction.) Written out, it is equivalent to the relation

$$\mathbf{a} \times \mathbf{b} = (a_{y}b_{z} - b_{y}a_{z}, b_{x}a_{z} - a_{x}b_{z}, a_{x}b_{y} - b_{x}a_{y})$$
(24)

which is harder to remember, though.

Show that the parallelepiped spanned by three arbitrary vectors a, b and c has the volume $V = c (a \times b)$. Show that the *pyramid* or *tetrahedron* formed by the same three vectors has one sixth of that volume.

Challenge 177 e

TABLE 22 Some measured rotation frequencies.

O B S E R V A T I O N	ANGULAR VELOCITY $\omega = 2\pi/T$
Galactic rotation	2π · 0.14 · 10 ⁻¹⁵ /s = $2\pi / (220 \cdot 10^6 \text{ a})$
Average Sun rotation around its axis	$2\pi \cdot 3.8 \cdot 10^{-7} / s = 2\pi / 30 d$
Typical lighthouse	$2\pi \cdot 0.08/s$
Pirouetting ballet dancer	$2\pi \cdot 3/s$
Ship's diesel engine	$2\pi \cdot 5/s$
Helicopter rotor	$2\pi \cdot 5.3/s$
Washing machine	up to $2\pi \cdot 20/s$
Bacterial flagella	$2\pi \cdot 100/s$
Fast CD recorder	up to $2\pi \cdot 458/s$
Racing car engine	up to $2\pi \cdot 600/s$
Fastest turbine built	$2\pi \cdot 10^3/\mathrm{s}$
Fastest pulsars (rotating stars)	up to at least $2\pi \cdot 716/s$
Ultracentrifuge	$> 2\pi \cdot 2 \cdot 10^3 / \mathrm{s}$
Dental drill	up to $2\pi \cdot 13 \cdot 10^3 / s$
Technical record	$2\pi \cdot 333 \cdot 10^3 / \mathrm{s}$
Proton rotation	$2\pi \cdot 10^{20}/\mathrm{s}$
Highest possible, Planck angular velocity	$2\pi \cdot 10^{35}/\mathrm{s}$

ROTATION AND ANGULAR MOMENTUM

Rotation keeps us alive. Without the change of day and night, we would be either fried or frozen to death, depending on our location on our planet. But rotation appears in many other settings, as Table 22 shows. A short exploration of rotation is thus appropriate.

All objects have the ability to rotate. We saw before that a body is described by its reluctance to move; similarly, a body also has a reluctance to turn. This quantity is called its *moment of inertia* and is often abbreviated Θ – pronounced 'theta.' The speed or rate of rotation is described by *angular velocity*, usually abbreviated ω – pronounced 'omega'. A few values found in nature are given in Table 22.

The observables that describe rotation are similar to those describing linear motion, as shown in Table 24. Like mass, the moment of inertia is defined in such a way that the sum of *angular momenta* L – the product of moment of inertia and angular velocity – is conserved in systems that do not interact with the outside world:

$$\sum_{i} \Theta_{i} \boldsymbol{\omega}_{i} = \sum_{i} \boldsymbol{L}_{i} = \text{const}.$$
 (25)

In the same way that the conservation of linear momentum defines mass, the conservation of angular momentum defines the moment of inertia.

The moment of inertia can be related to the mass and shape of a body. If the body is






FIGURE 66 Angular momentum and other quantities for a point particle in circular motion, and the two versions of the right-hand rule.

imagined to consist of small parts or mass elements, the resulting expression is

$$\Theta = \sum_{n} m_n r_n^2 \,, \tag{26}$$

where r_n is the distance from the mass element m_n to the axis of rotation. Can you confirm the expression? Therefore, the moment of inertia of a body depends on the chosen axis of rotation. Can you confirm that this is so for a brick?

In contrast to the case of mass, there is no conservation of the moment of inertia. The value of the moment of inertia depends on the location of the axis used for its definition. For each axis direction, one distinguishes an *intrinsic* moment of inertia, when the axis passes through the centre of mass of the body, from an *extrinsic* moment of inertia, when it does not.* In the same way, one distinguishes intrinsic and extrinsic angular momenta. (By the way, the *centre of mass* of a body is that imaginary point which moves straight during vertical fall, even if the body is rotating. Can you find a way to determine its location for a specific body?)

We now define the rotational energy as

$$E_{\rm rot} = \frac{1}{2} \Theta \ \omega^2 = \frac{L^2}{2\Theta} \ . \tag{28}$$

The expression is similar to the expression for the kinetic energy of a particle. Can you guess how much larger the rotational energy of the Earth is compared with the yearly electricity usage of humanity? In fact, if you could find a way to harness this energy, you would become famous. For undistorted rotated objects, rotational energy is conserved.

Every object that has an orientation also has an intrinsic angular momentum. (What about a sphere?) Therefore, point particles do not have intrinsic angular momenta – at

* Extrinsic and intrinsic moment of inertia are related by

$$\Theta_{\rm ext} = \Theta_{\rm int} + md^2 \,, \tag{27}$$

where *d* is the distance between the centre of mass and the axis of extrinsic rotation. This relation is called *Steiner's parallel axis theorem*. Are you able to deduce it?

Challenge 178 e

Challenge 179 s

Challenge 181 s

Challenge 180 s

Challenge 182 s

Challenge 183 s

TABLE 23 Some measured angular momentum values.

O B S E R V A T I O N	ANGULAR MOMEN- TUM
Smallest observed in nature $(\hbar/2)$ – applies to the z-component of most fermions	$0.53 \cdot 10^{-34} \text{Js}$
Spinning top	$5\cdot 10^{-6}$ Js
CD (compact disk) playing	0.029 Js
Walking man (around body axis)	c. 4 Js
Dancer in a pirouette	5 Js
Typical car wheel at 30 m/s	10 Js
Typical wind generator at 12 m/s (6 Beaufort)	10^4 Js
Atmosphere	1 to $1 \cdot 10^{26} \mathrm{kg} \mathrm{m}^2/\mathrm{s}$
Earth around its axis	$7.1 \cdot 10^{33} \text{kg m}^2/\text{s}$
Moon around Earth	$2.9 \cdot 10^{34} \mathrm{kg} \mathrm{m}^2/\mathrm{s}$
Earth around Sun	$2.7 \cdot 10^{40} \mathrm{kg} \mathrm{m}^2/\mathrm{s}$
Sun around its axis	$1.1 \cdot 10^{42} \text{kg m}^2/\text{s}$
Jupiter around Sun	$1.9 \cdot 10^{43} \text{kg m}^2/\text{s}$
Solar system around Sun	$3.2 \cdot 10^{43} \mathrm{kg} \mathrm{m}^2/\mathrm{s}$
Milky Way	$10^{68} \mathrm{kg} \mathrm{m}^2/\mathrm{s}$
All masses in the universe	0 (within measurement error)

QUANTITY	LINEAR M	OTION	ROTATIONAL MOTION	
State	time	t	time	t
	position	x	angle	φ
	momentum	$p = m\mathbf{v}$	angular momentum	$L = \Theta \omega$
	energy	$mv^2/2$	energy	$\Theta \omega^2/2$
Motion	velocity	\boldsymbol{v}	angular velocity	ω
	acceleration	a	angular acceleration	α
Reluctance to move	mass	m	moment of inertia	Θ
Motion change	force	m a	torque	$\Theta \alpha$

TABLE 24 Correspondence between linear and rotational motion.




FIGURE 67 Can the ape reach the banana?

Page 134

FIGURE 68 How a snake turns itself around its axis.

least in first approximation. (This conclusion will change in quantum theory.) The *extrinsic* angular momentum L of a point particle is given by

$$L = r \times p = \frac{2a(T)m}{T}$$
 so that $L = rp = \frac{2A(T)m}{T}$ (29)

where p is the momentum of the particle, a(T) is the surface swept by the position vector r of the particle during time T. The angular momentum thus points along the rotation axis, following the right-hand rule, as shown in Figure 66. A few observed values are given in Table 23.

A body can rotate simultaneously about several axes. The film of Figure 94 shows an example: The top rotates around its body axis and around the vertical at the same time. A detailed exploration shows that the exact rotation of the top is given by the vector sum of these two rotations. Rotations thus are vectors.

As in the case of linear motion, rotational energy and angular momentum are not

The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christoph Schiller November 1997–March 2012

always conserved in the macroscopic world: rotational energy can change due to friction, and angular momentum can change due to external forces (torques). However, for closed (undisturbed) systems both quantities are always conserved. In particular, on a microscopic scale, most objects are undisturbed, so that conservation of rotational energy and angular momentum is especially obvious there.

Angular momentum is conserved. This statement is valid for any axis of a physical system, provided that external forces (torques) play no role. To make the point, Jean-Marc Lévy-Leblond poses the problem of Figure 67. Can the ape reach the banana without leaving the plate, assuming that the plate on which the ape rests can turn around the axis without friction?

We note that the effects of rotation are the same as for acceleration. Therefore, many sensors for rotation are the same as the acceleration sensors we explored above. But a few sensors for rotation are fundamentally new. In particular, we will meet the gyroscope shortly.

On a frictionless surface, as approximated by smooth ice or by a marble floor covered by a layer of oil, it is impossible to move forward. In order to move, we need to push *against* something. Is this also the case for rotation?

Surprisingly, it is possible to turn even *without* pushing against something. You can check this on a well-oiled rotating office chair: simply rotate an arm above the head. After each turn of the hand, the orientation of the chair has changed by a small amount. Indeed, conservation of angular momentum and of rotational energy do not prevent bodies from changing their orientation. Cats learn this in their youth. After they have learned the trick, if they are dropped legs up, they can turn themselves in such a way that they always land feet first. Snakes also know how to rotate themselves, as Figure 68 shows. During the Olympic Games one can watch board divers and gymnasts perform similar tricks. Rotation is thus different from translation in this aspect. (Why?)

ROLLING WHEELS

Rotation is an interesting phenomenon in many ways. A rolling wheel does *not* turn around its axis, but around its point of contact. Let us show this.

A wheel of radius R is *rolling* if the speed of the axis v_{axis} is related to the angular velocity ω by

$$\omega = \frac{v_{\text{axis}}}{R} \ . \tag{30}$$

For any point P on the wheel, with distance r from the axis, the velocity $v_{\rm P}$ is the sum of the motion of the axis and the motion around the axis. Figure 69 shows that $v_{\rm P}$ is orthogonal to d, the distance between the point P and the contact point of the wheel. The figure also shows that the length ratio between $v_{\rm P}$ and d is the same as between $v_{\rm axis}$ and R. As a result, we can write

$$\boldsymbol{v}_{\mathrm{P}} = \boldsymbol{\omega} \times \boldsymbol{d} , \qquad (31)$$

which shows that a rolling wheel does indeed rotate about its point of contact with the ground.

Surprisingly, when a wheel rolls, some points on it move towards the wheel's axis,

Challenge 184 s

Page 82

Page 128

Ref. 84

Challenge 185 d

Challenge 186 e




FIGURE 69 The velocities and unit vectors for a rolling wheel.

FIGURE 70 A simulated photograph of a rolling wheel with spokes.

Challenge 187 s Ref. 85 Ref. 86

some stay at a *fixed* distance and others move *away* from it. Can you determine where these various points are located? Together, they lead to an interesting pattern when a rolling wheel with spokes, such as a bicycle wheel, is photographed, as show in Figure 70.

With these results you can tackle the following beautiful challenge. When a turning bicycle wheel is deposed on a slippery surface, it will slip for a while, then slip and roll, and finally roll only. How does the final speed depend on the initial speed and on the friction?

Challenge 188 d

HOW DO WE WALK?

Golf is a good walk spoiled.

Why do we move our arms when walking or running? To save energy or to be graceful? In fact, whenever a body movement is performed with as little energy as possible, it is natural and graceful. This correspondence can indeed be taken as the actual definition of grace. The connection is common knowledge in the world of dance; it is also a central aspect of the methods used by actors to learn how to move their bodies as beautifully as possible.

To convince yourself about the energy savings, try walking or running with your arms fixed or moving in the opposite direction to usual: the effort required is considerably higher. In fact, when a leg is moved, it produces a torque around the body axis which has to be counterbalanced. The method using the least energy is the swinging of arms. Since the arms are lighter than the legs, they must move further from the axis of the body, to compensate for the momentum; evolution has therefore moved the attachment of the arms, the shoulders, farther apart than those of the legs, the hips. Animals on two legs but no arms, such as penguins or pigeons, have more difficulty walking; they have to move their whole torso with every step.

Which muscles do most of the work when walking, the motion that experts call *gait*? In 1980, Serge Gracovetsky found that in human gait a large fraction of the power comes

Ref. 20




FIGURE 71 The measured motion of a walking human (© Ray McCoy).

from the muscles along the *spine*, not from those of the legs. (Indeed, people without legs are also able to walk. However, a number of muscles in the legs must work in oder to walk normally.) When you take a step, the lumbar muscles straighten the spine; this automatically makes it turn a bit to one side, so that the knee of the leg on that side automatically comes forward. When the foot is moved, the lumbar muscles can relax, and then straighten again for the next step. In fact, one can experience the increase in tension in the back muscles when walking without moving the arms, thus confirming where the human engine is located.

Human legs differ from those of apes in a fundamental aspect: humans are able to run. In fact the whole human body has been optimized for running, an ability that no other primate has. The human body has shed most of its hair to achieve better cooling, has evolved the ability to run while keeping the head stable, has evolved the right length of arms for proper balance when running, and even has a special ligament in the back that works as a shock absorber while running. In other words, running is the most human of all forms of motion.

CURIOSITIES AND FUN CHALLENGES ABOUT CONSERVATION AND ROTATION

It is a mathematical fact that the casting of this pebble from my hand alters the centre of gravity of the universe.

Thomas Carlyle,* Sartor Resartus III.

Walking is a source of many physics problems. When climbing a mountain, the most energy-effective way is not always to follow the steepest ascent; indeed, for steep slopes, zig-zagging is more energy efficient. Why? And can you estimate the slope angle at which this will happen?

Death is a physical process; let us explore it. In general, animals have a lifespan T that

* Thomas Carlyle (1797–1881), Scottish essayist. Do you agree with the quotation?

Challenge 189 e

Ref. 88

Challenge 191 s

Challenge 190 s

Ref. 89

Challenge 192 s

Challenge 193 s

Ref. 90




FIGURE 72 Is it safe to let the cork go?

scales with fourth root of their mass M. In other terms, $T = M^{1/4}$. This valid from bacteria to insects to blue whales. Animals also have a power consumption per mass, or *metabolic rate* per mass, that scales with the *inverse* fourth root. We conclude that death occurs for all animals when a certain fixed energy consumption per mass has been achieved. This is indeed the case; death occurs for most animals when they have consumed around 1 GJ/kg. (But quite a bit later for humans.) This surprisingly simple result is valid, *on average*, for all known animals.

Note that the argument is only valid when *different* species are compared. The dependence on mass is *not* valid when specimen of the same species are compared. (You cannot live longer by eating less.)

In short, animals die after they metabolized 1 GJ/kg. In other words, once we ate all the calories we were designed for, we die.

* *

A car at a certain speed uses 7 litres of gasoline per 100 km. What is the combined air and rolling resistance? (Assume that the engine has an efficiency of 25%.)

* *

A cork is attached to a thin string a metre long. The string is passed over a long rod held horizontally, and a wine glass is attached at the other end. If you let go the cork in Figure 72, nothing breaks. Why not? And what happens exactly?

* *

In 1907, Duncan MacDougalls, a medical doctor, measured the weight of dying people, in the hope to see whether death leads to a mass change. He found a sudden decrease between 10 and 20 g at the moment of death. He attributed it to the soul exiting the body. Can you find a more satisfying explanation?

* *

It is well known that the weight of a one-year old child depends on whether the it wants to be carried or whether it wants to reach the floor. Does this contradict mass conservation?

Challenge 195 e

Challenge 194 s

* *

The Earth's crust is less dense (2.7 kg/l) than the Earth's mantle (3.1 kg/l) and floats on it.




FIGURE 73 A simple model for continents and mountains.

Challenge 196 s

As a result, the lighter crust below a mountain ridge must be much deeper than below a plain. If a mountain rises 1 km above the plain, how much deeper must the crust be below it? The simple block model shown in Figure 73 works fairly well; first, it explains why, near mountains, measurements of the deviation of free fall from the vertical line lead to so much lower values than those expected without a deep crust. Later, sound measurements have confirmed directly that the continental crust is indeed thicker beneath mountains.

* *

Challenge 197 e

All homogeneous cylinders roll down an inclined plane in the same way. True or false? And what about spheres? Can you show that spheres roll faster than cylinders?

* *

Challenge 198 e

Take two cans of the same size and weight, one full of ravioli and one full of peas. Which one rolls faster on an inclined plane?

* *

Another difference between matter and images: matter smells. In fact, the nose is a matter sensor. The same can be said of the tongue and its sense of taste.

* *

Challenge 199 e

Take a pile of coins. You can push out the coins, starting with the one at the bottom, by shooting another coin over the table surface. The method also helps to visualize two-dimensional momentum conservation.

* *

In early 2004, two men and a woman earned £1.2 million in a single evening in a London casino. They did so by applying the formulae of Galilean mechanics. They used the method pioneered by various physicists in the 1950s who built various small computers that could predict the outcome of a roulette ball from the initial velocity imparted by the croupier. In the case in Britain, the group added a laser scanner to a smart phone that measured the path of a roulette ball and predicted the numbers where it would arrive. In this way, they increased the odds from 1 in 37 to about 1 in 6. After six months of investigations, Scotland Yard ruled that they could keep the money they won.

In fact around the same time, a few people earned around 400 000 euro over a few

Ref. 91




FIGURE 74 A well-known toy.



FIGURE 75 An elastic collision that seems not to obey energy conservation.

weeks by using the same method in Germany, but with no computer at all. In certain casinos, machines were throwing the roulette ball. By measuring the position of the zero to the incoming ball with the naked eye, these gamblers were able to increase the odds of the bets they placed during the last allowed seconds and thus win a considerable sum purely through fast reactions.

* *

The toy of Figure 74 shows interesting behaviour: when a number of spheres are lifted and dropped to hit the resting ones, the same number of spheres detach on the other side, whereas the previously dropped spheres remain motionless. At first sight, all this seems to follow from energy and momentum conservation. However, energy and momentum conservation provide only two equations, which are insufficient to explain or determine the behaviour of five spheres. Why then do the spheres behave in this way? And why do they all swing in phase when a longer time has passed?

Challenge 200 d

Ref. 92

* *

A surprising effect is used in home tools such as hammer drills. We remember that when a small ball elastically hits a large one at rest, both balls move after the hit, and the small one obviously moves faster than the large one. Despite this result, when a short cylinder hits a long one of the same diameter and material, but with a length that is some *integer* multiple of that of the short one, something strange happens. After the hit, the small cylinder remains almost at rest, whereas the large one moves, as shown in Figure 75. Even though the collision is elastic, conservation of energy seems not to hold in this case. (In fact this is the reason that demonstrations of elastic collisions in schools are always performed with spheres.) What happens to the energy?

Challenge 201 d



FIGURE 76 The centre of mass defines stability.




FIGURE 77 How does the ladder fall?

* *

Does a wall get a stronger jolt when it is hit by a ball rebounding from it or when it is hit by a ball that remains stuck to it?

* *

Housewives know how to extract a cork of a wine bottle using a cloth. Can you imagine how? They also know how to extract the cork with the cloth if the cork has fallen inside the bottle. How?

* *

The sliding ladder problem, shown schematically in Figure 77, asks for the detailed motion of the ladder over time. The problem is more difficult than it looks, even if friction is not taken into account. Can you say whether the lower end always touches the floor?

* *

A homogeneous ladder of length 5 m and mass 30 kg leans on a wall. The angle is 30°; the static friction coefficient on the wall is negligible, and on the floor it is 0.3. A person of mass 60 kg climbs the ladder. What is the maximum height the person can climb before the ladder starts sliding? This and many puzzles about ladders can be found on www. mathematische-basteleien.de/leiter.htm.

* *

Ref. 93 A common fly on the stern of a 30 000 ton ship of 100 m length tilts it by less than the diameter of an atom. Today, distances that small are easily measured. Can you think of at least two methods, one of which should not cost more than 2000 euro?

* *

Is the image of three stacked spinning tops shown in Figure 78 a true photograph, showing a real observation, or is it the result of digital composition, showing an impossible situation?

Challenge 206 ny

Challenge 204 ny





FIGURE 78 Is this a possible situation or is it a fake photograph? (© Wikimedia)

* *

Challenge 207 s How does the kinetic energy of a rifle bullet compare to that of a running man?

* *

Challenge 208 s What happens to the size of an egg when one places it in a jar of vinegar for a few days?

* *

What is the amplitude of a pendulum oscillating in such a way that the absolute value of the challenge 209 s its acceleration at the lowest point and at the return point are equal?

* *

Can you confirm that the value of the acceleration of a drop of water falling through vapour is g/7?

* *

You have two hollow spheres: they have the same weight, the same size and are painted in the same colour. One is made of copper, the other of aluminium. Obviously, they fall with the same speed and acceleration. What happens if they both roll down a tilted plane?

Challenge 211 ny

* *

Challenge 212 ny What is the shape of a rope when rope jumping?

* *

Challenge 213 s How can you determine the speed of a rifle bullet with only a scale and a metre stick?

* *

Why does a gun make a hole in a door but cannot push it open, in exact contrast to what Challenge 214 e a finger can do?



FIGURE 79 A commercial clock that needs no special energy source, because it takes its energy from the environment (© Jaeger-LeCoultre).

* *

Challenge 215 s What is the curve described by the mid point of a ladder sliding down a wall?

* *

A high-tech company, see www.enocean.com, sells electric switches for room lights that have no cables and no power cell (battery). You can glue such a switch to the centre of a window pane. How is this possible?

* *

For over 50 years, a famous Swiss clock maker is selling table clocks with a rotating pendulum that need no battery and no manual rewinding, as they take up energy from the environment. A specimen is shown in Figure 79. Can you imagine how this clock works?

Challenge 217 s

Challenge 216 s

* *

Ship lifts, such as the one shown in Figure 80, are impressive machines. How does the Weight of the lift change when the ships enters?

* *

How do you measure the mass of a ship?





FIGURE 80 The spectacular ship lift at Strépy-Thieux in Belgium. What engine power is needed to lift a ship, if the right and left lifts were connected by ropes or by a hydraulic system? (© Jean-Marie Hoornaert)



FIGURE 81 The famous Celtic wobble stone and a version made with a spoon (© Ed Keath).

All masses are measured by comparing them, directly or indirectly, to the standard kilogram in Sèvres near Paris. Since a few years, there is the serious doubt that the standard kilogram is losing weight, possibly through outgassing, with an estimated rate of around 0.5 µg/a. This is an awkward situation, and there is a vast, world-wide effort to find a better definition of the kilogram. Such an improved definition must be simple, precise, and avoid trips to Sèvres. No such alternative has been defined yet.

Which engine is more efficient: a moped or a human on a bicycle?

Both mass and moment of inertia can be defined and measured both with and without

Challenge 219 e

contact. Can you do so?

Figure 81 shows the so-called *Celtic wobble stone*, also called *anagyre* or *rattleback*, a stone that starts rotating on a plane surface when it is put into up-and-down oscillation. The size can vary between a few centimetres and a few metres. By simply bending a spoon one can realize a primitive form of this strange device, if the bend is not completely symmetrical. The rotation is always in the same direction. If the stone is put into rotation in the wrong direction, after a while it stops and starts rotating in the other sense! Can you explain the effect that seems to contradict the conservation of angular momentum?

Challenge 220 d

SUMMARY ON CONSERVATION

The gods are not as rich as one might think: what they give to one, they take away from the

Antiquity

We have encountered four conservation principles that are valid for closed systems in everyday life:

- conservation of total linear momentum,
- conservation of total angular momentum,
- conservation of total energy,
- conservation of total mass.

None of these conservation laws applies to motion of images.

These conservation principles are among the great results in science. They limit the surprises that nature can offer: conservation means that linear momentum, angular momentum, and mass-energy can neither be created from nothing, nor can they disappear into nothing. Conservation limits creation. The above quote, almost blasphemous, expresses this idea.

Page 241

Later on we will find out that these results could have been deduced from three simple observations: closed systems behave the same independently of where they are, in what direction they are oriented and of the time at which they are set up. Motion is universal. In more abstract and somewhat more general terms, physicists like to say that all conservation principles are consequences of the *invariances*, or *symmetries*, of nature.

Later on, the theory of special relativity will show that energy and mass are conserved only when taken together. Many adventures still await us.



FROM THE ROTATION OF THE EARTH TO THE RELATIVITY OF MOTION

Eppur si muove!

Ref. 94

s the Earth rotating? The search for definite answers to this question gives an nteresting cross section of the history of classical physics. Around the year 265 BCE, n Samos, the Greek thinker Aristarchus maintained that the Earth rotates. He had measured the parallax of the Moon (today known to be up to 0.95°) and of the Sun (today known to be 8.8').** The parallax is an interesting effect; it is the angle describing the difference between the directions of a body in the sky when seen by an observer on the surface of the Earth and when seen by a hypothetical observer at the Earth's centre. (See Figure 82.) Aristarchus noticed that the Moon and the Sun wobble across the sky, and this wobble has a period of 24 hours. He concluded that the Earth rotates. It seems that Aristarchus received death threats for his result.

Aristarchus' observation yields a more powerful argument than the trails of the stars shown in Figure 83. Can you explain why?

Measurements of the aberration of light also show the rotation of the Earth; it can be detected with a telescope while looking at the stars. The aberration is a change of the expected light direction, which we will discuss shortly. At the Equator, Earth rotation adds an angular deviation of 0.32 ', changing sign every 12 hours, to the aberration due to the motion of the Earth around the Sun, about 20.5 '. In modern times, astronomers have found a number of additional proofs, but none is accessible to the man on the street.

Furthermore, the measurements showing that the Earth is not a sphere, but is *flat*tened at the poles, confirmed the rotation of the Earth. Figure 84 illustrates the situation. Again, however, this eighteenth century measurement by Maupertuis*** is not accessible

Challenge 221 e

Vol. II, page 16

^{* &#}x27;And yet she moves' is the sentence about the Earth attributed, most probably incorrectly, to Galileo since the 1640s. It is true, however, that at his trial he was forced to publicly retract the statement of a moving Earth to save his life. For more details of this famous story, see the section on page 285.

^{**} For the definition of the concept of angle, see page 64, and for the definition of the measurement units for angle see Appendix B.

^{***} Pierre Louis Moreau de Maupertuis (1698-1759), French physicist and mathematician. He was one of the key figures in the quest for the principle of least action, which he named in this way. He was also founding president of the Berlin Academy of Sciences. Maupertuis thought that the principle reflected the maximization of goodness in the universe. This idea was thoroughly ridiculed by Voltaire in this Histoire du Docteur Akakia et du natif de Saint-Malo, 1753. Maupertuis (www.voltaire-integral.com/Html/23/08DIAL.htm) performed his measurement of the Earth to distinguish between the theory of gravitation of Newton and that of Descartes, who had predicted that the Earth is elongated at the poles, instead of flattened.






FIGURE 82 The parallax – not drawn to scale.

to everyday observation.

Then, in the years 1790 to 1792 in Bologna, Giovanni Battista Guglielmini (1763–1817) finally succeeded in measuring what Galileo and Newton had predicted to be the simplest proof for the Earth's rotation. On the Earth, objects do not fall vertically, but are slightly deviated to the east. This deviation appears because an object keeps the larger horizontal velocity it had at the height from which it started falling, as shown in Figure 85. Guglielmini's result was the first non-astronomical proof of the Earth's rotation. The experiments were repeated in 1802 by Johann Friedrich Benzenberg (1777–1846). Using metal balls which he dropped from the Michaelis tower in Hamburg – a height of 76 m – Benzenberg found that the deviation to the east was 9.6 mm. Can you confirm that the value measured by Benzenberg almost agrees with the assumption that the Earth turns once every 24 hours? There is also a much smaller deviation towards the Equator, not measured by Guglielmini, Benzenberg or anybody after them up to this day; however, it completes the list of effects on free fall by the rotation of the Earth.

Challenge 222 ny

Both deviations from vertical fall are easily understood if we use the result (described below) that falling objects describe an ellipse around the centre of the rotating Earth. The elliptical shape shows that the path of a thrown stone does not lie on a plane for an observer standing on Earth; for such an observer, the exact path thus cannot be drawn on a piece of paper.

Page 168

In 1835, Gustave-Gaspard Coriolis discovered a closely related effect that nobody had as yet noticed in everyday life. Imagine a ball that rolls over a table. For a person on the floor, the ball rolls in a straight line. Now imagine that the table rotates. For the person on the floor, the ball still rolls in a straight line. But for a person on the rotating table, the ball traces a *curved* path. In short, any object that travels in a rotating background is subject to a transversal acceleration. This is the so-called *Coriolis acceleration* or *Coriolis effect*. On a rotating background, travelling objects deviate from the straight line. The best way to understand the Coriolis effect is to experience it yourself; this can be done on a carousel, as shown in Figure 86. Watching films on the internet on the topic is also helpful. Said simply, on a rotating carousel it is not easy to hit a target by throwing or rolling a ball.

Ref. 95

Also the Earth is a rotating background. On the northern hemisphere, the rotation is anticlockwise. As the result, any moving object is slightly deviated to the right (while the magnitude of its velocity stays constant). On Earth, like on all rotating backgrounds, the

FIGURE 83 The trails of the stars at night, photographed as a series of 6 min exposures (© Anthony Ayiomamitis).




FIGURE 84 Earth's deviation from spherical shape due to its rotation (exaggerated).




FIGURE 85 The deviations of free fall towards the east and towards the Equator due to the rotation of the Earth.




FIGURE 86 A typical carousel allows observing the Coriolis effect in its most striking appearance: if a person lets a ball roll with the proper speed and direction, the ball is deflected so strongly that it comes back to her.

Coriolis acceleration a results from the change of distance to the rotation axis. Can you deduce the analytical expression for the Coriolis effect, namely $a_C = -2\omega \times v$?

On Earth, the Coriolis acceleration generally has a small value. Therefore it is best observed either in large-scale or high-speed phenomena. Indeed, the Coriolis acceleration determines the handedness of many large-scale phenomena with a spiral shape, such as the directions of cyclones and anticyclones in meteorology, the general wind patterns on

Challenge 223 s




FIGURE 87 The turning motion of a pendulum showing the rotation of the Earth.

Ref. 96

Ref. 97

handedness on the northern and the southern hemisphere. Most beautifully, the Coriolis acceleration explains why icebergs do not follow the direction of the wind as they drift away from the polar caps. The Coriolis acceleration also plays a role in the flight of cannon balls (that was the original interest of Coriolis), in satellite launches, in the motion of sunspots and even in the motion of electrons in molecules. All these Coriolis accelerations are of opposite sign on the northern and southern hemispheres and thus prove the rotation of the Earth. (In the First World War, many naval guns missed their targets in the southern hemisphere because the engineers had compensated them for the Coriolis effect in the northern hemisphere.)

Earth and the deflection of ocean currents and tides. These phenomena have opposite

Ref. 98

Only in 1962, after several earlier attempts by other researchers, Asher Shapiro was the first to verify that the Coriolis effect has a tiny influence on the direction of the vortex formed by the water flowing out of a bathtub. Instead of a normal bathtub, he had to use a carefully designed experimental set-up because, contrary to an often-heard assertion, no such effect can be seen in a real bathtub. He succeeded only by carefully eliminating all disturbances from the system; for example, he waited 24 hours after the filling of the reservoir (and never actually stepped in or out of it!) in order to avoid any left-over motion of water that would disturb the effect, and built a carefully designed, completely rotationally-symmetric opening mechanism. Others have repeated the experiment in the southern hemisphere, finding opposite rotation direction and thus confirming the result. In other words, the handedness of usual bathtub vortices is *not* caused by the rotation of the Earth, but results from the way the water starts to flow out. (A number of crooks in Quito, a city located on the equator, show gullible tourists that the vortex in a sink changes when crossing the equator line drawn on the road.) But let us go on with the story about the Earth's rotation.

Ref. 98

In 1851, the French physician-turned-physicist Jean Bernard Léon Foucault (b. 1819 Paris, d. 1868 Paris) performed an experiment that removed all doubts and rendered him world-famous practically overnight. He suspended a 67 m long pendulum* in the Panthéon in Paris and showed the astonished public that the direction of its swing changed

Challenge 224 d Ref. 99

^{*} Why was such a long pendulum necessary? Understanding the reasons allows one to repeat the experiment at home, using a pendulum as short as 70 cm, with the help of a few tricks. To observe Foucault's effect with a simple set-up, attach a pendulum to your office chair and rotate the chair slowly. Several pendulum animations, with exaggerated deviation, can be found at commons.wikimedia.org/wiki/Foucault_pendulum.

over time, rotating slowly. To anybody with a few minutes of patience to watch the change of direction, the experiment proved that the Earth rotates. If the Earth did not rotate, the swing of the pendulum would always continue in the same direction. On a rotating Earth, in Paris, the direction changes to the right, in clockwise sense, as shown in Figure 87. The swing direction does not change if the pendulum is located at the Equator, and it changes to the left in the southern hemisphere.* A modern version of the pendulum can be observed via the web cam at pendelcam.kip.uni-heidelberg.de/; high speed films of the pendulum's motion during day and night can be downloaded at www.kip.uni-heidelberg.de/ OeffWiss/Pendel-Internetauftritt/zeitraffer.php.

The time over which the orientation of the pendulum's swing performs a full turn – the *precession time* – can be calculated. Study a pendulum starting to swing in the North–South direction and you will find that the precession time $T_{\rm Foucault}$ is given by

Challenge 225 d

$$T_{\text{Foucault}} = \frac{23 \text{ h } 56 \text{ min}}{\sin \varphi} \tag{32}$$

where φ is the latitude of the location of the pendulum, e.g. 0° at the Equator and 90° at the North Pole. This formula is one of the most beautiful results of Galilean kinematics.**

Foucault was also the inventor and namer of the *gyroscope*. He built the device, shown in Figure 88, in 1852, one year after his pendulum. With it, he again demonstrated the rotation of the Earth. Once a gyroscope rotates, the axis stays fixed in space – but only when seen from distant stars or galaxies. (This is not the same as talking about absolute space. Why?) For an observer on Earth, the axis direction changes regularly with a period of 24 hours. Gyroscopes are now routinely used in ships and in aeroplanes to give the direction of north, because they are more precise and more reliable than magnetic compasses. In the most modern versions, one uses laser light running in circles instead of rotating masses.***

In 1909, Roland von Eötvös measured a simple effect: due to the rotation of the Earth, the weight of an object depends on the direction in which it moves. As a result, a balance in rotation around the vertical axis does not stay perfectly horizontal: the balance starts to oscillate slightly. Can you explain the origin of the effect?

Challenge 228 s Ref. 100

Challenge 226 s

In 1910, John Hagen published the results of an even simpler experiment, proposed by Louis Poinsot in 1851. Two masses are put on a horizontal bar that can turn around a vertical axis, a so-called *isotomeograph*. Its total mass was 260 kg. If the two masses are slowly moved towards the support, as shown in Figure 89, and if the friction is kept low enough, the bar rotates. Obviously, this would not happen if the Earth were not rotating. Can you explain the observation? This little-known effect is also useful for winning bets between physicists.

Challenge 229 s

Challenge 227 s

^{*} The discovery also shows how precision and genius go together. In fact, the first person to observe the effect was Vincenzo Viviani, a student of Galileo, as early as 1661! Indeed, Foucault had read about Viviani's work in the publications of the Academia dei Lincei. But it took Foucault's genius to connect the effect to the rotation of the Earth; nobody had done so before him.

^{**} The calculation of the period of Foucault's pendulum assumes that the precession rate is constant during a rotation. This is only an approximation (though usually a good one).

^{***} Can you guess how rotation is detected in this case?










FIGURE 88 The gyroscope: the original system by Foucault with its freely movable spinning top, the mechanical device to bring it to speed, the optical device to detect its motion, the general construction principle, and a modern (triangular) ring laser gyroscope, based on colour change of rotating laser light instead of angular changes of a rotating mass (© CNAM, JAXA).




FIGURE 89 Showing the rotation of the Earth through the rotation of an axis.




FIGURE 90 Demonstrating the rotation of the Earth with water.

pdf file availabl






FIGURE 91 A modern precision ring laser interferometer (© Bundesamt für Kartographie und Geodäsie, Carl Zeiss).

Ref. 101

In 1913, Arthur Compton showed that a closed tube filled with water and some small floating particles (or bubbles) can be used to show the rotation of the Earth. The device is called a *Compton tube* or *Compton wheel*. Compton showed that when a horizontal tube filled with water is rotated by 180°, something happens that allows one to prove that the Earth rotates. The experiment, shown in Figure 90, even allows measuring the latitude of the point where the experiment is made. Can you guess what happens?

Challenge 230 d

Vol. IV, page 49

Ref. 102

Challenge 231 s

Another method to detect the rotation of the Earth using light was first realized in 1913 by the French physicist Georges Sagnac:* he used an *interferometer* to produce bright and dark fringes of light with two light beams, one circulating in clockwise direction, and the second circulating in anticlockwise direction. The interference fringes are *shifted* when the whole system rotates; the faster it rotates, the larger is the shift. A modern, high-precision version of the experiment, which uses lasers instead of lamps, is shown in Figure 91. Sagnac also determined the relation between the fringe shift and the details of the experiment. The rotation of a complete ring interferometer with angular frequency (vector) Ω produces a fringe shift of angular phase $\Delta \varphi$ given by

$$\Delta \varphi = \frac{8\pi \ \Omega \ a}{c \ \lambda} \tag{33}$$

where a is the area (vector) enclosed by the two interfering light rays, λ their wavelength and c the speed of light. The effect is now called the *Sagnac effect* after its discoverer. It had already been predicted 20 years earlier by Oliver Lodge.** Today, Sagnac interferometers are the central part of laser gyroscopes – shown in Figure 88 – and are found in every

^{*} Georges Sagnac (b. 1869 Périgeux, d. 1928 Meudon-Bellevue) was a physicist in Lille and Paris, friend of the Curies, Langevin, Perrin, and Borel. Sagnac also deduced from his experiment that the speed of light was independent from the speed of its source, and thus confirmed a prediction of special relativity.

^{**} Oliver Lodge (1851–1940) was a British physicist who studied electromagnetic waves and tried to communicate with the dead. A strange but influential figure, his ideas are often cited when fun needs to be made of physicists; for example, he was one of those (rare) physicists who believed that at the end of the nineteenth century physics was complete.




FIGURE 92 Observing the rotation of the Earth in two seconds.

passenger aeroplane, missile and submarine, in order to measure the changes of their motion and thus to determine their actual position.

A part of the fringe shift is due to the rotation of the Earth. Modern high-precision Sagnac interferometers use ring lasers with areas of a few square metres, as shown in Figure 91. Such a ring interferometer is able to measure variations of the rotation rates of the Earth of less than one part per million. Indeed, over the course of a year, the length of a day varies irregularly by a few milliseconds, mostly due to influences from the Sun or the Moon, due to weather changes and due to hot magma flows deep inside the Earth.* But also earthquakes, the El Niño effect in the climate and the filling of large water dams have effects on the rotation of the Earth. All these effects can be studied with such high-precision interferometers; they can also be used for research into the motion of the soil due to lunar tides or earthquakes, and for checks on the theory of special relativity.

Finally, in 1948, Hans Bucka developed the simplest experiment so far to show the rotation of the Earth. A metal rod allows one to detect the rotation of the Earth after only a few seconds of observation, using the set-up of Figure 92. The experiment can be easily be performed in class. Can you guess how it works?

In summary, observations show that the Earth surface rotates at 463 m/s at the Equator, a larger value than that of the speed of sound in air, which is about 340 m/s at usual conditions. The rotation also implies an acceleration, at the Equator, of 0.034 m/s^2 . We are in fact *whirling* through the universe.

How does the Earth rotate?

Is the rotation of the Earth, the length of the day, *constant* over geological time scales? That is a hard question. If you find a method leading to an answer, publish it! (The same is true for the question whether the length of the year is constant.) Only a few methods are known, as we will find out shortly.

Ref. 104

Ref. 105

Challenge 232 s

Ref. 106

^{*} The growth of leaves on trees and the consequent change in the Earth's moment of inertia, already thought of in 1916 by Harold Jeffreys, is way too small to be seen, as it is hidden by larger effects.

The rotation of the Earth is not even constant during a human lifespan. It varies by a few parts in 10⁸. In particular, on a 'secular' time scale, the length of the day increases by about 1 to 2 ms per century, mainly because of the friction by the Moon and the melting of the polar ice caps. This was deduced by studying historical astronomical observations of the ancient Babylonian and Arab astronomers. Additional 'decadic' changes have an amplitude of 4 or 5 ms and are due to the motion of the liquid part of the Earth's core.

The seasonal and biannual changes of the length of the day – with an amplitude of 0.4 ms over six months, another 0.5 ms over the year, and 0.08 ms over 24 to 26 months – are mainly due to the effects of the *atmosphere*. In the 1950s the availability of precision measurements showed that there is even a 14 and 28 day period with an amplitude of 0.2 ms, due to the Moon. In the 1970s, when *wind oscillations* with a length scale of about 50 days were discovered, they were also found to alter the length of the day, with an amplitude of about 0.25 ms. However, these last variations are quite irregular.

Also the oceans influence the rotation of the Earth, due to the tides, the ocean currents, wind forcing, and atmospheric pressure forcing. Further effects are due to the ice sheet variations and due to water evaporation and rain falls. Last but not least, flows in the interior of the Earth, both in the mantle and in the core, change the rotation. For example, earthquakes, plate motion, post-glacial rebound and volcanic eruptions all influence the rotation.

But why does the Earth rotate at all? The rotation derives from the rotating gas cloud at the origin of the solar system. This connection explains that the Sun and all planets, except one, turn around themselves in the same direction, and that they also all turn around the Sun in that same direction. But the complete story is outside the scope of this text.

The rotation around its axis is not the only motion of the Earth; it performs other motions as well. This was already known long ago. In 128 BCE, the Greek astronomer Hipparchos discovered what is today called the *(equinoctial) precession*. He compared a measurement he made himself with another made 169 years before. Hipparchos found that the Earth's axis points to different stars at different times. He concluded that the sky was moving. Today we prefer to say that the axis of the Earth is moving. During a period of 25 800 years the axis draws a cone with an opening angle of 23.5°. This motion, shown in Figure 93, is generated by the tidal forces of the Moon and the Sun on the equatorial bulge of the Earth that results form its flattening. The Sun and the Moon try to align the axis of the Earth at right angles to the Earth's path; this torque leads to the precession of the Earth's axis.

Precession is a motion common to all rotating systems: it appears in planets, spinning tops and atoms. (Precession is also at the basis of the surprise related to the suspended wheel shown on page 209.) Precession is most easily seen in spinning tops, be they suspended or not. An example is shown in Figure 94; for atomic nuclei or planets, just imagine that the suspending wire is missing and the rotating body less flat.

In addition, the axis of the Earth is not even fixed relative to the Earth's surface. In 1884, by measuring the exact angle above the horizon of the celestial North Pole, Friedrich Küstner (1856–1936) found that the axis of the Earth *moves* with respect to the Earth's crust, as Bessel had suggested 40 years earlier. As a consequence of Küstner's discovery, the International Latitude Service was created. The *polar motion* Küstner discovered turned out to consist of three components: a small linear drift – not yet understood –

Ref. 108

Ref. 109




FIGURE 93 The precession and the nutation of the Earth's axis.

Ref. 110

Ref. 111

a yearly elliptical motion due to seasonal changes of the air and water masses, and a circular motion* with a period of about 1.2 years due to fluctuations in the pressure at the bottom of the oceans. In practice, the North Pole moves with an amplitude of about 15 m around an average central position, as shown in Figure 95. Short term variations of the North Pole position, due to local variations in atmospheric pressure, to weather change and to the tides, have also been measured. Only with help of the exact position of the Earth's axis is the high precision of the GPS system possible; and only with this knowledge can artificial satellites be guided to Mars or other planets.

The details of the motion of the Earth have been studied in great detail. Table 25 gives an overview of the knowledge and the precision that is available today.

In 1912, the German meteorologist and geophysicist Alfred Wegener (1880–1930) discovered an even larger effect. After studying the shapes of the continental shelves and the geological layers on both sides of the Atlantic, he conjectured that the continents *move*, and that they are all fragments of a single continent that broke up 200 million years ago.**


^{*} The circular motion, a wobble, was predicted by the great Swiss mathematician Leonhard Euler (1707–1783). In an incredible story, using Euler's and Bessel's predictions and Küstner's data, in 1891 Seth Chandler claimed to be the discoverer of the circular component.

^{**} In this old continent, called Gondwanaland, there was a huge river that flowed westwards from the Chad to Guayaquil in Ecuador. After the continent split up, this river still flowed to the west. When the Andes



FIGURE 94

Precession of a suspended spinning top (mpg film © Lucas Barbosa)






FIGURE 95 The motion of the North Pole from 2003 to 2007, including the prediction until 2008 (left) and the average position since 1900 (right) – with 0.1 arcsecond being around 3.1 m on the surface of the Earth – not showing the diurnal and semidiurnal variations of a fraction of a millisecond of arc due to the tides (from hpiers.obspm.fr/eop-pc).

Even though at first derided across the world, Wegener's discoveries were correct. Modern satellite measurements, shown in Figure 96, confirm this model. For example, the American continent moves away from the European continent by about 10 mm every year. There are also speculations that this velocity may have been much higher at certain

appeared, the water was blocked, and many millions of years later, it flowed back. Today, the river still flows eastwards and is called the Amazon River.



FIGURE 96 The continental plates are the objects of tectonic motion.




FIGURE 97 The angular size of the Sun changes due to the elliptical motion of the Earth (© Anthony Ayiomamitis).

Page 114 Vol. III, page 174 Ref. 112 periods in the past. The way to check this is to look at the magnetization of sedimental rocks. At present, this is still a hot topic of research. Following the modern version of the model, called *plate tectonics*, the continents (with a density of $2.7 \cdot 10^3 \, \text{kg/m}^3$) float on the fluid mantle of the Earth (with a density of $3.1 \cdot 10^3 \, \text{kg/m}^3$) like pieces of cork on water, and the convection inside the mantle provides the driving mechanism for the motion.

Does the Earth Move?

The centre of the Earth is not at rest in the universe. In the third century BCE Aristarchus of Samos maintained that the Earth turns around the Sun. Experiments such as that of Figure 97 confirm that the orbit is an ellipse. However, a fundamental difficulty of the heliocentric system is that the stars look the same all year long. How can this be, if the

TABLE 25 Modern measurement data about the motion of the Earth (from hpiers.obspm.fr/eop-pc).

O B S E R V A B L E	S у м в о L	Value
Mean angular velocity of Earth	Ω	72.921 150(1) µrad/s
Nominal angular velocity of Earth (epoch 1820)	$\Omega_{ m N}$	72.921 151 467 064 µrad/s
Conventional mean solar day (epoch 1820)	d	86 400 s
Conventional sidereal day	d_{si}	86 164.090 530 832 88 s
Ratio conv. mean solar day to conv. sidereal day	$vk = d/d_{si}$	1.002 737 909 350 795
Conventional duration of the stellar day	d_{st}	86 164.098 903 691 s
Ratio conv. mean solar day to conv. stellar day	$k' = d/d_{st}$	1.002 737 811 911 354 48
General precession in longitude	p	5.028 792(2) "/a
Obliquity of the ecliptic (epoch 2000)	ε_0	23° 26′ 21.4119″
Küstner-Chandler period in terrestrial frame	T_{KC}	433.1(1.7) d
Quality factor of the Küstner-Chandler peak	Q_{KC}	170
Free core nutation period in celestial frame	$T_{ m F}$	430.2(3) d
Quality factor of the free core nutation	Q_{F}	$2\cdot 10^4$
Astronomical unit	AU	149 597 870.691(6) km
Sidereal year (epoch 2000)	$a_{\rm si}$	365.256 363 004 d
		= 365 d6h9min 9.76s
Tropical year	$a_{\rm tr}$	365.242 190 402 d
		= 365 d 5 h 48 min 45.25 s
Mean Moon period	T_{M}	27.321 661 55(1) d
Earth's equatorial radius	а	6 378 136.6(1) m
First equatorial moment of inertia	A	$8.0101(2) \cdot 10^{37} \text{ kg m}^2$
Longitude of principal inertia axis <i>A</i>	λ_A	-14.9291(10)°
Second equatorial moment of inertia	B	$8.0103(2) \cdot 10^{37} \mathrm{kg} \mathrm{m}^2$
Axial moment of inertia	C	$8.0365(2) \cdot 10^{37} \mathrm{kg} \mathrm{m}^2$
Equatorial moment of inertia of mantle	$A_{\rm m}$	$7.0165 \cdot 10^{37} \mathrm{kg} \mathrm{m}^2$
Axial moment of inertia of mantle	$C_{\rm m}$	$7.0400 \cdot 10^{37} \text{ kg m}^2$
Earth's flattening	f	1/298.25642(1)
Astronomical Earth's dynamical flattening	H = (C - A)/C	2 0.003 273 794 9(1)
Geophysical Earth's dynamical flattening	e = (C - A)/A	0.003 284 547 9(1)
Earth's core dynamical flattening	$e_{ m f}$	0.002646(2)
Second degree term in Earth's gravity potential	$J_2 = -(A + B - 2C)/(2MR^2)$	$1.0826359(1) \cdot 10^{-3}$
Secular rate of J_2	$\mathrm{d}J_2/\mathrm{d}t$	$-2.6(3) \cdot 10^{-11} / a$
Love number (measures shape distortion by tides)	k_2	0.3
Secular Love number	$k_{\rm s}$	0.9383
Mean equatorial gravity	$g_{ m eq}$	$9.7803278(10)\mathrm{m/s^2}$
Geocentric constant of gravitation	GM 3.986 004 418(8) \cdot 10 ¹⁴ m ³ /s ²	
Heliocentric constant of gravitation	GM_{\odot} 1.327 124 420 76(50) · 10 ²⁰ m ³ /s	
Moon-to-Earth mass ratio	μ	0.012 300 038 3(5)



FIGURE 98 Friedrich Wilhelm Bessel (1784–1846).

Earth travels around the Sun? The distance between the Earth and the Sun has been known since the seventeenth century, but it was only in 1837 that Friedrich Wilhelm Bessel* became the first to observe the *parallax* of a star. This was a result of extremely careful measurements and complex calculations: he discovered the *Bessel functions* in order to realize it. He was able to find a star, 61 Cygni, whose apparent position changed with the month of the year. Seen over the whole year, the star describes a small ellipse in the sky, with an opening of 0.588 " (this is the modern value). After carefully eliminating all other possible explanations, he deduced that the change of position was due to the motion of the Earth around the Sun, and from the size of the ellipse he determined the distance to the star to be 105 Pm, or 11.1 light years.

Challenge 233 s

Bessel had thus managed for the first time to measure the distance of a star. By doing so he also proved that the Earth is not fixed with respect to the stars in the sky and that the Earth indeed revolves around the Sun. The motion itself was not a surprise. It confirmed the result of the mentioned aberration of light, discovered in 1728 by James Bradley** and to be discussed below; the Earth moves around the Sun.

Vol. II, page 16

With the improvement of telescopes, other motions of the Earth were discovered. In 1748, James Bradley announced that there is a small regular *change* of the precession, which he called *nutation*, with a period of 18.6 years and an angular amplitude of 19.2 ". Nutation occurs because the plane of the Moon's orbit around the Earth is not exactly the same as the plane of the Earth's orbit around the Sun. Are you able to confirm that this situation produces nutation?

Challenge 234 ny

Astronomers also discovered that the 23.5° tilt – or *obliquity* – of the Earth's axis, the angle between its intrinsic and its orbital angular momentum, actually changes from 22.1° to 24.5° with a period of 41 000 years. This motion is due to the attraction of the Sun and the deviations of the Earth from a spherical shape. In 1941, during the Second World War, the Serbian astronomer Milutin Milankovitch (1879–1958) retreated into solitude and explored the consequences. In his studies he realized that this 41 000 year period of

^{*} Friedrich Wilhelm Bessel (1784–1846), Westphalian astronomer who left a successful business career to dedicate his life to the stars, and became the foremost astronomer of his time.

^{**} James Bradley (1693–1762), English astronomer. He was one of the first astronomers to understand the value of precise measurement, and thoroughly modernized Greenwich. He discovered the aberration of light, a discovery that showed that the Earth moves and also allowed him to measure the speed of light; he also discovered the nutation of the Earth.

Ref. 113




FIGURE 99 Changes in the Earth's motion around the Sun.

the obliquity, together with an average period of 22 000 years due to precession,* gives rise to the more than 20 *ice ages* in the last 2 million years. This happens through stronger or weaker irradiation of the poles by the Sun. The changing amounts of melted ice then lead to changes in average temperature. The last ice age had its peak about 20 000 years ago and ended around 11 800 years ago; the next is still far away. A spectacular confirmation of the relation between ice age cycles and astronomy came through measurements of oxygen isotope ratios in ice cores and sea sediments, which allow the average temperature over the past million years to be tracked. Figure 100 shows how closely the temperature follows the changes in irradiation due to changes in obliquity and precession.

The Earth's orbit also changes its *eccentricity* with time, from completely circular to slightly oval and back. However, this happens in very complex ways, not with periodic regularity, and is due to the influence of the large planets of the solar system on the Earth's orbit. The typical time scale is 100 000 to 125 000 years.

In addition, the Earth's orbit changes in *inclination* with respect to the orbits of the other planets; this seems to happen regularly every $100\,000$ years. In this period the inclination changes from $+2.5^{\circ}$ to -2.5° and back.

Even the direction in which the ellipse points changes with time. This so-called per-

^{*} In fact, the 25 800 year precession leads to three insolation periods, of 23 700, 22 400 and 19 000 years, due to the interaction between precession and perihelion shift.




FIGURE 100 Modern measurements showing how Earth's precession parameter (black curve A) and obliquity (black curve D) influence the average temperature (coloured curve B) and the irradiation of the Earth (blue curve C) over the past 800 000 years: the obliquity deduced by Fourier analysis from the irradiation data RF (blue curve D) and the obliquity deduced by Fourier analysis from the temperature (red curve D) match the obliquity known from astronomical data (black curve D); sharp cooling events took place whenever the obliquity rose while the precession parameter was falling (marked red below the temperature curve) (© Jean Jouzel/Science from Ref. 113).

ihelion shift is due in large part to the influence of the other planets; a small remaining part will be important in the chapter on general relativity. It was the first piece of data confirming the theory.

Obviously, the length of the year also changes with time. The measured variations are of the order of a few parts in 10^{11} or about 1 ms per year. However, knowledge of these changes and of their origins is much less detailed than for the changes in the Earth's rotation.

The next step is to ask whether the Sun itself moves. Indeed it does. Locally, it moves




FIGURE 101 The motion of the Sun around the galaxy.

with a speed of 19.4 km/s towards the constellation of Hercules. This was shown by William Herschel in 1783. But globally, the motion is even more interesting. The diameter of the galaxy is at least 100 000 light years, and we are located 26 000 light years from the centre. (This has been known since 1918; the centre of the galaxy is located in the direction of Sagittarius.) At our position, the galaxy is 1 300 light years thick; presently, we are 68 light years 'above' the centre plane. The Sun, and with it the solar system, takes about 225 million years to turn once around the galactic centre, its orbital velocity being around 220 km/s. It seems that the Sun will continue moving away from the galaxy plane until it is about 250 light years above the plane, and then move back, as shown in Figure 101. The oscillation period is estimated to be around 62 million years, and has been suggested as the mechanism for the mass extinctions of animal life on Earth, possibly because some gas cloud or some cosmic radiation source may be periodically encountered on the way. The issue is still a hot topic of research.

We turn around the galaxy centre because the formation of galaxies, like that of solar systems, always happens in a whirl. By the way, can you confirm from your own observation that our galaxy itself rotates?

Finally, we can ask whether the galaxy itself moves. Its motion can indeed be observed because it is possible to give a value for the motion of the Sun through the universe, defining it as the motion against the background radiation. This value has been measured to be 370 km/s. (The velocity of the *Earth* through the background radiation of course depends on the season.) This value is a combination of the motion of the Sun around the galaxy centre and of the motion of the galaxy itself. This latter motion is due to the gravitational attraction of the other, nearby galaxies in our local group of galaxies.*

In summary, the Earth really moves, and it does so in rather complex ways. As Henri Poincaré would say, if we are in a given spot today, say the Panthéon in Paris, and come

Ref. 114

Challenge 235 s

Ref. 115

^{*} This is roughly the end of the ladder. Note that the expansion of the universe, to be studied later, produces no motion.

back to the same spot tomorrow at the same time, we are in fact 31 million kilometres away. This state of affairs would make time travel extremely difficult even if it were possible (which it is not); whenever you went back to the past, you would have to get to the old spot exactly!

IS VELOCITY ABSOLUTE? - THE THEORY OF EVERYDAY RELATIVITY

Why don't we feel all the motions of the Earth? The two parts of the answer were already given in 1632. First of all, as Galileo explained, we do not feel the accelerations of the Earth because the effects they produce are too small to be detected by our senses. Indeed, many of the mentioned accelerations do induce measurable effects only in high-precision experiments, e.g. in atomic clocks.

Vol. II, page 142

But the second point made by Galileo is equally important: it is impossible to feel the high speed at which we are moving. We do not feel translational, unaccelerated motions because this is impossible *in principle*. Galileo discussed the issue by comparing the observations of two observers: one on the ground and another on the most modern means of unaccelerated transportation of the time, a ship. Galileo asked whether a man on the ground and a man in a ship moving at constant speed experience (or 'feel') anything different. Einstein used observers in trains. Later it became fashionable to use travellers in rockets. (What will come next?) Galileo explained that only *relative* velocities between bodies produce effects, not the absolute values of the velocities. For the senses, there is no difference between constant, undisturbed motion, however rapid it may be, and rest. This is now called *Galileo's principle of relativity*. In everyday life we feel motion only if the means of transportation trembles (thus if it accelerates), or if we move against the air. Therefore Galileo concludes that two observers in straight and undisturbed motion against each other cannot say who is 'really' moving. Whatever their relative speed, neither of them 'feels' in motion.*

Challenge 236 e

^{*} In 1632, in his Dialogo, Galileo writes: 'Shut yourself up with some friend in the main cabin below decks on some large ship, and have with you there some flies, butterflies, and other small flying animals. Have a large bowl of water with some fish in it; hang up a bottle that empties drop by drop into a wide vessel beneath it. With the ship standing still, observe carefully how the little animals fly with equal speed to all sides of the cabin. The fish swim indifferently in all directions; the drops fall into the vessel beneath; and, in throwing something to your friend, you need throw it no more strongly in one direction than another, the distances being equal: jumping with your feet together, you pass equal spaces in every direction. When you have observed all these things carefully (though there is no doubt that when the ship is standing still everything must happen in this way), have the ship proceed with any speed you like, so long as the motion is uniform and not fluctuating this way and that, you will discover not the least change in all the effects named, nor could you tell from any of them whether the ship was moving or standing still. In jumping, you will pass on the floor the same spaces as before, nor will you make larger jumps toward the stern than toward the prow even though the ship is moving quite rapidly, despite the fact that during the time you are in the air the floor under you will be going in a direction opposite to your jump. In throwing something to your companion, you will need no more force to get it to him whether he is in the direction of the bow or the stern, with yourself situated opposite. The droplets will fall as before into the vessel beneath without dropping toward the stern, although while the drops are in the air the ship runs many spans. The fish in their water will swim toward the front of their bowl with no more effort than toward the back, and will go with equal ease to bait placed anywhere around the edges of the bowl. Finally the butterflies and flies will continue their flights indifferently toward every side, nor will it ever happen that they are concentrated toward the stern, as if tired out from keeping up with the course of the ship, from which they will have been separated during long intervals by keeping themselves in the air. And if smoke is made by burning some

Rest is relative. Or more clearly: rest is an observer-dependent concept. This result of Galilean physics is so important that Poincaré introduced the expression 'theory of relativity' and Einstein repeated the principle explicitly when he published his famous theory of special relativity. However, these names are awkward. Galilean physics is also a theory of relativity! The relativity of rest is common to *all* of physics; it is an essential aspect of motion.

Undisturbed or uniform motion has no observable effect; only *change* of motion does. Velocity cannot be felt; acceleration can. As a result, every physicist can deduce something simple about the following statement by Wittgenstein:

Daß die Sonne morgen aufgehen wird, ist eine Hypothese; und das heißt: wir wissen nicht, ob sie aufgehen wird.*

The statement is *wrong*. Can you explain why Wittgenstein erred here, despite his strong desire not to?

Challenge 237 s

IS ROTATION RELATIVE?

When we turn rapidly, our arms lift. Why does this happen? How can our body detect whether we are rotating or not? There are two possible answers. The first approach, promoted by Newton, is to say that there is an absolute space; whenever we rotate against this space, the system reacts. The other answer is to note that whenever the arms lift, the stars also rotate, and in exactly the same manner. In other words, our body detects rotation because we move against the average mass distribution in space.

The most cited discussion of this question is due to Newton. Instead of arms, he explored the water in a rotating bucket. As usual for philosophical issues, Newton's answer was guided by the mysticism triggered by his father's early death. Newton saw absolute space as a religious concept and was not even able to conceive an alternative. Newton thus sees rotation as an absolute concept. Most modern scientist have fewer problems and more common sense than Newton; as a result, today's consensus is that rotation effects are due to the mass distribution in the universe: rotation is relative. Even the theory of general relativity confirms this conclusion.

CURIOSITIES AND FUN CHALLENGES ABOUT RELATIVITY

When travelling in the train, you can test Galileo's statement about everyday relativity of motion. Close your eyes and ask somebody to turn you around many times: are you able to say in which direction the train is running?

Challenge 238 e

* *

incense, it will be seen going up in the form of a little cloud, remaining still and moving no more toward one side than the other. The cause of all these correspondences of effects is the fact that the ship's motion is common to all the things contained in it, and to the air also. That is why I said you should be below decks; for if this took place above in the open air, which would not follow the course of the ship, more or less noticeable differences would be seen in some of the effects noted.'

^{* &#}x27;That the Sun will rise to-morrow, is an hypothesis; and that means that we do not *know* whether it will rise.' This well-known statement is found in Ludwig Wittgenstein, *Tractatus*, 6.36311.

A good bathroom scales, used to determine the weight of objects, does not show a con-Challenge 239 s stant weight when you step on it and stay motionless. Why not?

* *

Challenge 240 s If a gun located at the Equator shoots a bullet vertically, where does the bullet fall?

* *

Challenge 241 s Why are most rocket launch sites as near as possible to the Equator?

* *

At the equator, the speed of rotation of the Earth is 465 m/s, or about Mach 1.4; the latter number means that it is 1.4 times the speed of sound. This supersonic motion has two intriguing consequences.

First of all, the speed determines the size of typical weather phenomena. This size, the so-called *Rosby length*, is given by the speed of sound divided by the local rotation speed. At moderate latitudes, the Rosby length is about 2000 km. This is a sizeable fraction of the Earth's radius, so that only a few large weather systems are present on Earth at any specific time. If the Earth rotated more slowly, the weather would be determined by short-lived, local flows and have no general regularities. If the Earth rotated more rapidly, the weather would be much more violent – as on Jupiter – but the small Rosby length implies that large weather structures have a huge lifetime, such as the red spot on Jupiter, which lasted for several centuries. In a sense, the rotation of the Earth has the speed that provides the most interesting weather.

The other consequence of the value of the Earth's rotation speed concerns the atmosphere. Mach 1 is also, roughly speaking, the thermal speed of air molecules. This speed is sufficient for an air molecule to reach the characteristic height of the atmosphere, about 6 km. On the other hand, the speed of rotation Ω of the Earth determines its departure h from sphericity: the Earth is flattened, as we saw above. Roughly speaking, we have $gh = \Omega^2 R^2/2$, or about 12 km. (This is correct to within 50%, the actual value is 21 km.) We thus find that the speed of rotation of the Earth implies that its flattening is comparable to the thickness of the atmosphere.

* *

The Coriolis effect influences rivers and their shores. This surprising connection was made in 1860 by Karl Ernst von Baer who found that in Russia, many rivers flowing north in lowlands had right shores that are steep and high, and left shores that are low and flat. (Can you explain the details?) He also found that rivers in the southern hemisphere show the opposite effect.

* *

Would travelling through interplanetary space be healthy? People often fantasize about long trips through the cosmos. Experiments have shown that on trips of long duration, cosmic radiation, bone weakening, muscle degeneration and psychological problems are the biggest dangers. Many medical experts question the viability of space travel lasting longer than a couple of years. Other dangers are rapid sunburn, at least near the Sun,

Page 126

Challenge 242 e

and exposure to the vacuum. So far only one man has experienced vacuum without Ref. 116 protection. He lost consciousness after 14 seconds, but survived unharmed.

* *

Challenge 243 s In which direction does a flame lean if it burns inside a jar on a rotating turntable?

* *

Galileo's principle of everyday relativity states that it is impossible to determine an absolute velocity. It is equally impossible to determine an absolute position, an absolute time and an absolute direction. Is this correct?

Challenge 244 ny

* *

Does *centrifugal acceleration* exist? Most university students go through the shock of meeting a teacher who says that it doesn't because it is a 'fictitious' quantity, in the face of what one experiences every day in a car when driving around a bend. Simply ask the teacher who denies it to define 'existence'. (The definition physicists usually use is given later on.) Then check whether the definition applies to the term and make up your own mind

Vol. III, page 259 Challenge 245 s

Whether you like the term 'centrifugal acceleration' or avoid it by using its negative, the so-called *centripetal acceleration*, you should know it is calculated. We use a simple trick. For an object in circular motion of radius r, the magnitude v of the velocity v = dx/dt is $v = 2\pi r/T$. The vector v behaves over time exactly like the position of the object: it rotates continuously. Therefore, the magnitude a of the centrifugal/centripetal acceleration a = dv/dt is given by the corresponding expression, namely $a = 2\pi v/T$. Eliminating T, we find that the centrifugal/centripetal acceleration a of a body rotating at speed v at radius r is given by

 $a = \frac{v^2}{r} = \omega^2 r \ . \tag{34}$

This is the acceleration we feel when sitting in a car that goes around a bend.

* *

Rotation holds a surprise for anybody who studies it carefully. Angular momentum is a quantity with a magnitude and a direction. However, it is not a vector, as any mirror shows. The angular momentum of a body circling in a plane parallel to a mirror behaves in a different way from a usual arrow: its mirror image is not reflected if it points towards the mirror! You can easily check this for yourself. For this reason, angular momentum is called a *pseudovector*. The fact has no important consequences in classical physics; but we have to keep it in mind for later occasions.


Challenge 246

* *

What is the best way to transport a number of full coffee or tea cups while at the same time avoiding spilling any precious liquid?

* *

A ping-pong ball is attached by a string to a stone, and the whole is put under water in a



Challenge 248 s

Challenge 250 s

Challenge 251 ny

Challenge 252 s

FIGURE 102 How does the ball move when the jar is accelerated in direction of the arrow?

jar. The set-up is shown in Figure 102. Now the jar is accelerated horizontally. In which direction does the ball move? What do you deduce for a jar at rest?

* *

The Moon recedes from the Earth by 3.8 cm a year, due to friction. Can you find the Challenge 249 s mechanism responsible for the effect?

* *

What are earthquakes? *Earthquakes* are large examples of the same process that make a door squeak. The continental plates correspond to the metal surfaces in the joints of the door

Earthquakes can be described as energy sources. The Richter scale is a direct measure of this energy. The *Richter magnitude* $M_{\rm s}$ of an earthquake, a pure number, is defined from its energy E in joule via

$$M_{\rm s} = \frac{\log(E/1\,\mathrm{J}) - 4.8}{1.5} \,. \tag{35}$$

The strange numbers in the expression have been chosen to put the earthquake values as near as possible to the older, qualitative Mercalli scale (now called EMS98) that classifies the intensity of earthquakes. However, this is not fully possible; the most sensitive instruments today detect earthquakes with magnitudes of -3. The highest value ever measured was a Richter magnitude of 10, in Chile in 1960. Magnitudes above 12 are probably impossible. (Can you show why?)

* *

What is the motion of the point on the surface of the Earth that has Sun in its zenith (i.e., vertically above it), when seen on a map of the Earth during one day, and day after day?

* *

The moment of inertia of a body depends on the shape of the body; usually, angular momentum and the angular velocity do not point in the same direction. Can you confirm this with an example?

Can it happen that a satellite dish for geostationary TV satellites focuses the sunshine

* *






FIGURE 103 What happens when the ape climbs?

Challenge 253 s onto the receiver?

Challenge 255 s

Challenge 258 s

* *

Why is it difficult to fire a rocket from an aeroplane in the direction opposite to the Challenge 254 s motion of the plane?

* *

An ape hangs on a rope. The rope hangs over a wheel and is attached to a mass of equal weight hanging down on the other side, as shown in Figure 103. The rope and the wheel are massless and frictionless. What happens when the ape climbs the rope?

* *

Challenge 256 s Can a water skier move with a higher speed than the boat pulling him?

* *

Challenge 257 s What is the moment of inertia of a homogeneous sphere?

* *

The moment of inertia is determined by the values of its three principal axes. These are all equal for a sphere and for a cube. Does it mean that it is impossible to distinguish a sphere from a cube by their inertial behaviour?

* *

You might know the 'Dynabee', a hand-held gyroscopic device that can be accelerated to high speed by proper movements of the hand. How does it work?

* *

It is possible to make a spinning top with a metal paper clip. It is even possible to make Challenge 260 s one of those tops that turn onto their head when spinning. Can you find out how?



FIGURE 104 A long exposure of the stars at night - over the Gemini telescope in Hawaii (© Gemini Observatory/AURA).

* *

Is it true that the Moon in the first quarter in the northern hemisphere looks like the Challenge 261 s Moon in the last quarter in the southern hemisphere?

* *

An impressive confirmation that the Earth is round can be seen at sunset, if one turns, against usual habits, one's back on the Sun. On the eastern sky one can see the impressive rise of the Earth's shadow. (In fact, more precise investigations show that it is not the shadow of the Earth alone, but the shadow of its ionosphere.) One can admire a vast shadow rising over the whole horizon, clearly having the shape of a segment of a huge circle.

* *

Challenge 262 s How would Figure 104 look if taken at the Equator?

* *

Since the Earth is round, there are many ways to drive from one point on the Earth to another along a circle segment. This has interesting consequences for volley balls and for girl-watching. Take a volleyball and look at its air inlet. If you want to move the inlet to a different position with a simple rotation, you can choose the rotation axis in many different ways. Can you confirm this? In other words, when we look in a given direction and then want to look in another, the eye can accomplish this change in different ways.

Challenge 263 e

Challenge 264 s

The option chosen by the human eye had already been studied by medical scientists in the eighteenth century. It is called *Listing's 'law'*.* It states that all axes that nature chooses lie in one plane. Can you imagine its position in space? Men have a real interest that this mechanism is strictly followed; if not, looking at girls on the beach could cause the muscles moving the eyes to get knotted up.

LEGS OR WHEELS? - AGAIN

The acceleration and deceleration of standard wheel-driven cars is never much greater than about $1 g = 9.8 \,\mathrm{m/s^2}$, the acceleration due to gravity on our planet. Higher accelerations are achieved by motorbikes and racing cars through the use of suspensions that divert weight to the axes and by the use of spoilers, so that the car is pushed downwards with more than its own weight. Modern spoilers are so efficient in pushing a car towards the track that racing cars could race on the roof of a tunnel without falling down.

Through the use of special tyres these downwards forces are transformed into grip; modern racing tyres allow forward, backward and sideways accelerations (necessary for speed increase, for braking and for turning corners) of about 1.1 to 1.3 times the load. Engineers once believed that a factor 1 was the theoretical limit and this limit is still sometimes found in textbooks; but advances in tyre technology, mostly by making clever use of interlocking between the tyre and the road surface as in a gear mechanism, have allowed engineers to achieve these higher values. The highest accelerations, around $4\,g$, are achieved when part of the tyre melts and glues to the surface. Special tyres designed to make this happen are used for dragsters, but high performance radio-controlled model cars also achieve such values.

Ref. 117

How do all these efforts compare to using legs? High jump athletes can achieve peak accelerations of about 2 to 4 g, cheetahs over 3 g, bushbabies up to 13 g, locusts about 18 g, and fleas have been measured to accelerate about 135 g. The maximum acceleration known for animals is that of click beetles, a small insect able to accelerate at over $2000 \,\mathrm{m/s^2} = 200 \,g$, about the same as an airgun pellet when fired. Legs are thus definitively more efficient accelerating devices than wheels – a cheetah can easily beat any car or motorbike – and evolution developed legs, instead of wheels, to improve the chances of an animal in danger getting to safety. In short, legs *outperform* wheels.

Challenge 265 s

There are other reasons for using legs instead of wheels. (Can you name some?) For example, legs, unlike wheels, allow walking on water. Most famous for this ability is the *basilisk*, ** a lizard living in Central America and shown in Figure 105. This reptile is up to 70 cm long and has a mass of about 90 g. It looks like a miniature *Tyrannosaurus rex* and is able to run over water surfaces on its hind legs. The motion has been studied in detail with high-speed cameras and by measurements using aluminium models of the animal's feet. The experiments show that the feet slapping on the water provides only 25 % of the force necessary to run above water; the other 75 % is provided by a pocket of compressed air that the basilisks create between their feet and the water once the feet

Ref. 118

^{*} If you are interested in learning in more detail how nature and the eye cope with the complexities of three dimensions, see the schorlab.berkeley.edu/vilis/whatisLL.htm and www.physpharm.fmd.uwo.ca/undergrad/llconsequencesweb/ListingsLaw/perceptual1.htm websites.

^{**} In the Middle Ages, the term 'basilisk' referred to a mythical monster supposed to appear shortly before the end of the world. Today, it is a small reptile in the Americas.



FIGURE 105 A basilisk lizard (*Basiliscus basiliscus*) running on water, with a total length of about 25 cm, showing how the propulsing leg pushes into the water (© TERRA).



FIGURE 106 A water strider, total size about 10 mm (© Charles Lewallen).




FIGURE 107 A water walking robot, total size about 20 mm (© AIP).

are inside the water. In fact, basilisks mainly walk on air. (Both effects used by basilisks are also found in fast canoeing.) It was calculated that humans are also able to walk on water, provided their feet hit the water with a speed of 100 km/h using the simultaneous physical power of 15 sprinters. Quite a feat for all those who ever did so.

There is a second method of walking and running on water; this second method even allows its users to remain immobile on top of the water surface. This is what water striders, insects of the family *Gerridae* with a overall length of up to 15 mm, are able to do (together with several species of spiders), as shown in Figure 106. Like all insects, the water strider has six legs (spiders have eight). The water strider uses the back and front legs to hover over the surface, helped by thousands of tiny hairs attached to its body. The hairs, together with the surface tension of water, prevent the strider from getting wet. If you put shampoo into the water, the water strider sinks and can no longer move. The water strider uses its large middle legs as oars to advance over the surface, reaching speeds of up to 1 m/s doing so. In short, water striders actually row over water. The same mechanism is used by the small robots that can move over water and were developed by Metin Sitti and his group, as shown in Figure 107.

Robot design is still in its infancy. No robot can walk or even run as fast as the animal system it tries to copy. For two-legged robots, the most difficult ones, the speed record is around 3.5 leg lengths per second. In fact, there is a race going on in robotics departments: each department tries to build the first robot that is faster, either in metres per

Ref. 120

Ref. 119

second or in leg lengths per second, than the original fourlegged animal or two-legged human. The difficulties of realizing this development goal show how complicated walking motion is and how well nature has optimized living systems.

Legs pose many interesting problems. Engineers know that a staircase is comfortable to walk only if for each step the depth l plus *twice* the height h is a constant: $l + 2h = 0.63 \pm 0.02$ m. This is the so-called *staircase formula*. Why does it hold?

Challenge 266 s Challenge 267 s

Most animals have an *even* number of legs. Do you know an exception? Why not? In fact, one can argue that no animal has less than four legs. Why is this the case?

On the other hand, all animals with two legs have the legs side by side, whereas systems with two wheels have them one behind the other. Why is this not the other way round?

Challenge 268 e

Legs also provide simple distance rulers: just count your steps. In 2006, it was discovered that this method is used by certain ant species, such as *Cataglyphis fortis*. They can count to at least 25 000, as shown by Matthias Wittlinger and his team. These ants use the ability to find the shortest way back to their home even in structureless desert terrain.

Ref. 122

Ref. 121

Why do $100 \,\mathrm{m}$ sprinters run faster than ordinary people? A thorough investigation shows that the speed v of a sprinter is given by

$$v = f L_{\text{stride}} = f L_{\text{c}} \frac{F_{\text{c}}}{W} , \qquad (36)$$

where f is the frequency of the legs, $L_{\rm stride}$ is the stride length, $L_{\rm c}$ is the contact length – the length that the sprinter advances during the time the foot is in contact with the floor – W the weight of the sprinter, and $F_{\rm c}$ the average force the sprinter exerts on the floor during contact. It turns out that the frequency f is almost the same for all sprinters; the only way to be faster than the competition is to increase the stride length $L_{\rm stride}$. Also the contact length $L_{\rm c}$ varies little between athletes. Increasing the stride length thus requires that the athlete hits the ground with strong strokes. This is what athletic training for sprinters has to achieve.

SUMMARY ON GALILEAN RELATIVITY

The Earth rotates. The acceleration is so small that we do not feel it. The speed is large, but we do not feel it, because the is no way to so.

Undisturbed or inertial motion cannot be felt or measured. It is thus impossible to distinguish motion from rest; the distinction depends on the observer: *motion of bodies is relative*. That is why the soil below our feet seems so stable to us, even though it moves with high speed across the universe.

Only later on will we discover that one example of motion in nature is *not* relative: the motion of light. But we continue first with the study of motion transmitted over distance, without the use of any contact at all.



MOTION DUE TO GRAVITATION

Caddi come corpo morto cade.

Dante, *Inferno*, c. V, v. 142.*

hat we discover in our environment is *height*. Waterfalls, snow, rain, he ball of your favourite game and falling apples all rely on it. It was one of the fundamental discoveries of physics that height has this property because there is an interaction between every body and the Earth. *Gravitation* produces an acceleration along the line connecting the centres of gravity of the body and the Earth. Note that in order to make this statement, it is necessary to realize that the Earth is a body in the same way as a stone or the Moon, that this body is finite and that therefore it has a centre and a mass. Today, these statements are common knowledge, but they are by no means evident from everyday personal experience.**

How does gravitation change when two bodies are far apart? The experts on distant objects are the astronomers. Over the years they have performed numerous measurements of the movements of the Moon and the planets. The most industrious of all was Tycho Brahe,*** who organized an industrial-scale search for astronomical facts sponsored by his king. His measurements were the basis for the research of his young assistant, the Swabian astronomer Johannes Kepler**** who found the first precise description of

Ref. 123

A famous crook, Robert Peary, claimed to have reached the North Pole in 1909. (In fact, Roald Amundsen reached both the South and the North Pole first.) Among others, Peary claimed to have taken a picture there, but that picture, which went round the world, turned out to be one of the proofs that he had not been there. Can you imagine how?

Challenge 270 s

Challenge 269 s

By the way, if the Earth is round, the top of two buildings is further apart than their base. Can this effect be measured?

Challenge 271 s

^{* &#}x27;I fell like dead bodies fall.' Dante Alighieri (1265, Firenze-1321, Ravenna), the powerful Italian poet.

^{**} In several myths about the creation or the organization of the world, such as the biblical one or the Indian one, the Earth is not an object, but an imprecisely defined entity, such as an island floating or surrounded by water with unclear boundaries and unclear method of suspension. Are you able to convince a friend that the Earth is round and not flat? Can you find another argument apart from the roundness of the Earth's shadow when it is visible on the Moon, shown in Figure 109?

^{***} Tycho Brahe (b. 1546 Scania, d. 1601 Prague), famous Danish astronomer, builder of Uraniaborg, the astronomical castle. He consumed almost 10 % of the Danish gross national product for his research, which produced the first star catalogue and the first precise position measurements of planets.

^{****} Johannes Kepler (1571 Weil der Stadt-1630 Regensburg) studied Protestant theology and became a teacher of mathematics, astronomy and rhetoric. He helped his mother to defend herself successfully in a trial where she was accused of witchcraft. His first book on astronomy made him famous, and he became



FIGURE 108 The retrograde motion of planet Mars across the sky – the Pleiades star cluster is at the top left – when the planet is on the other side of the sun. The pictures were taken about a week apart and superimposed. The motion is one of the many example that are fully explained by universal gravitation (© Tunc Tezel).

Vol. III, page 259 planetary motion. This is not an easy task, as the observation of Figure 108 shows.

In 1684, all observations about planets and stones were condensed into an astonishingly simple result by the English physicist Robert Hooke and a few others:* every body of mass M attracts any other body towards its centre with an acceleration whose magnitude a is given by

$$a = G \frac{M}{r^2} \tag{37}$$

where *r* is the centre-to-centre distance of the two bodies. This is called *universal gravita*tion, or the *universal 'law' of gravitation*, because it is valid in general, both on Earth and in the sky. The proportionality constant *G* is called the *gravitational constant*; it is one of the fundamental constants of nature, like the speed of light or the quantum of action. More about *G* will be said shortly. The effect of gravity thus decreases with increasing distance; gravity depends on the inverse distance squared of the bodies under consideration. If bodies are small compared with the distance *r*, or if they are spherical, expression

Page 155

assistant to Tycho Brahe and then, at his teacher's death, the Imperial Mathematician. He was the first to use mathematics in the description of astronomical observations, and introduced the concept and field of 'celestial physics'.

^{*} Robert Hooke (1635–1703), important English physicist and secretary of the Royal Society. Apart from discovering the inverse square relation and many others, such as Hooke's 'law', he also wrote the *Micrographia*, a beautifully illustrated exploration of the world of the very small.



FIGURE 109 How to compare the radius of the Earth with that of the Moon during a partial lunar eclipse (© Anthony Ayiomamitis).

(37) is correct as it stands; for non-spherical shapes the acceleration has to be calculated separately for each part of the bodies and then added together.

This inverse square dependence is often called Newton's 'law' of gravitation, because the English physicist Isaac Newton proved more elegantly than Hooke that it agreed with all astronomical and terrestrial observations. Above all, however, he organized a better public relations campaign, in which he falsely claimed to be the originator of the idea.

Ref. 124

Newton published a simple proof showing that this description of astronomical motion also gives the correct description for stones thrown through the air, down here on 'father Earth'. To achieve this, he compared the acceleration a_{m} of the Moon with that of stones g. For the ratio between these two accelerations, the inverse square relation predicts a value $g/a_{\rm m} = d_{\rm m}^2/R^2$, where $d_{\rm m}$ the distance of the Moon and R is the radius of the Earth. The Moon's distance can be measured by triangulation, comparing the position of the Moon against the starry background from two different points on Earth.* The result is $d_{\rm m}/R = 60 \pm 3$, depending on the orbital position of the Moon, so that an average ratio $g/a_{\rm m}=3.6\cdot 10^3$ is predicted from universal gravity. But both accelerations can also be measured directly. At the surface of the Earth, stones are subject to an acceleration due to gravitation with magnitude $g = 9.8 \,\mathrm{m/s^2}$, as determined by measuring the time that stones need to fall a given distance. For the Moon, the definition of acceleration, a = dv/dt, in the case of circular motion – roughly correct here – gives $a_{\rm m}=d_{\rm m}(2\pi/T)^2$, where $T=2.4\,{\rm Ms}$ is the time the Moon takes for one orbit around the Earth.** The measurement of the radius of the Earth*** yields $R = 6.4 \,\mathrm{Mm}$, so that the average Earth-Moon distance is $d_{\rm m}=0.38\,{\rm Gm}$. One thus has $g/a_{\rm m}=3.6\cdot 10^3$, in

Challenge 272 s

^{*} The first precise – but not the first – measurement was achieved in 1752 by the French astronomers Lalande and La Caille, who simultaneously measured the position of the Moon seen from Berlin and from Le Cap. ** This expression for the centripetal acceleration is deduced easily by noting that for an object in circular motion, the magnitude v of the velocity v = dx/dt is given as $v = 2\pi r/T$. The drawing of the vector v over time, the so-called *hodograph*, shows that it behaves exactly like the position of the object. Therefore the magnitude a of the acceleration a = dv/dt is given by the corresponding expression, namely $a = 2\pi v/T$. *** This is the hardest quantity to measure oneself. The most surprising way to determine the Earth's size is






FIGURE 110 A physicist's and an artist's view of the fall of the Moon: a diagram by Christiaan Huygens (not to scale) and a marble statue by Auguste Rodin.

agreement with the above prediction. With this famous 'Moon calculation' we have thus shown that the inverse square property of gravitation indeed describes both the motion of the Moon and that of stones. You might want to deduce the value of the product *GM* for Earth.

Challenge 275 s

Universal gravitation describes motion due to gravity on Earth and in the sky. This was an important step towards the unification of physics. Before this discovery, from the observation that on the Earth all motion eventually comes to rest, whereas in the sky all motion is eternal, Aristotle and many others had concluded that motion in the sublunar world has different properties from motion in the translunar world. Several thinkers had criticized this distinction, notably the French philosopher and rector of the University of Paris, Jean Buridan.* The Moon calculation was the most important result showing this distinction to be wrong. This is the reason for calling the expression (37) the universal gravitation.

Universal gravitation allows us to answer another old question. Why does the Moon not fall from the sky? Well, the preceding discussion showed that fall is motion due to

Ref. 125

Ref. 127

the following: watch a sunset in the garden of a house, with a stopwatch in hand. When the last ray of the Sun disappears, start the stopwatch and run upstairs. There, the Sun is still visible; stop the stopwatch when the Sun disappears again and note the time t. Measure the height distance h of the two eye positions where the Sun was observed. The Earth's radius R is then given by $R = k h/t^2$, with $k = 378 \cdot 10^6 \text{ s}^2$.

Challenge 273 s Ref. 126

Page 66

There is also a simple way to measure the distance to the Moon, once the size of the Earth is known. Take a photograph of the Moon when it is high in the sky, and call θ its zenith angle, i.e., its angle from the vertical. Make another photograph of the Moon a few hours later, when it is just above the horizon. On this picture, unlike a common optical illusion, the Moon is smaller, because it is further away. With a sketch the reason for this becomes immediately clear. If q is the ratio of the two angular diameters, the Earth–Moon distance $d_{\rm m}$ is given by the relation $d_{\rm m}^2 = R^2 + (2Rq\cos\theta/(1-q^2))^2$. Enjoy finding its derivation from the

Challenge 274 s

Another possibility is to determine the size of the Moon by comparing it with the size of the shadow of the Earth during a lunar eclipse, as shown in Figure 109. The distance to the Moon is then computed from its angular size, about 0.5°.

* Jean Buridan (c. 1295 to c. 1366) was also one of the first modern thinkers to discuss the rotation of the Earth about an axis.



FIGURE 111 A precision second pendulum, thus about 1 m in length; almost at the upper end, the vacuum chamber that compensates for changes in atmospheric pressure; towards the lower end, the wide construction that compensates for temperature variations of pendulum length; at the very bottom, the screw that compensates for local variations of the gravitational acceleration, giving a final precision of about 1 s per month (© Erwin Sattler OHG).

gravitation. Therefore the Moon actually *is* falling, with the peculiarity that instead of falling *towards* the Earth, it is continuously falling *around* it. Figure 110 illustrates the idea. The Moon is continuously missing the Earth.*

The Moon is not the only object that falls around the Earth. Figure 112 shows another. Universal gravity also explains why the Earth and most planets are (almost) spherical. Since gravity increases with decreasing distance, a liquid body in space will always try to form a spherical shape. Seen on a large scale, the Earth is indeed liquid. We also know that the Earth is cooling down – that is how the crust and the continents formed. The sphericity of smaller solid objects encountered in space, such as the Moon, thus means that they used to be liquid in older times.

PROPERTIES OF GRAVITATION

Gravitation implies that the path of a stone is not a parabola, as stated earlier, but actually an *ellipse* around the centre of the Earth. This happens for exactly the same reason that the planets move in ellipses around the Sun. Are you able to confirm this statement?

Universal gravitation allows us to understand the puzzling acceleration value

Ref. 128 Challenge 276 d

Page 74

Page 166

^{*} Another way to put it is to use the answer of the Dutch physicist Christiaan Huygens (1629–1695): the Moon does not fall from the sky because of the centrifugal acceleration. As explained on page 144, this explanation is often out of favour at universities.

There is a beautiful problem connected to the left side of the figure: Which points on the surface of the Earth can be hit by shooting from a mountain? And which points can be hit by shooting horizontally?




FIGURE 112 The man in orbit feels no weight, the blue atmosphere, which is not, does (NASA).

TABLE 26 Some measured values of the acceleration due to gravity.

PLACE	Value
Poles	9.83 m/s ²
Trondheim	$9.8215243 \mathrm{m/s^2}$
Hamburg	$9.8139443 \mathrm{m/s^2}$
Munich	$9.8072914 \mathrm{m/s^2}$
Rome	$9.8034755 \mathrm{m/s^2}$
Equator	$9.78 \mathrm{m/s^2}$
Moon	$1.6 \mathrm{m/s^2}$
Sun	$273 \mathrm{m/s^2}$

 $g = 9.8 \,\mathrm{m/s^2}$ we encountered in equation (6). The value is thus due to the relation

$$g = GM_{\text{Earth}}/R_{\text{Earth}}^2 . (38)$$

The expression can be deduced from equation (37), universal gravity, by taking the Earth to be spherical. The everyday acceleration of gravity g thus results from the size of the Earth, its mass, and the universal constant of gravitation G. Obviously, the value for g is almost constant on the surface of the Earth, as shown in Table 26, because the Earth is almost a sphere. Expression (38) also explains why g gets smaller as one rises above the Earth, and the deviations of the shape of the Earth from sphericity explain why g is different at the poles and higher on a plateau. (What would g be on the Moon? On Mars? On Jupiter?)

Challenge 277 s

By the way, it is possible to devise a simple machine, other than a yo-yo, that slows

down the effective acceleration of gravity by a known amount, so that one can measure

Challenge 278 s its value more easily. Can you imagine it?

Note that 9.8 is roughly π^2 . This is *not* a coincidence: the metre has been chosen in such a way to make this (roughly) correct. The period T of a swinging pendulum, i.e., a back and forward swing, is given by*

Challenge 279 s

$$T = 2\pi \sqrt{\frac{l}{g}} , \qquad (39)$$

where l is the length of the pendulum, and $g = 9.8 \,\mathrm{m/s^2}$ is the gravitational acceleration. (The pendulum is assumed to consist of a compact mass attached to a string of negligible mass.) The oscillation time of a pendulum depends only on the length of the string and on g, thus on the planet it is located on.

Challenge 281 e

Ref. 129

If the metre had been defined such that T/2 = 1 s, the value of the normal acceleration g would have been exactly π^2 m/s². Indeed, this was the first proposal for the definition of the metre; it was made in 1673 by Huygens and repeated in 1790 by Talleyrand, but was rejected by the conference that defined the metre because variations in the value of g with geographical position, temperature-induced variations of the length of a pendulum and even air pressure variations induce errors that are too large to yield a definition of useful precision. (Indeed, all these effects must be corrected in pendulum clocks, as shown in Figure 111.)

Finally, the proposal was made to define the metre as 1/40 000 000 of the circumference of the Earth through the poles, a so-called *meridian*. This proposal was almost identical to – but much more precise than – the pendulum proposal. The meridian definition of the metre was then adopted by the French national assembly on 26 March 1791, with the statement that 'a meridian passes under the feet of every human being, and all meridians are equal'. (Nevertheless, the distance from Equator to the poles is not exactly 10 Mm; that is a strange story. One of the two geographers who determined the size of the first metre stick was dishonest. The data he gave for his measurements – the general method of which is shown in Figure 113 – was fabricated. Thus the first official metre stick in Paris was shorter than it should be.)

Continuing our exploration of the gravitational acceleration g, we can still ask: Why does the Earth have the mass and size it has? And why does G have the value it has? The first question asks for a history of the solar system; it is still unanswered and is topic of research. The second question is addressed in Appendix B.

If gravitation is indeed universal, and if all objects really attract each other, attraction should also occur for objects in everyday life. Gravity must also work *sideways*. This is

^{*} Formula (39) is noteworthy mainly for all that is missing. The period of a pendulum does *not* depend on the mass of the swinging body. In addition, the period of a pendulum does *not* depend on the amplitude. (This is true as long as the oscillation angle is smaller than about 15°.) Galileo discovered this as a student, when observing a chandelier hanging on a long rope in the dome of Pisa. Using his heartbeat as a clock he found that even though the amplitude of the swing got smaller and smaller, the time for the swing stayed the same.

A leg also moves like a pendulum, when one walks normally. Why then do taller people tend to walk faster? Is the relation also true for animals of different size?



FIGURE 113 The measurements that lead to the definition of the metre (© Ken Alder).

indeed the case, even though the effects are so small that they were measured only long after universal gravity had predicted them. In fact, measuring this effect allows the gravitational constant *G* to be determined. Let us see how.

We note that measuring the gravitational constant *G* is also the only way to determine the mass of the *Earth*. The first to do so, in 1798, was the English physicist Henry Cavendish; he used the machine, ideas and method of John Michell who died when attempting the experiment. Michell and Cavendish* called the aim and result of their experiments 'weighing the Earth'.

The idea of Michell was to suspended a horizontal handle, with two masses at the end, at the end of a long metal wire. He then approached two large masses at the two ends of the handle, avoiding any air currents, and measured how much the handle rotated. Figure 114 shows how to repeat this experiment in your basement, and Figure 115 how to perform it when you have a larger budget.

The value the gravitational constant G found in more elaborate versions of the Michell–Cavendish experiments is

$$G = 6.7 \cdot 10^{-11} \,\text{Nm}^2/\text{kg}^2 = 6.7 \cdot 10^{-11} \,\text{m}^3/\text{kg s}^2 \,. \tag{40}$$

Cavendish's experiment was thus the first to confirm that gravity also works sideways. The experiment also allows deducing the mass *M* of the Earth from its radius *R* and the

^{*} Henry Cavendish (b. 1731 Nice, d. 1810 London) was one of the great geniuses of physics; rich, autistic, misogynist, unmarried and solitary, he found many rules of nature, but never published them. Had he done so, his name would be much more well known. John Michell (1724–1793) was church minister, geologist and amateur astronomer.













FIGURE 114 An experiment that allows weighing the Earth and proving that gravity also works sideways and curves space. Top left and right: a torsion balance made of foam and lead, with pétanque (boules) masses as fixed masses; centre right: a torsion balance made of wood and lead, with stones as fixed masses; bottom: a time sequence showing how the stones do attract the lead (© John Walker).

Challenge 282 e Vol. II, page 124

Ref. 130

relation $g = GM/R^2$. Finally, as we will see later on, this experiment proves, if we keep in mind that the speed of light is finite and invariant, that space is curved. All this is achieved with this simple set-up!

Cavendish found a mass density of the Earth of 5.5 times that of water. At his time, this was a surprising result, because rock only has 2.8 times the density of water.

Gravitation is weak. For example, two average people 1 m apart feel an acceleration towards each other that is less than that exerted by a common fly when landing on the skin. Therefore we usually do not notice the attraction to other people. When we notice it, it is much stronger than that. The measurement of G thus proves that gravitation cannot be the true cause of people falling in love, and also that erotic attraction is not of gravitational origin, but of a different source. The physical basis for love will be studied

Challenge 283 s




FIGURE 115 A modern precision torsion balance experiment to measure the gravitational constant, performed at the University of Washington (© Eöt-Wash Group).




FIGURE 116 The potential and the gradient.

Vol. III, page 14 later in our walk: it is called *electromagnetism*.

THE GRAVITATIONAL POTENTIAL

Gravity has an important property: all effects of gravitation can also be described by another observable, namely the *(gravitational) potential* φ . We then have the simple relation that the acceleration is given by the *gradient* of the potential

$$\mathbf{a} = -\nabla \varphi$$
 or $\mathbf{a} = -\operatorname{grad} \varphi$. (41)

The gradient is just a learned term for 'slope along the steepest direction'. The gradient is defined for any point on a slope, is large for a steep one and small for a shallow one. The gradient points in the direction of steepest ascent, as shown in Figure 116. The gradient is abbreviated ∇ , pronounced 'nabla', and is mathematically defined through the relation

 $\nabla \varphi = (\partial \varphi / \partial x, \partial \varphi / \partial y, \partial \varphi / \partial z) = \operatorname{grad} \varphi$. The minus sign in (41) is introduced by convention, in order to have higher potential values at larger heights.* In everyday life, when the spherical shape of the Earth can be neglected, the gravitational potential is given by

$$\varphi = gh. \tag{42}$$

The potential φ is an interesting quantity; with a single number at every position in space we can describe the vector aspects of gravitational acceleration. It automatically gives that gravity in New Zealand acts in the opposite direction to gravity in Paris. In addition, the potential suggests the introduction of the so-called *potential energy U* by setting

$$U = m\varphi \tag{43}$$

and thus allowing us to determine the change of kinetic energy T of a body falling from a point 1 to a point 2 via

$$T_1 - T_2 = U_2 - U_1$$
 or $\frac{1}{2}m_1 v_1^2 - \frac{1}{2}m_2 v_2^2 = m\varphi_2 - m\varphi_1$. (44)

In other words, the *total energy*, defined as the sum of kinetic and potential energy, is *conserved* in motion due to gravity. This is a characteristic property of gravitation. Gravity conserves energy and momentum.

Not all accelerations can be derived from a potential; systems with this property are called *conservative*. Observation shoes that accelerations due to friction are not conservative, but accelerations due to electromagnetism are. In short, we can either say that gravity can be described by a potential, or say that it conserves energy and momentum. When the spherical shape of the Earth can be neglected, the potential energy of an object at height h is given by

$$U = mqh. (45)$$

To get a feeling of how much energy this is, answer the following question. A car with mass 1 Mg falls down a cliff of 100 m. How much water can be heated from freezing point to boiling point with the energy of the car?

THE SHAPE OF THE EARTH

Page 101

Challenge 284 s

Challenge 285 e For a spherical or a point-like body of mass M, the potential φ is

$$\varphi = -G \frac{M}{r} \ . \tag{46}$$

A potential considerably simplifies the description of motion, since a potential is additive: given the potential of a point particle, we can calculate the potential and then the

– The Adventure of Physics - noff file available free of charge at www.motionmountain net - Convribt © Christoph Schiller November 1997–March 201

^{*} In two or more dimensions slopes are written $\partial \varphi/\partial z$ – where ∂ is still pronounced 'd' – because in those cases the expression $d\varphi/dz$ has a slightly different meaning. The details lie outside the scope of this walk.




FIGURE 117 The shape of the Earth, with exaggerated height scale (© GeoForschungsZentrum Potsdam).

Challenge 287 s Ref. 131

Ref. 132

Ref. 13

Challenge 288 ny

motion around any other irregularly shaped object.* Interestingly, the number d of dimensions of space is coded into the potential φ of a spherical mass: the dependence of φ on the radius r is in fact $1/r^{d-2}$. The exponent d-2 has been checked experimentally to extremely high precision; no deviation of d from 3 has ever been found.

The concept of potential helps in understanding the *shape* of the Earth. Since most of the Earth is still liquid when seen on a large scale, its surface is always horizontal with respect to the direction determined by the combination of the accelerations of gravity and rotation. In short, the Earth is *not* a sphere. It is not an ellipsoid either. The mathematical shape defined by the equilibrium requirement is called a *geoid*. The geoid shape, illustrated in Figure 117, differs from a suitably chosen ellipsoid by at most 50 m. Can you describe the geoid mathematically? The geoid is an excellent approximation to the actual shape of the Earth; sea level differs from it by less than 20 metres. The differences can be measured with satellite radar and are of great interest to geologists and geographers. For example, it turns out that the South Pole is nearer to the equatorial plane than the

$$\Delta \varphi = 4\pi G \rho \tag{47}$$

where $\rho = \rho(x,t)$ is the mass volume density of the body and the operator Δ , pronounced 'delta', is defined as $\Delta f = \nabla \nabla f = \partial^2 f/\partial x^2 + \partial^2 f/\partial y^2 + \partial^2 f/\partial z^2$. Equation (47) is called the *Poisson equation* for the potential φ . It is named after Siméon-Denis Poisson (1781–1840), eminent French mathematician and physicist. The positions at which ρ is not zero are called the *sources* of the potential. The so-called source term $\Delta \varphi$ of a function is a measure for how much the function $\varphi(x)$ at a point x differs from the average value in a region around that point. (Can you show this, by showing that $\Delta \varphi \sim \bar{\varphi} - \varphi(x)$?) In other words, the Poisson equation (47) implies that the actual value of the potential at a point is the same as the average value around that point minus the mass density multiplied by $4\pi G$. In particular, in the case of empty space the potential at a point is equal to the average of the potential around that point.

Often the concept of *gravitational field* is introduced, defined as $g = -\nabla \varphi$. We avoid this in our walk, because we will discover that, following the theory of relativity, gravity is not due to a field at all; in fact even the concept of gravitational potential turns out to be only an approximation.

Challenge 286 ny

^{*} Alternatively, for a general, extended body, the potential is found by requiring that the *divergence* of its gradient is given by the mass (or charge) density times some proportionality constant. More precisely, one has

North Pole by about 30 m. This is probably due to the large land masses in the northern hemisphere.

Page 126

Appendix B

Above we saw how the inertia of matter, through the so-called 'centrifugal force', increases the radius of the Earth at the Equator. In other words, the Earth is *flattened* at the poles. The Equator has a radius a of 6.38 Mm, whereas the distance b from the poles to the centre of the Earth is 6.36 Mm. The precise flattening (a - b)/a has the value 1/298.3 = 0.0034. As a result, the top of Mount Chimborazo in Ecuador, even though its height is only 6267 m above sea level, is about 20 km farther away from the centre of the Earth than the top of Mount Sagarmatha* in Nepal, whose height above sea level is 8850 m. The top of Mount Chimborazo is in fact the point on the surface most distant from the centre of the Earth.

The shape of the Earth has another important consequence. If the Earth stopped rotating (but kept its shape), the water of the oceans would flow from the equator to the poles; all of Europe would be under water, except for the few mountains of the Alps that are higher than about 4 km. The northern parts of Europe would be covered by between 6 km and 10 km of water. Mount Sagarmatha would be over 11 km above sea level. We would also walk inclined. If we take into account the resulting change of shape of the Earth, the numbers come out somewhat smaller. In addition, the change in shape would produce extremely strong earthquakes and storms. As long as there are none of these effects, we can be *sure* that the Sun will indeed rise tomorrow, despite what some philosophers pretended.

Page 142

DYNAMICS - HOW DO THINGS MOVE IN VARIOUS DIMENSIONS?

The concept of potential is a powerful tool. If a body can move only along a – straight or curved – line, the concepts of kinetic and potential energy are sufficient to determine completely the way the body moves. In short, motion in *one dimension* follows directly from energy conservation.

If a body can move in *two dimensions* – i.e., on a flat or curved surface – *and* if the forces involved are *internal* (which is always the case in theory, but not in practice), the conservation of angular momentum can be used. The full motion in two dimensions thus follows from energy and angular momentum conservation. For example, all properties of free fall follow from energy and angular momentum conservation. (Are you able to show this?) Again, the potential is essential.

Challenge 289 s

In the case of motion in *three dimensions*, a more general rule for determining motion is necessary. If more than *two spatial dimensions* are involved conservation is insufficient to determine how a body moves. It turns out that general motion follows from a simple principle: the time average of the difference between kinetic and potential energy must be as small as possible. This is called the *least action principle*. We will explain the details of this calculation method later. But again, the potential is the main ingredient in the calculation of change, and thus in the description of any example of motion.

Page 213

For simple gravitational motions, motion is two-dimensional, in a plane. Most three-dimensional problems are outside the scope of this text; in fact, some of these problems are so hard that they still are subjects of research. In this adventure, we will explore three-dimensional motion only for selected cases that provide important insights.

^{*} Mount Sagarmatha is sometimes also called Mount Everest.




FIGURE 118 Some important concepts when observing the stars at night.

GRAVITATION IN THE SKY

Page 193

The expression $a = GM/r^2$ for the acceleration due to universal gravity also describes the motion of all the planets across the sky. We usually imagine to be located at the centre of the Sun and say that the planets 'orbit the Sun'. How can we check this?

First of all, looking at the sky at night, we can check that the planets always stay within the *zodiac*, a narrow stripe across the sky. The centre line of the zodiac gives the path of the Sun and is called the *ecliptic*, since the Moon must be located on it to produce an eclipse. This shows that planets move (approximately) in a single, common plane.*

The detailed motion of the planets is not easy to describe. As Figure 118 shows, observing a planet or star requires measuring various angles. For a planet, these angles change every night. From the way the angles change, one can deduce the motion of the planets. A few generations before Hooke, using the observations of *Tycho Brahe*, the Swabian astronomer Johannes Kepler, in his painstaking research on the movements of the planets in the zodiac, had deduced several 'laws'. The three main ones are as follows:

- 1. Planets move on ellipses with the Sun located at one focus (1609).
- 2. Planets sweep out equal areas in equal times (1609).
- 3. All planets have the same ratio T^2/d^3 between the orbit duration T and the semimajor axis d (1619).

Kepler's results are illustrated in Figure 119. The sheer work required to deduce the three 'laws' was enormous. Kepler had no calculating machine available. The calculation tech-

^{*} The apparent height of the ecliptic changes with the time of the year and is the reason for the changing seasons. Therefore seasons are a gravitational effect as well.




FIGURE 119 The motion of a planet around the Sun, showing its semimajor axis d, which is also the spatial average of its distance from the Sun

nology he used was the recently discovered logarithms. Anyone who has used tables of logarithms to perform calculations can get a feeling for the amount of work behind these three discoveries.

Now comes the central point. The huge volume of work by Brahe and Kepler can be summarized in the expression

$$a = GM/r^2 , (48)$$

as Hooke and a few others had stated. Let us see why.

Ref. 134

Why is the usual orbit an ellipse? The simplest argument is given in Figure 120. We know that the acceleration due to gravity varies as $a = GM/r^2$. We also know that an orbiting body of mass m has a constant energy E < 0. We then can draw, around the Sun, the circle with radius R = -GMm/E, which gives the largest distance that a body with energy E can be from the Sun. We now project the planet position P onto this circle, thus constructing a position S. We then reflect S along the tangent to get a position F. This last position F is constant in time, as a simple argument shows. (Can you find it?) As a result of the construction, the distance sum OP+PF is constant in time, and given by the radius R = -GMm/E. Since this distance sum is constant, the orbit is an ellipse, because an ellipse is precisely the curve that appears when this sum is constant. (Remember that an ellipse can be drawn with a piece of rope in this way.) Point F, like the Sun, is a focus of the ellipse. This is the first of Kepler's 'laws'.

Challenge 290 s

Can you confirm that also the other two of Kepler's 'laws' follow from Hooke's expression of universal gravity? Publishing this result was the main achievement of Newton. Try to repeat his achievement; it will show you not only the difficulties, but also the possibilities of physics, and the joy that puzzles give.

Challenge 291 s

The second of Kepler's 'laws', about equal swept areas, implies that planets move faster when they are near the Sun. It is a simple way to state the conservation of angular momentum. What does the third 'law' state?

Challenge 292 e

Newton solved these puzzles with geometric drawing – though in quite a complex manner. It is well known that Newton was not able to write down, let alone handle, differential equations at the time he published his results on gravitation. In fact, Newton's notation and calculation methods were poor. (Much poorer than yours!) The English

Ref. 26




FIGURE 120 The proof that a planet moves in an ellipse (magenta) around the Sun, given an inverse square distance relation for gravitation (see text).

mathematician Godfrey Hardy* used to say that the insistence on using Newton's integral and differential notation, rather than the earlier and better method, still common today, due to his rival Leibniz – threw back English mathematics by 100 years.

To sum up, Kepler, Hooke and Newton became famous because they brought order to the description of planetary motion. They showed that all motion due to gravity follows from the same description, the inverse square distance. For this reason, the inverse square distance relation $a = GM/r^2$ is called the *universal* law of gravity. Achieving this unification of motion description, though of small practical significance, was widely publicized. The main reason were the age-old prejudices and fantasies linked with astrology.

In fact, the inverse square distance relation explains many additional phenomena. It explains the motion and shape of the Milky Way and of the other galaxies, the motion of many weather phenomena, and explains why the Earth has an atmosphere but the Moon does not. (Can you explain this?) In fact, universal gravity explains much more about the Moon.

Challenge 293 s

^{*} Godfrey Harold Hardy (1877–1947) was an important English number theorist, and the author of the well-known *A Mathematician's Apology*. He also 'discovered' the famous Indian mathematician Srinivasa Ramanujan, and brought him to Britain.



FIGURE 121 The change of the moon during the month, showing its libration (QuickTime film @ Martin Elsässer)




FIGURE 122 High resolution maps (not photographs) of the near side (left) and far side (right) of the moon, showing how often the latter saved the Earth from meteorite impacts (courtesy USGS).

THE MOON

How long is a day on the Moon? The answer is roughly 29 Earth-days. That is the time that it takes for an observer on the Moon to see the Sun again in the same position in the sky.

One often hears that the Moon always shows the same side to the Earth. But this is wrong. As one can check with the naked eye, a given feature in the centre of the face of the Moon at full Moon is not at the centre one week later. The various motions leading to this change are called *librations*; they are shown in the film in Figure 121. The motions appear mainly because the Moon does not describe a circular, but an elliptical orbit around the Earth and because the axis of the Moon is slightly inclined, compared with that of its rotation around the Earth. As a result, only around 45 % of the Moon's surface is permanently hidden from Earth.

The first photographs of the hidden area were taken in the 1960s by a Soviet artificial satellite; modern satellites provided exact maps, as shown in Figure 122. (Just zoom into the figure for fun.) The hidden surface is much more irregular than the visible one, as

the hidden side is the one that intercepts most asteroids attracted by the Earth. Thus the gravitation of the Moon helps to deflect asteroids from the Earth. The number of animal life extinctions is thus reduced to a small, but not negligible number. In other words, the gravitational attraction of the Moon has saved the human race from extinction many times over.*

The trips to the Moon in the 1970s also showed that the Moon originated from the Earth itself: long ago, an object hit the Earth almost tangentially and threw a sizeable fraction of material up into the sky. This is the only mechanism able to explain the large size of the Moon, its low iron content, as well as its general material composition.

The Moon is receding from the Earth at 3.8 cm a year. This result confirms the old deduction that the tides slow down the Earth's rotation. Can you imagine how this measurement was performed? Since the Moon slows down the Earth, the Earth also changes shape due to this effect. (Remember that the shape of the Earth depends on its speed of rotation.) These changes in shape influence the tectonic activity of the Earth, and maybe also the drift of the continents.

The Moon has many effects on animal life. A famous example is the midge *Clunio*, which lives on coasts with pronounced tides. Clunio spends between six and twelve weeks as a larva, sure then hatches and lives for only one or two hours as an adult flying insect, during which time it reproduces. The midges will only reproduce if they hatch during the low tide phase of a spring tide. Spring tides are the especially strong tides during the full and new moons, when the solar and lunar effects combine, and occur only every 14.8 days. In 1995, Dietrich Neumann showed that the larvae have two built-in clocks, a circadian and a circalunar one, which together control the hatching to precisely those few hours when the insect can reproduce. He also showed that the circalunar clock is synchronized by the brightness of the Moon at night. In other words, the larvae monitor the Moon at night and then decide when to hatch: they are the smallest known astronomers.

If insects can have circalunar cycles, it should come as no surprise that women also have such a cycle; however, in this case the precise origin of the cycle length is still unknown and a topic of research.

The Moon also helps to stabilize the tilt of the Earth's axis, keeping it more or less fixed relative to the plane of motion around the Sun. Without the Moon, the axis would change its direction irregularly, we would not have a regular day and night rhythm, we would have extremely large climate changes, and the evolution of life would have been impossible. Without the Moon, the Earth would also rotate much faster and we would have much less clement weather. The Moon's main remaining effect on the Earth, the precession of its axis, is responsible for the ice ages.

ORBITS - AND CONIC SECTIONS

The path of a body continuously orbiting another under the influence of gravity is an *ellipse* with the central body at one focus. A circular orbit is also possible, a circle being a special case of an ellipse. Single encounters of two objects can also be *parabolas* or *hyperbolas*, as shown in Figure 124. Circles, ellipses, parabolas and hyperbolas are collectively

Ref. 135 Ref. 136

Challenge 295 s

Ref. 137

Ref. 138

Ref. 139

Ref. 140

Page 132

^{*} The web pages cfa-www.harvard.edu/iau/lists/Closest.html and InnerPlot.html give an impression of the number of objects that almost hit the Earth every year. Without the Moon, we would have many additional catastrophes.



FIGURE 123 The phases of the Moon and of Venus, as observed from Athens in summer 2007 (© Anthony Ayiomamitis).

Challenge 296 e

known as *conic sections*. Indeed each of these curves can be produced by cutting a cone with a knife. Are you able to confirm this?

If orbits are mostly ellipses, it follows that comets *return*. The English astronomer Edmund Halley (1656–1742) was the first to draw this conclusion and to predict the return of a comet. It arrived at the predicted date in 1756, and is now named after him. The period of Halley's comet is between 74 and 80 years; the first recorded sighting was 22 centuries ago, and it has been seen at every one of its 30 passages since, the last time in 1986.


Depending on the initial energy and the initial angular momentum of the body with respect to the central planet, paths are either *elliptic*, *parabolic* or *hyperbolic*. Can you determine the conditions of the energy and the angular momentum needed for these paths to appear?

Challenge 297 ny

In practice, parabolic paths do not exist in nature. (Though some comets seem to approach this case when moving around the Sun; almost all comets follow elliptical paths). Hyperbolic paths do exist; artificial satellites follow them when they are shot towards a planet, usually with the aim of changing the direction of the satellite's journey across the solar system.

Why does the inverse square 'law' lead to conic sections? First, for two bodies, the total angular momentum L is a constant:

$$L = mr^2 \dot{\varphi} \tag{49}$$






FIGURE 124 The possible orbits due to universal gravity around a large mass (left) and a few recent examples of measured orbits (right), namely those of some extrasolar planets and of the Earth, all drawn around their respective central star, with distances given in astronomical units (© Geoffrey Marcy).



FIGURE 125 Universal gravitation also explains the obervations of Venus, the evening and morning star. In particular, universal gravitation, and the elliptical orbits it implies, explains its phases and its change of angular size. The pictures shown here were taken in 2004 and 2005. The observations can easily be made with a binocular or a small telescope (© Wahl; film available at apod.nasa.gov/apod/ap060110. html).

and therefore the motion lies in a plane. Also the energy *E* is a constant

$$E = \frac{1}{2}m\left(\frac{\mathrm{d}r}{\mathrm{d}t}\right)^2 + \frac{1}{2}m\left(r\frac{\mathrm{d}\varphi}{\mathrm{d}t}\right)^2 - G\frac{mM}{r} \ . \tag{50}$$

Challenge 298 ny Together, the two equations imply that

$$r = \frac{L^2}{Gm^2M} \frac{1}{1 + \sqrt{1 + \frac{2EL^2}{G^2m^3M^2}\cos\varphi}} .$$
 (51)

Now, any curve defined by the general expression

$$r = \frac{C}{1 + e\cos\varphi} \quad \text{or} \quad r = \frac{C}{1 - e\cos\varphi} \tag{52}$$

is an ellipse for 0 < e < 1, a parabola for e = 1 and a hyperbola for e > 1, one focus being at the origin. The quantity e, called the *eccentricity*, describes how squeezed the curve is. In other words, a body in orbit around a central mass follows a conic section.

In all orbits, also the heavy mass moves. In fact, both bodies orbit around the common centre of mass. Both bodies follow the same type of curve (ellipsis, parabola or hyperbola), but the sizes of the two curves differ.

If more than two objects move under mutual gravitation, many additional possibilities for motions appear. The classification and the motions are quite complex. In fact, this so-called *many-body problem* is still a topic of research, and the results are mathematically fascinating. Let us look at a few examples.

When several planets circle a star, they also attract each other. Planets thus do not move in perfect ellipses. The largest deviation is a perihelion shift, as shown in Figure 99. It is observed for Mercury and a few other planets, including the Earth. Other deviations from elliptical paths appear during a single orbit. In 1846, the observed deviations of the motion of the planet Uranus from the path predicted by universal gravity were used to predict the existence of another planet, Neptune, which was discovered shortly afterwards.

We have seen that mass is always positive and that gravitation is thus always attractive; there is no *antigravity*. Can gravity be used for *levitation* nevertheless, maybe using more than two bodies? Yes; there are two examples.* The first are the geostationary satellites, which are used for easy transmission of television and other signals from and towards Earth.

The *Lagrangian libration points* are the second example. Named after their discoverer, these are points in space near a two-body system, such as Moon–Earth or Earth–Sun, in which small objects have a stable equilibrium position. An overview is given in Figure 126. Can you find their precise position, remembering to take rotation into account? There are three additional Lagrangian points on the Earth–Moon line (or Sun–planet line). How many of them are stable?

There are thousands of asteroids, called *Trojan asteroids*, at and around the Lagrangian points of the Sun–Jupiter system. In 1990, a Trojan asteroid for the Mars–Sun system was discovered. Finally, in 1997, an 'almost Trojan' asteroid was found that follows the Earth on its way around the Sun (it is only transitionary and follows a somewhat more complex orbit). This 'second companion' of the Earth has a diameter of 5 km. Similarly, on the main Lagrangian points of the Earth–Moon system a high concentration of dust has been observed.

To sum up, the single equation $a = -GMr/r^3$ correctly describes a large number of phenomena in the sky. The first person to make clear that this expression describes *everything* happening in the sky was Pierre Simon Laplace** in his famous treatise *Traité*

Chanerige 255 e

Challenge 300 s


Page 97

Challenge 301 d

Vol. III, page 175

^{*} Levitation is discussed in detail in the section on electrodynamics.

^{**} Pierre Simon Laplace (b. 1749 Beaumont-en-Auge, d. 1827 Paris), important French mathematician. His






FIGURE 126 Geostationary satellites (left) and the main stable Lagrangian points (right).

de mécanique céleste. When Napoleon told him that he found no mention about the creator in the book, Laplace gave a famous, one sentence summary of his book: Je n'ai pas eu besoin de cette hypothèse. 'I had no need for this hypothesis.' In particular, Laplace studied the stability of the solar system, the eccentricity of the lunar orbit, and the eccentricities of the planetary orbits, always getting full agreement between calculation and measurement.

These results are quite a feat for the simple expression of universal gravitation; they also explain why it is called 'universal'. But how *accurate* is the formula? Since astronomy allows the most precise measurements of gravitational motion, it also provides the most stringent tests. In 1849, Urbain Le Verrier concluded after intensive study that there was only one known example of a discrepancy between observation and universal gravity, namely one observation for the planet Mercury. (Nowadays a few more are known.) The point of least distance to the Sun of the orbit of planet Mercury, its *perihelion*, changes at a rate that is slightly less than that predicted: he found a tiny difference, around 38 " per century. (This was corrected to 43 " per century in 1882 by Simon Newcomb.) Le Verrier thought that the difference was due to a planet between Mercury and the Sun, *Vulcan*, which he chased for many years without success. Indeed, Vulcan does not exist. The correct explanation of the difference had to wait for Albert Einstein.

Vol. II, page 163

Ref. 142

TIDES

Ref. 143 Why do physics texts always talk about tides? Because, as general relativity will show, tides prove that space is curved! It is thus useful to study them in a bit more detail. Figure 127 how striking tides can be. Gravitation explains the sea tides as results of the attraction of the ocean water by the Moon and the Sun. Tides are interesting; even though the amplitude of the tides is only about 0.5 m on the open sea, it can be up to 20 m at

treatise appeared in five volumes between 1798 and 1825. He was the first to propose that the solar system was formed from a rotating gas cloud, and one of the first people to imagine and explore black holes.

FIGURE 127 Tides at Saint-Valéry en Caux on September 20, 2005 (© Gilles Régnier).




FIGURE 128 Tidal deformations due to gravity.




FIGURE 129 The origin of tides.

Challenge 302 s Ref. 52

Ref. 144

special places near the coast. Can you imagine why? The *soil* is also lifted and lowered by the Sun and the Moon, by about 0.3 m, as satellite measurements show. Even the *atmosphere* is subject to tides, and the corresponding pressure variations can be filtered out from the weather pressure measurements.

Tides appear for any *extended* body moving in the gravitational field of another. To understand the origin of tides, picture a body in orbit, like the Earth, and imagine its components, such as the segments of Figure 128, as being held together by springs. Universal

gravity implies that orbits are slower the more distant they are from a central body. As a result, the segment on the outside of the orbit would like to be slower than the central one; but it is *pulled* by the rest of the body through the springs. In contrast, the inside segment would like to orbit more rapidly but is *retained* by the others. Being slowed down, the inside segments want to fall towards the Sun. In sum, both segments feel a pull away from the centre of the body, limited by the springs that stop the deformation. Therefore, *extended bodies are deformed in the direction of the field inhomogeneity.*

For example, as a result of tidal forces, the Moon always has (roughly) the same face to the Earth. In addition, its radius in direction of the Earth is larger by about 5 m than the radius perpendicular to it. If the inner springs are too weak, the body is torn into pieces; in this way a *ring* of fragments can form, such as the asteroid ring between Mars and Jupiter or the rings around Saturn.

Let us return to the Earth. If a body is surrounded by water, it will form bulges in the direction of the applied gravitational field. In order to measure and compare the strength of the tides from the Sun and the Moon, we reduce tidal effects to their bare minimum. As shown in Figure 129, we can study the deformation of a body due to gravity by studying the deformation of four pieces. We can study it in free fall, because orbital motion and free fall are equivalent. Now, gravity makes some of the pieces approach and others diverge, depending on their relative positions. The figure makes clear that the strength of the deformation – water has no built-in springs – depends on the change of gravitational acceleration with distance; in other words, the *relative* acceleration that leads to the tides is proportional to the derivative of the gravitational acceleration.

Page 383 Using the numbers from Appendix B, the gravitational accelerations from the Sun and the Moon measured on Earth are

$$a_{\text{Sun}} = \frac{GM_{\text{Sun}}}{d_{\text{Sun}}^2} = 5.9 \text{ mm/s}^2$$

$$a_{\text{Moon}} = \frac{GM_{\text{Moon}}}{d_{\text{Moon}}^2} = 0.033 \text{ mm/s}^2$$
(53)

and thus the attraction from the Moon is about 178 times weaker than that from the Sun.

When two nearby bodies fall near a large mass, the relative acceleration is proportional to their distance, and follows da = da/dr dr. The proportionality factor $da/dr = \nabla a$, called the *tidal acceleration* (gradient), is the true measure of tidal effects. Near a large spherical mass M, it is given by

Challenge 303 e

$$\frac{da}{dr} = -\frac{2GM}{r^3} \tag{54}$$

which yields the values

$$\frac{da_{\text{Sun}}}{dr} = -\frac{2GM_{\text{Sun}}}{d_{\text{Sun}}^3} = -0.8 \cdot 10^{-13} / \text{s}^2$$

$$\frac{da_{\text{Moon}}}{dr} = -\frac{2GM_{\text{Moon}}}{d_{\text{Moon}}^3} = -1.7 \cdot 10^{-13} / \text{s}^2 .$$
(55)




FIGURE 130 The Earth, the Moon and the friction effects of the tides (not to scale).

In other words, despite the much weaker pull of the Moon, its tides are predicted to be over *twice as strong* as the tides from the Sun; this is indeed observed. When Sun, Moon and Earth are aligned, the two tides add up; these so-called *spring tides* are especially strong and happen every 14.8 days, at full and new moon.

Tides lead to a pretty puzzle. Moon tides are much stronger than Sun tides. This implies that the Moon is much denser than the Sun. Why?

Tides also produce *friction*, as shown in Figure 130. The friction leads to a slowing of the Earth's rotation. Nowadays, the slowdown can be measured by precise clocks (even though short time variations due to other effects, such as the weather, are often larger). The results fit well with fossil results showing that 400 million years ago, in the Devonian period, a year had 400 days, and a day about 22 hours. It is also estimated that 900 million years ago, each of the 481 days of a year were 18.2 hours long. The friction at the basis of this slowdown also results in an increase in the distance of the Moon from the Earth by about 3.8 cm per year. Are you able to explain why?

As mentioned above, the tidal motion of the soil is also responsible for the triggering of *earthquakes*. Thus the Moon can have also dangerous effects on Earth. (Unfortunately, knowing the mechanism does not allow predicting earthquakes.) The most fascinating example of tidal effects is seen on Jupiter's satellite Io. Its tides are so strong that they induce intense volcanic activity, as shown in Figure 131, with eruption plumes as high as 500 km. If tides are even stronger, they can destroy the body altogether, as happened to the body between Mars and Jupiter that formed the planetoids, or (possibly) to the moons that led to Saturn's rings.

In summary, tides are due to relative accelerations of nearby mass points. This has an

Challenge 304 s

Ref. 104

Vol. II, page 211

Challenge 305 s



FIGURE 131 A spectacular result of tides: volcanism on Io (NASA).




FIGURE 132 Particles falling side-by-side approach over time.




FIGURE 133 Masses bend light.

Vol. II, page 124

Vol. II, page 174

important consequence. In the chapter on general relativity we will find that time multiplied by the speed of light plays the same role as length. Time then becomes an additional dimension, as shown in Figure 132. Using this similarity, two free particles moving in the same direction correspond to parallel lines in space-time. Two particles falling side-by-side also correspond to parallel lines. Tides show that such particles approach each other. In other words, tides imply that parallel lines approach each other. But parallel lines can approach each other *only* if space-time is curved. In short, tides imply *curved* space-time and space. This simple reasoning could have been performed in the eighteenth century; however, it took another 200 years and Albert Einstein's genius to uncover it.

CAN LIGHT FALL?

Oie Maxime, jederzeit selbst zu denken, ist die Aufklärung.

Immanuel Kant*

Vol. II, page 14

Towards the end of the seventeenth century people discovered that light has a finite velocity – a story which we will tell in detail later. An entity that moves with infinite velocity cannot be affected by gravity, as there is no time to produce an effect. An entity with a finite speed, however, should feel gravity and thus fall.

Ref. 145

Does its speed increase when light reaches the surface of the Earth? For almost three centuries people had no means of detecting any such effect; so the question was not investigated. Then, in 1801, the Prussian astronomer Johann Soldner (1776–1833) was the first to put the question in a different way. Being an astronomer, he was used to measuring stars and their observation angles. He realized that light passing near a massive body would be *deflected* due to gravity.

Soldner studied a body on a hyperbolic path, moving with velocity c past a spherical mass M at distance b (measured from the centre), as shown in Figure 133. Soldner deduced the deflection angle

Challenge 306 ny

$$\alpha_{\text{univ. grav.}} = \frac{2}{b} \frac{GM}{c^2} . \tag{56}$$

Vol. II, page 161

One sees that the angle is largest when the motion is just grazing the mass M. For light deflected by the mass of the Sun, the angle turns out to be at most a tiny 0.88 "= $4.3 \,\mu$ rad. In Soldner's time, this angle was too small to be measured. Thus the issue was forgotten. Had it been pursued, general relativity would have begun as an experimental science, and not as the theoretical effort of Albert Einstein! Why? The value just calculated is *different* from the measured value. The first measurement took place in 1919;** it found the correct dependence on the distance, but found a deflection of up to 1.75 ", exactly double that of expression (56). The reason is not easy to find; in fact, it is due to the curvature of space, as we will see. In summary, light can fall, but the issue hides some surprises.

What is mass? - Again

Mass describes how an object interacts with others. In our walk, we have encountered two of its aspects. *Inertial mass* is the property that keeps objects moving and that offers resistance to a change in their motion. *Gravitational mass* is the property responsible for the acceleration of bodies nearby (the active aspect) or of being accelerated by objects nearby (the passive aspect). For example, the active aspect of the mass of the Earth determines the surface acceleration of bodies; the passive aspect of the bodies allows us to weigh them in order to measure their mass using distances only, e.g. on a scale or a balance. The gravitational mass is the basis of *weight*, the difficulty of lifting things.***

Is the gravitational mass of a body equal to its inertial mass? A rough answer is given by the experience that an object that is difficult to move is also difficult to lift. The simplest

^{*} The maxim to think at all times for oneself is the enlightenment.

^{**} By the way, how would you measure the deflection of light near the bright Sun?

^{***} What are the values shown by a balance for a person of 85 kg juggling three balls of 0.3 kg each?

experiment is to take two bodies of different masses and let them fall. If the acceleration is the same for all bodies, inertial mass is equal to (passive) gravitational mass, because in the relation $ma = \nabla(GMm/r)$ the left-hand m is actually the inertial mass, and the right-hand m is actually the gravitational mass.

But in the seventeenth century Galileo had made widely known an even older argument showing without a single experiment that the gravitational acceleration is indeed the same for all bodies. If larger masses fell more rapidly than smaller ones, then the following paradox would appear. Any body can be seen as being composed of a large fragment attached to a small fragment. If small bodies really fell less rapidly, the small fragment would slow the large fragment down, so that the complete body would have to fall *less* rapidly than the larger fragment (or break into pieces). At the same time, the body being larger than its fragment, it should fall *more* rapidly than that fragment. This is obviously impossible: all masses must fall with the same acceleration.

Many accurate experiments have been performed since Galileo's original discussion. In all of them the independence of the acceleration of free fall from mass and material composition has been confirmed with the precision they allowed. In other words, as far as we can tell, gravitational mass and inertial mass are *identical*. What is the origin of this mysterious equality?

This so-called 'mystery' is a typical example of disinformation, now common across the whole world of physics education. Let us go back to the definition of mass as a negative inverse acceleration ratio. We mentioned that the physical origins of the accelerations do not play a role in the definition because the origin does not appear in the expression. In other words, the value of the mass is by definition independent of the interaction. That means in particular that inertial mass, based on electromagnetic interaction, and gravitational mass are identical *by definition*.

We also note that we have never defined a separate concept of 'passive gravitational mass'. The mass being accelerated by gravitation is the inertial mass. Worse, there is no way to define a 'passive gravitational mass'. Try it! All methods, such as weighing an object, cannot be distinguished from those that determine inertial mass from its reaction to acceleration. Indeed, all methods of measuring mass use non-gravitational mechanisms. Scales are a good example.

If the 'passive gravitational mass' were different from the inertial mass, we would have strange consequences. For those bodies for which it were different we would get into trouble with energy conservation. Also assuming that 'active gravitational mass' differs from inertial mass gets us into trouble.

Another way of looking at the issue is as follows. How could 'gravitational mass' differ from inertial mass? Would the difference depend on relative velocity, time, position, composition or on mass itself? Each of these possibilities contradicts either energy or momentum conservation.

No wonder that all measurements confirm the equality of all mass types. The issue is usually resurrected in general relativity, with no new results. 'Both' masses remain equal. *Mass is a unique property of bodies*. Another, deeper issue remains, though. What is the *origin* of mass? Why does it exist? This simple but deep question cannot be answered by classical physics. We will need some patience to find out.

Ref. 146

Page 92

Challenge 309 s

Vol. II, page 144




FIGURE 134 Brooms fall more rapidly than stones (© Luca Gastaldi).

CURIOSITIES AND FUN CHALLENGES ABOUT GRAVITATION

Fallen ist weder gefährlich noch eine Schande; Liegen bleiben ist beides.*

Konrad Adenauer

Gravity on the Moon is only one sixth of that on the Earth. Why does this imply that it is difficult to walk quickly and to run on the Moon (as can be seen in the TV images recorded there)?

* *

Is the acceleration due to gravity constant? Not really. Every day, it is estimated that 10^8 kg of material fall onto the Earth in the form of meteorites and asteroids. Nevertheless, it is unknown whether the mass of the Earth increases with time (due to collection of meteorites and cosmic dust) or decreases (due to gas loss). If you find a way to settle the issue, publish it.

* *

Incidentally, discovering objects hitting the Earth is not at all easy. Astronomers like to point out that an asteroid as large as the one that led to the extinction of the dinosaurs could hit the Earth without any astronomer noticing in advance, if the direction is slightly unusual, such as from the south, where few telescopes are located.

* *

Several humans have survived free falls from aeroplanes for a thousand metres or more, even though they had no parachute. A minority of them even did so without any harm at all. How was this possible?

Challenge 310 s

* *

Imagine that you have twelve coins of identical appearance, of which one is a forgery. The forged one has a different mass from the eleven genuine ones. How can you decide

^{* &#}x27;Falling is neither dangerous nor a shame; to keep lying is both.' Konrad Adenauer (b. 1876 Köln, d. 1967 Rhöndorf), West German Chancellor.

which is the forged one and whether it is lighter or heavier, using a simple balance only three times? Challenge 311 e

> You have nine identically-looking spheres, all of the same mass, except one, which is heavier. Can you determine which one, using the balance only two times?

For a physicist, antigravity is repulsive gravity; it does not exist in nature. Nevertheless, the term 'antigravity' is used incorrectly by many people, as a short search on the internet shows. Some people call any effect that overcomes gravity, 'antigravity'. However, this definition implies that tables and chairs are antigravity devices. Following the definition, most of the wood, steel and concrete producers are in the antigravity business. The internet definition makes absolutely no sense.

What is the cheapest way to switch gravity off for 25 seconds? Challenge 312 s

Do all objects on Earth fall with the same acceleration of 9.8 m/s², assuming that air resistance can be neglected? No; every housekeeper knows that. You can check this by yourself. As shown in Figure 134, a broom angled at around 35° hits the floor before a

stone, as the sounds of impact confirm. Are you able to explain why? Challenge 313 s

Also bungee jumpers are accelerated more strongly than g. For a bungee cord of mass mand a jumper of mass M, the maximum acceleration a is

$$a = g\left(1 + \frac{1}{8}\frac{m}{M}\left(4 + \frac{m}{M}\right)\right) . (57)$$

Can you deduce the relation from Figure 135?

Guess: What is the weight of a ball of cork with a radius of 1 m? Challenge 315 s

Guess: One thousand 1 mm diameter steel balls are collected. What is the mass? Challenge 316 s

How can you use your observations made during your travels with a bathroom scale to show that the Earth is not flat? Challenge 317 s

* *

Both the Earth and the Moon attract bodies. The centre of mass of the Earth-Moon system is 4800 km away from the centre of the Earth, quite near its surface. Why do bodies on Earth still fall towards the centre of the Earth?



FIGURE 135 The starting situation for a bungee jumper.




FIGURE 136 An honest balance?

* *

Challenge 319 e confirm

Page 178 Galileo

Does every spherical body fall with the same acceleration? No. If the weight of the object is comparable to that of the Earth, the distance decreases in a different way. Can you confirm this statement? Figure 136 shows a related puzzle. What then is wrong about Galileo's argument about the constancy of acceleration of free fall?

* *

What is the fastest speed that a human can achieve making use of gravitational acceleration? There are various methods that try this; a few are shown in Figure 137. Terminal speed of free falling skydivers can be even higher, but no reliable record speed value exists. The last word is not spoken yet, as all these records will be surpassed in the coming years. It is important to require normal altitude; at stratospheric altitudes, speed values can be four times the speed values at low altitude.

Vol. II, page 124

* *

It is easy to lift a mass of a kilogram from the floor on a table. Twenty kilograms is harder. A thousand is impossible. However, $6 \cdot 10^{24}$ kg is easy. Why?

Challenge 320 s

Page 175

* 1

Challenge 321 ny

The friction between the Earth and the Moon slows down the rotation of both. The Moon stopped rotating millions of years ago, and the Earth is on its way to doing so as well. When the Earth stops rotating, the Moon will stop moving away from Earth. How far will the Moon be from the Earth at that time? Afterwards however, even further in the future, the Moon will move back towards the Earth, due to the friction between the Earth–Moon system and the Sun. Even though this effect would only take place if the Sun burned for







FIGURE 137 Reducing air resistance increases the terminal speed: left, the 2007 speed skiing world record holder Simone Origone with 69.83 m/s and right, the 2007 speed world record holder for bicycles on snow Éric Barone with 61.73 m/s (© Simone Origone, Éric Barone).

Challenge 322 s ever, which is known to be false, can you explain it?

* *

When you run towards the east, you *lose weight*. There are two different reasons for this: the 'centrifugal' acceleration increases so that the force with which you are pulled down diminishes, and the Coriolis force appears, with a similar result. Can you estimate the size of the two effects?

Challenge 323 ny

* *

What is the relation between the time a stone takes falling through a distance l and the time a pendulum takes swinging though half a circle of radius l? (This problem is due to Galileo.) How many digits of the number π can one expect to determine in this way?

* *

Challenge 325 s

Ref. 147

Why can a spacecraft accelerate through the *slingshot effect* when going round a planet, despite momentum conservation? It is speculated that the same effect is also the reason for the few exceptionally fast stars that are observed in the galaxy. For example, the star HE0457-5439 moves with 720 km/s, which is much higher than the 100 to 200 km/s of most stars in the Milky way. It seems that the role of the accelerating centre was taken by a black hole.

* *

Ref. 148 The orbit of a planet around the Sun has many interesting properties. What is the hodo-Challenge 326 s graph of the orbit? What is the hodograph for parabolic and hyperbolic orbits?




FIGURE 138 The four satellites of Jupiter discovered by Galileo and their motion (© Robin Scagell).

* *

The *Galilean satellites* of Jupiter, shown in Figure 138, can be seen with small amateur telescopes. Galileo discovered them in 1610 and called them the *Medicean satellites*. (Today, they are named, in order of increasing distance from Jupiter, as Io, Europa, Ganymede and Callisto.) They are almost mythical objects. They were the first bodies found that obviously did not orbit the Earth; thus Galileo used them to deduce that the Earth is not at the centre of the universe. The satellites have also been candidates to be the first *standard clock*, as their motion can be predicted to high accuracy, so that the 'standard time' could be read off from their position. Finally, due to this high accuracy, in 1676, the speed of light was first measured with their help, as told in the section on special relativity.

Vol. II, page 15

* *

Challenge 327 s

A simple, but difficult question: if all bodies attract each other, why don't or didn't all stars fall towards each other? Indeed, the inverse square expression of universal gravity has a limitation: it does not allow one to make sensible statements about the matter in the universe. Universal gravity predicts that a homogeneous mass distribution is unstable; indeed, an inhomogeneous distribution is observed. However, universal gravity does not predict the average mass density, the darkness at night, the observed speeds of the distant galaxies, etc. In fact, 'universal' gravity does not explain or predict a single property of the universe. To do this, we need general relativity.

Vol. II, page 194

* *

The acceleration g due to gravity at a depth of 3000 km is $10.05 \,\mathrm{m/s^2}$, over 2 % more than at the surface of the Earth. How is this possible? Also, on the Tibetan plateau, g is influenced by the material below it.




FIGURE 139 Which of the two Moon paths is correct?

* *

When the Moon circles the Sun, does its path have sections *concave* towards the Sun, as shown at the right of Figure 139, or not, as shown on the left? (Independent of this issue, both paths in the diagram disguise that the Moon path does *not* lie in the same plane as the path of the Earth around the Sun.)

* *

You can prove that objects *attract each other* (and that they are not only attracted by the Earth) with a simple experiment that anybody can perform at home, as described on the www.fourmilab.ch/gravitation/foobar website.

* *

It is instructive to calculate the *escape velocity* from the Earth, i.e., that velocity with which a body must be thrown so that it never falls back. It turns out to be around 11 km/s. (This was called the *second cosmic velocity* in the past; the *first cosmic velocity* was the name given to the lowest speed for an orbit, 7.9 km/s.) The exact value of the escape velocity depends on the latitude of the thrower, and on the direction of the throw. (Why?)

Challenge 329 e

What is the escape velocity for the solar system? (It was once called the *third cosmic velocity*.) By the way, the escape velocity from our galaxy is over 500 km/s. What would happen if a planet or a system were so heavy that its escape velocity would be larger than the speed of light?

Challenge 330 s

* *

Challenge 331 s What is the largest asteroid one can escape from by jumping?

* *



FIGURE 140 The analemma over Delphi, taken between January and December 2002 (© Anthony Ayiomamitis).

For bodies of irregular shape, the centre of gravity of a body is *not* the same as the centre of mass. Are you able to confirm this? (Hint: Find and use the simplest example possible.)

* *

Can gravity produce repulsion? What happens to a small test body on the inside of a large C-shaped mass? Is it pushed towards the centre of mass?

* *

Ref. 150 The *shape* of the Earth is not a sphere. As a consequence, a plumb line usually does not point to the centre of the Earth. What is the largest deviation in degrees?

* *

If you look at the sky every day at 6 a.m., the Sun's position varies during the year. The result of photographing the Sun on the same film is shown in Figure 140. The curve, called the analemma, is due to two combined effects: the inclination of the Earth's axis and the elliptical shape of the Earth's orbit around the Sun. The top and the (hidden) bottom points of the analemma correspond to the solstices. How does the analemma look if photographed every day at local noon? Why is it not a straight line pointing exactly south?

The constellation in which the Sun stands at noon (at the centre of the time zone) is supposedly called the 'zodiacal sign' of that day. Astrologers say there are twelve of them,

Challenge 335 s

Challenge 333 ny

Page 138

Ref. 151

namely Aries, Taurus, Gemini, Cancer, Leo, Virgo, Libra, Scorpius, Sagittarius, Capricornus, Aquarius and Pisces and that each takes (quite precisely) a twelfth of a year or a twelfth of the ecliptic. Any check with a calendar shows that at present, the midday Sun is never in the zodiacal sign during the days usually connected to it. The relation has shifted by about a month since it was defined, due to the precession of the Earth's axis. A check with a map of the star sky shows that the twelve constellations do not have the same length and that on the ecliptic there are fourteen of them, not twelve. There is *Ophiuchus* or *Serpentarius*, the serpent bearer constellation, between Scorpius and Sagittarius, and *Cetus*, the whale, between Aquarius and Pisces. In fact, not a single astronomical statement about zodiacal signs is correct. To put it clearly, astrology, in contrast to its name, is *not* about stars. (In one language, the term for 'crook' is derived from the word 'astrologer'.)

* *

For a long time, it was thought that there is no additional planet in our solar system outside Neptune and Pluto, because their orbits show no disturbances from another body. Today, the view has changed. It is known that there are only eight planets: Pluto is not a planet, but the first of a set of smaller objects in the so-called Kuiper belt. Kuiper belt objects are regularly discovered; around 80 are known today.

In 2003, two major Kuiper objects were discovered; one, called *Sedna*, is almost as large as Pluto, the other, called *Eris*, is even larger than Pluto and has a moon. Both have strongly elliptical orbits (see Figure 141). Since Pluto and Eris, like the asteroid Ceres, have cleaned their orbit from debris, these three objects are now classified as *dwarf planets*.

* *

In astronomy new examples of motion are regularly discovered even in the present century. Sometimes there are also false alarms. One example was the alleged fall of *mini comets* on the Earth. They were supposedly made of a few dozen kilograms of ice, hitting the Earth every few seconds. It is now known not to happen.

* *

Universal gravity allows only elliptical, parabolic or hyperbolic orbits. It is impossible for a small object approaching a large one to be captured. At least, that is what we have learned so far. Nevertheless, all astronomy books tell stories of capture in our solar system; for example, several outer satellites of Saturn have been captured. How is this possible?

Challenge 336 s

Ref. 154

* *

How would a tunnel have to be shaped in order that a stone would fall through it without touching the walls? (Assume constant density.) If the Earth did not rotate, the tunnel would be a straight line through its centre, and the stone would fall down and up again, in a oscillating motion. For a rotating Earth, the problem is much more difficult. What is the shape when the tunnel starts at the Equator?

Challenge 337 s




FIGURE 141 The orbit of Sedna in comparison with the orbits of the planets in the solar system (NASA).

The International Space Station circles the Earth every 90 minutes at an altitude of about 380 km. You can see where it is from the website www.heavens-above.com. By the way, whenever it is just above the horizon, the station is the third brightest object in the night sky, superseded only by the Moon and Venus. Have a look at it.

Challenge 338 e

* *

Challenge 339 s

Vol. II, page 27

Is it true that the centre of mass of the solar system, its barycentre, is always inside the Sun? Even though a star or the Sun move very little when planets move around them, this motion can be detected with precision measurements making use of the Doppler effect for light or radio waves. Jupiter, for example, produces a speed change of 13 m/s in the Sun, the Earth 1 m/s. The first planets outside the solar system, around the pulsar PSR1257+12 and the star Pegasi 51, was discovered in this way, in 1992 and 1995. In the meantime, over 400 planets have been discovered with this and other methods. Some






FIGURE 142 The vanishing of gravitational force inside a spherical shell of matter.

have even masses comparable to that of the Earth.

* *

Not all points on the Earth receive the same number of daylight hours during a year. The Challenge 340 d effects are difficult to spot, though. Can you find one?

* *

Can the phase of the Moon have a measurable effect on the human body, for example through tidal effects?

* *

There is an important difference between the heliocentric system and the old idea that all planets turn around the Earth. The heliocentric system states that certain planets, such as Mercury and Venus, can be *between* the Earth and the Sun at certain times, and *behind* the Sun at other times. In contrast, the geocentric system states that they are always in between. Why did such an important difference not immediately invalidate the geocentric system?

Challenge 342 s syst

* *

The strangest reformulation of the description of motion given by $m\mathbf{a} = \nabla U$ is the almost absurd looking equation

$$\nabla v = \mathrm{d}\boldsymbol{v}/\mathrm{d}s \tag{58}$$

where s is the motion path length. It is called the *ray form* of Newton's equation of motion. Challenge 343 s Can you find an example of its application?

* *

Seen from Neptune, the size of the Sun is the same as that of Jupiter seen from the Earth Challenge 344 s at the time of its closest approach. True?

* *

The gravitational acceleration for a particle inside a spherical shell is zero. The vanishing Ref. 156 of gravity in this case is independent of the particle shape and its position, and indepen-

Challenge 345 s

dent of the thickness of the shell. Can you find the argument using Figure 142? This works only because of the $1/r^2$ dependence of gravity. Can you show that the result does not hold for non-spherical shells? Note that the vanishing of gravity inside a spherical shell usually does not hold if other matter is found outside the shell. How could one eliminate the effects of outside matter?

Challenge 346 s

What is gravity? This is not a simple question. In 1690, Nicolas Fatio de Duillier and in 1747, Georges-Louis Lesage proposed an explanation for the $1/r^2$ dependence. Lesage ar-Ref. 157 gued that the world is full of small particles - he called them 'corpuscules ultra-mondains' - flying around randomly and hitting all objects. Single objects do not feel the hits, since they are hit continuously and randomly from all directions. But when two objects are near to each other, they produce shadows for part of the flux to the other body, resulting in an attraction. Can you show that such an attraction has a $1/r^2$ dependence?

Challenge 347 e

Ref. 2

However, Lesage's proposal has a number of problems. The argument only works if the collisions are inelastic. (Why?) However, that would mean that all bodies would heat up with time, as Jean-Marc Lévy-Leblond explains.

There are two additional problems with the idea of Lesage. First, a moving body in free space would be hit by more or faster particles in the front than in the back; as a result, the body should be decelerated. Second, gravity would depend on size, but in a strange way. In particular, three bodies lying on a line should not produce shadows, as

no such shadows are observed; but the naive model predicts such shadows.

Despite all the criticisms, this famous idea has regularly resurfaced in physics ever since, even though such particles have never been found. Only in the final part of our mountain ascent will we settle the issue of the origin of gravitation.

For which bodies does gravity decrease as you approach them?

Could one put a satellite into orbit using a cannon? Does the answer depend on the Challenge 349 s direction in which one shoots?

Two computer users share experiences. 'I threw my Pentium III and Pentium IV out of the window. 'And?' 'The Pentium III was faster.'

How often does the Earth rise and fall when seen from the Moon? Does the Earth show Challenge 350 s phases?

What is the weight of the Moon? How does it compare with the weight of the Alps?




FIGURE 143 The motion of the planetoids compared to that of the planets (Shockwave animation © Hans-Christian Greier)

* *

Owing to the slightly flattened shape of the Earth, the source of the Mississippi is about 20 km nearer to the centre of the Earth than its mouth; the water effectively runs uphill. How can this be?

Challenge 352 s

Ref. 158

Page 357

* *

If a star is made of high density material, the speed of a planet orbiting near to it could be greater than the speed of light. How does nature avoid this strange possibility?

* *

What will happen to the solar system in the future? This question is surprisingly hard to answer. The main expert of this topic, French planetary scientist Jacques Laskar, simulated a few hundred million years of evolution using computer-aided calculus. He found that the planetary orbits are stable, but that there is clear evidence of chaos in the evolution of the solar system, at a small level. The various planets influence each other in subtle and still poorly understood ways. Effects in the past are also being studied, such as the energy change of Jupiter due to its ejection of smaller asteroids from the solar system, or energy gains of Neptune. There is still a lot of research to be done in this field.

* *

One of the open problems of the solar system is the description of planet distances discovered in 1766 by Johann Daniel Titius (1729–1796) and publicized by Johann Elert Bode (1747–1826). Titius discovered that planetary distances d from the Sun can be approximated by

$$d = a + 2^n b$$
 with $a = 0.4 \text{ AU}$, $b = 0.3 \text{ AU}$ (59)

where distances are measured in astronomical units and n is the number of the planet. The resulting approximation is compared with observations in Table 27.

Motion Mountain – The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christoph Schiller November 1997–March

TABLE 27 An unexplained property of nature: planet distances and the values resulting from the Titius–Bode rule.

PLANET	n	PREDICTED	MEASURED	
		DISTANCE IN AU		
Mercury	-∞	0.4	0.4	
Venus	0	0.7	0.7	
Earth	1	1.0	1.0	
Mars	2	1.6	1.5	
Planetoids	3	2.8	2.2 to 3.2	
Jupiter	4	5.2	5.2	
Saturn	5	10.0	9.5	
Uranus	6	19.6	19.2	
Neptune	7	38.8	30.1	
Pluto	8	77.2	39.5	

Interestingly, the last three planets, as well as the planetoids, were discovered *after* Bode's and Titius' deaths; the rule had successfully predicted Uranus' distance, as well as that of the planetoids. Despite these successes – and the failure for the last two planets – nobody has yet found a model for the formation of the planets that explains Titius' rule. The large satellites of Jupiter and of Uranus have regular spacing, but not according to the Titius–Bode rule.

Ref. 159 Ref. 160 Explaining or disproving the rule is one of the challenges that remains in classical mechanics. Some researchers maintain that the rule is a consequence of scale invariance, others maintain that it is a accident or even a red herring. The last interpretation is also suggested by the non-Titius–Bode behaviour of practically all extrasolar planets. The issue is not closed.

* *

Around 3000 years ago, the Babylonians had measured the orbital times of the seven celestial bodies. Ordered from longest to shortest, they wrote them down in Table 28.

The Babylonians also introduced the week and the division of the day into 24 hours. The Babylonians dedicated every one of the 168 hours of the week to a celestial body, following the order of Table 28. They also dedicated the whole day to that celestial body that corresponds to the first hour of that day. The first day of the week was dedicated to Saturn; the present ordering of the other days of the week then follows from Table 28. This story was told by Cassius Dio (c. 160 to c. 230). Towards the end of Antiquity, the ordering was taken up by the Roman empire. In Germanic languages, including English, the Latin names of the celestial bodies were replaced by the corresponding Germanic gods. The order Saturday, Sunday, Monday, Tuesday, Wednesday, Thursday and Friday is thus a consequence of both the astronomical measurements and the astrological superstitions of the ancients.

Challenge 354 e Ref. 161

TABLE 28 The orbital periods known to the Babylonians.

Ворч	PERIOD	
Saturn	29 a	
Jupiter	12 a	
Mars	687 d	
Sun	365 d	
Venus	224 d	
Mercury	88 d	
Moon	29 d	

* *

In 1722, the great mathematician Leonhard Euler made a mistake in his calculation that led him to conclude that if a tunnel, or better, a deep hole were built from one pole of the Earth to the other, a stone falling into it would arrive at the Earth's centre and then immediately turn and go back up. Voltaire made fun of this conclusion for many years. Can you correct Euler and show that the real motion is an oscillation from one pole to the other, and can you calculate the time a pole-to-pole fall would take (assuming homogeneous density)?

Challenge 355 s

Challenge 356 s

chancing c 550 i

Challenge 357 ny

Ref. 162

What would be the oscillation time for an arbitrary straight surface-to-surface tunnel of length *l*, thus *not* going from pole to pole?

The previous challenges circumvented the effects of the Earth's rotation. The topic becomes much more interesting if rotation is included. What would be the shape of a tunnel so that a stone falling through it never touches the wall?

* *

Figure 144 shows a photograph of a solar eclipse taken from the Russian space station *Mir* and a photograph taken at the centre of the shadow from the Earth. Indeed, a global view of a phenomenon can be quite different from a local one. What is the speed of the shadow?

Challenge 358 s

Ref. 163

* *

In 2005, satellite measurements have shown that the water in the Amazon river presses down the land up to 75 mm more in the season when it is full of water than in the season when it is almost empty.

* *

Assume that wires existed that do not break at all. How long would such a wire have to be so that, when attached to the equator, it would stand upright in the air?

* *

Everybody knows that there are roughly two tides per day. But there are places, such as on the coast of Vietnam, where there is only one tide per day. See www.jason.oceanobs.





FIGURE 144 The solar eclipse of 11 August 1999, photographed by Jean-Pierre Haigneré, member of the Mir 27 crew, and the (enhanced) solar eclipse of 29 March 2006 (© CNES and Laurent Laveder/PixHeaven.net).

Challenge 360 ny com/html/applications/marees/marees_m2k1_fr.html. Why?

* *

It is sufficient to use the concept of centrifugal force to show that the rings of Saturn cannot be made of massive material, but must be made of separate pieces. Can you find out how?

Challenge 361 s

* *

A painting is hanging on the wall of Dr. Dolittle's waiting room. He hung up the painting using two nails, and wound the picture wire around the nails in such a way that the painting would fall if either nail were pulled out. How did Dr. Dolittle do it?

Challenge 362 e

* *

Why did Mars lose its atmosphere? Nobody knows. It has recently been shown that the solar wind is too weak for this to happen. This is one of the many open riddles of the solar system.

* *

Page 217

The observed motion due to gravity can be shown to be the simplest possible, in the following sense. If we measure change of a falling object with $\int mv^2/2 - mgh \, dt$, then a constant acceleration due to gravity *minimizes* the change in every example of fall. Can you confirm this?

Challenge 363 ny

* *

Motion due to gravity is fun: think about roller coasters. If you want to know more at how they are built, visit www.vekoma.com.

The scientific theory I like best is that the rings of Saturn are made of lost airline luggage.


Mark Russel

))

SUMMARY ON GRAVITATION

Spherical bodies of mass m attract other bodies at a distance r by inducing an acceleration towards them given by $a = Gm/r^2$. This expression, *universal gravitation*, describes snow-boarders, skiers, paragliders, athletes, couch potatoes, pendula, stones, canons, rockets, tides, eclipses, planet shapes, planet motion and much more. It is the first example of a unified description, in this case, of *how everything falls*.





CHAPTER 7

CLASSICAL MECHANICS AND THE PREDICTABILITY OF MOTION

LL those types of motion in which the only permanent property of body is mass define the field of *mechanics*. The same name is given lso to the experts studying the field. We can think of mechanics as the athletic part of physics.* Both in athletics and in mechanics only lengths, times and masses are measured.

More specifically, our topic of investigation so far is called *classical* mechanics, to distinguish it from *quantum* mechanics. The main difference is that in classical physics arbitrary small values are assumed to exist, whereas this is not the case in quantum physics. Classical mechanics is often also called *Galilean physics* or *Newtonian physics*.**

Classical mechanics states that motion is *predictable*: it thus states that there are no surprises in motion. Is this correct in all cases? Let us start with the exploration of this issue.

We know that there is more to the world than gravity. A simple observation makes the point: *friction*. Friction cannot be due to gravity, because friction is not observed in the skies, where motion follows gravity rules only.*** Moreover, on Earth, friction is not related to gravity, as you might want to check. There must be another interaction responsible for friction. We shall study it shortly. But one issue merits a discussion right away.

Challenge 364 e

SHOULD ONE USE FORCE? POWER?

The direct use of force is such a poor solution to any problem, it is generally employed only by small children and large nations.

David Friedman

Vol. II, page 153

^{*} This is in contrast to the actual origin of the term 'mechanics', which means 'machine science'. It derives from the Greek $\mu\eta\kappa\alpha\nu\dot{\eta}$, which means 'machine' and even lies at the origin of the English word 'machine' itself. Sometimes the term 'mechanics' is used for the study of motion of *solid* bodies only, excluding, e.g., hydrodynamics. This use fell out of favour in physics in the twentieth century.

^{**} The basis of classical mechanics, the description of motion using only space and time, is called *kinematics*. An example is the description of free fall by $z(t) = z_0 + v_0(t - t_0) - \frac{1}{2}g(t - t_0)^2$. The other, main part of classical mechanics is the description of motion as a consequence of interactions between bodies; it is called *dynamics*. An example of dynamics is the formula of universal gravity. The distinction between kinematics and dynamics can also be made in relativity, thermodynamics and electrodynamics.

^{***} This is not completely correct: in the 1980s, the first case of gravitational friction was discovered: the emission of gravity waves. We discuss it in detail in the chapter on general relativity.

TABLE 29 Some force values in nature.

O B S E R V A T I O N	FORCE
Value measured in a magnetic resonance force microscope	820 zN
Force needed to rip a DNA molecule apart by pulling at its two ends	600 pN
Maximum force exerted by human bite	2.1 kN
Typical peak force exerted by sledgehammer	2 kN
Force exerted by quadriceps	up to 3 kN
Force sustained by 1 cm ² of a good adhesive	up to 10 kN
Force needed to tear a good rope used in rock climbing	30 kN
Maximum force measurable in nature	$3.0 \cdot 10^{43} \mathrm{N}$

Everybody has to take a stand on this question, even students of physics. Indeed, many types of forces are used and observed in daily life. One speaks of muscular, gravitational, psychic, sexual, satanic, supernatural, social, political, economic and many others. Physicists see things in a simpler way. They call the different types of forces observed between objects *interactions*. The study of the details of all these interactions will show that, in everyday life, they are of electrical origin.

For physicists, all change is due to motion. The term force then also takes on a more restrictive definition. (*Physical*) force is defined as the *change of momentum with time*, i.e., as

$$F = \frac{\mathrm{d}\boldsymbol{p}}{\mathrm{d}t} \ . \tag{60}$$

A few measured values are listed in Figure 29. Since momentum is conserved, we can say that force measures the *flow* of momentum. If a force accelerates a body, momentum flows into it. Indeed, momentum can be imagined to be some invisible and intangible liquid. Force measures how much of this liquid flows into or out of a body per unit time.

Using the Galilean definition of linear momentum p = mv, we can rewrite the definition of force (for constant mass) as

$$F = ma, (61)$$

where F = F(t, x) is the force acting on an object of mass m and where $a = a(t, x) = \frac{dv}{dt} = \frac{d^2x}{dt^2}$ is the acceleration of the same object, that is to say its change of velocity.* The expression states in precise terms that force is what changes the *velocity* of masses. The quantity is called 'force' because it corresponds in many, but *not* all aspects to muscular force. For example, the more force is used, the further a stone can be thrown.

However, whenever the concept of force is used, it should be remembered that *physical force is different from everyday force or everyday effort*. Effort is probably best ap-

^{*} This equation was first written down by the Swiss mathematician and physicist Leonhard Euler (1707–1783) in 1747, 20 years after the death of Newton, to whom it is usually and falsely ascribed. It was Euler, one of the greatest mathematicians of all time, not Newton, who first understood that this definition of force is useful in *every* case of motion, whatever the appearance, be it for point particles or extended objects, and be it rigid, deformable or fluid bodies. Surprisingly and in contrast to frequently-made statements, equation (61) is even correct in relativity.

proximated by the concept of (physical) power, usually abbreviated P, and defined (for constant force) as

$$P = \frac{\mathrm{d}W}{\mathrm{d}t} = \mathbf{F} \cdot \mathbf{v} \tag{62}$$

in which (*physical*) work W is defined as $W = \mathbf{F} \cdot \mathbf{s}$. Physical work is a form of energy, as you might want to check. Work, as a form of energy, has to be taken into account when the conservation of energy is checked. Note that a man who walks carrying a heavy rucksack is hardly doing any work; why then does he get tired?

With the definition of work just given you can solve the following puzzles. What happens to the electricity consumption of an escalator if you walk on it instead of standing still? What is the effect of the definition of power for the salary of scientists?

When students in exams say that the force acting on a thrown stone is least at the highest point of the trajectory, it is customary to say that they are using an incorrect view, namely the so-called *Aristotelian view*, in which force is proportional to velocity. Sometimes it is even said that they are using a different concept of *state* of motion. Critics then add, with a tone of superiority, how wrong all this is. This is an example of intellectual disinformation. Every student knows from riding a bicycle, from throwing a stone or from pulling an object that increased *effort* results in increased speed. The student is right; those theoreticians who deduce that the student has a mistaken concept of *force* are wrong. In fact, instead of the *physical* concept of force, the student is just using the *everyday* version, namely effort. Indeed, the effort exerted by gravity on a flying stone is least at the highest point of the trajectory. Understanding the difference between physical force and everyday effort is the main hurdle in learning mechanics.*

Often the flow of momentum, equation (60), is not recognized as the definition of force. This is mainly due to an everyday observation: there seem to be forces without any associated acceleration or change in momentum, such as in a string under tension or in water at high pressure. When one pushes against a tree, there is no motion, yet a force is applied. If force is momentum flow, where does the momentum go? It flows into the slight deformations of the arms and the tree. In fact, when one starts pushing and thus deforming, the associated momentum change of the molecules, the atoms, or the electrons of the two bodies can be observed. After the deformation is established, and looking at even higher magnification, one can indeed find that a continuous and equal flow of momentum is going on in both directions. The nature of this flow will be clarified in our exploration of quantum theory.

As force is net momentum flow, it is only needed as a separate concept in everyday life, where it is useful in situations where net momentum flows are less than the total flows. At the microscopic level, momentum alone suffices for the description of motion. For example, the concept of *weight* describes the flow of momentum due to gravity. Thus we will hardly ever use the term 'weight' in the microscopic part of our adventure.

Before we can answer the question in the section title, on the usefulness of force and power, we need more arguments. Through its definition, the concept of force is distin-

Challenge 365 s

Challenge 366 s Challenge 367 d

Ref. 164

^{*} This stepping stone is so high that many professional physicists do not really take it themselves; this is confirmed by the innumerable comments in papers that state that physical force is defined using mass, and, at the same time, that mass is defined using force (the latter part of the sentence being a fundamental mistake).

guished clearly from 'mass', 'momentum', 'energy' and 'power'. But where do forces originate? In other words, which effects in nature have the capacity to accelerate bodies by pumping momentum into objects? Table 30 gives an overview.

FORCES, SURFACES AND CONSERVATION

We saw that force is the change of momentum. We also saw that momentum is conserved. How do these statements come together? The answer is the same for all conserved quantities. We imagine a closed surface that is the boundary of a volume in space. Conservation implies that the conserved quantity enclosed *inside* the surface can only change by flowing *through* that surface.*

All conserved quantities in nature – such as energy, linear momentum, electric charge, angular momentum – can only change by flowing through surfaces. In particular, when the momentum of a body changes, this happens through a surface. Momentum change is due to momentum flow. In other words, the concept of force always assumes a surface through which momentum flows.

▶ Force is the flow of momentum through a surface.

Ref. 272 This point is essential in understanding physical force! Every force requires a surface for its definition.

To refine your own concept of force, you can search for the relevant surface when a rope pulls a chariot, or when an arm pushes a tree, or when a car accelerates. It is also helpful to compare the definition of force with the definition of power: both are flows through surfaces.

FRICTION AND MOTION

Every example of motion, from the one that lets us choose the direction of our gaze to the one that carries a butterfly through the landscape, can be put into one of the two leftmost columns of Table 30. Physically, the two columns are separated by the following criterion: in the first class, the acceleration of a body can be in a different direction from its velocity. The second class of examples produces only accelerations that are exactly *opposed* to the velocity of the moving body, as seen from the frame of reference of the braking medium. Such a resisting force is called *friction*, *drag* or a *damping*. All examples in the second class are types of friction. Just check.

Challenge 369 e

Challenge 368 e

$$\frac{\mathrm{d}q}{\mathrm{d}t} = \int_{V} \frac{\partial \rho}{\partial t} \mathrm{d}V = -\int_{A=\partial V} j \mathrm{d}A = -I \tag{63}$$

or, equivalently,

$$\frac{\partial \rho}{\partial t} + \nabla \mathbf{j} = 0 . {(64)}$$

This is the *continuity equation* for the quantity q. All this only states that a conserved quantity in a closed volume V can only change by flowing through the surface A. This is a typical example of how complex mathematical expressions can obfuscate the simple physical content.

^{*} Mathematically, the conservation of a quantity q is expressed with the help of the volume density $\rho = q/V$, the current I = q/t, and the flow or flux $\mathbf{j} = \rho \mathbf{v}$, so that j = q/At. Conservation then implies

Motion Mountain – The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christoph Schiller November 1997–March 2012

TABLE 30 Selected processes and devices changing the motion of bodies.

'	9 9	
SITUATIONS THAT CAN LEAD TO ACCELERATION	SITUATIONS THAT ONLY LEAD TO DECELERATION	MOTORS AND ACTUATORS
piezoelectricity		
quartz under applied voltage	thermoluminescence	walking piezo tripod
collisions		
satellite in planet encounter	car crash	rocket motor
growth of mountains	meteorite crash	swimming of larvae
magnetic effects		
compass needle near magnet	electromagnetic braking	electromagnetic gun
magnetostriction	transformer losses	linear motor
current in wire near magnet	electric heating	galvanometer
electric effects		
rubbed comb near hair	friction between solids	electrostatic motor
bombs	fire	muscles, sperm flagella
television tube	electron microscope	Brownian motor
light		
levitating objects by light	light bath stopping atoms	(true) light mill
solar sail for satellites	light pressure inside stars	solar cell
elasticity		
bow and arrow	trouser suspenders	ultrasound motor
bent trees standing up again	pillow, air bag	bimorphs
osmosis		
water rising in trees	salt conservation of food	osmotic pendulum
electro-osmosis		tunable X-ray screening
heat & pressure		
freezing champagne bottle	surfboard water resistance	hydraulic engines
tea kettle	quicksand	steam engine
barometer	parachute	air gun, sail
earthquakes	sliding resistance	seismometer
attraction of passing trains	shock absorbers	water turbine
nuclei		
radioactivity	plunging into the Sun	supernova explosion
biology		
bamboo growth	decreasing blood vessel diameter	molecular motors
gravitation		
falling	emission of gravity waves	pulley




FIGURE 145 Shapes and air/water resistance.

Challenge 370 s A puzzle on cycling: does side wind brake – and why?

Friction can be so strong that all motion of a body against its environment is made impossible. This type of friction, called *static friction* or *sticking friction*, is common and important: without it, turning the wheels of bicycles, trains or cars would have no effect. Without static friction, wheels driven by a motor would have no grip. Similarly, not a single screw would stay tightened and no hair clip would work. We could neither run nor walk in a forest, as the soil would be more slippery than polished ice. In fact not only our own motion, but all *voluntary motion* of living beings is *based* on friction. The same is the case for self-moving machines. Without static friction, the propellers in ships, aeroplanes and helicopters would not have any effect and the wings of aeroplanes would produce no lift to keep them in the air. (Why?) In short, static friction is required whenever we or an engine want to move relative to our environment.

Challenge 371 s

FRICTION, SPORT, MACHINES AND PREDICTABILITY

Once an object moves through its environment, it is hindered by another type of friction; it is called *dynamic friction* and acts between bodies in relative motion. Without it, falling bodies would always rebound to the same height, without ever coming to a stop; neither parachutes nor brakes would work; worse, we would have no memory, as we will see later.*

All motion examples in the second column of Table 30 include friction. In these examples, macroscopic energy is not conserved: the systems are *dissipative*. In the first column, macroscopic energy is constant: the systems are *conservative*.

The first two columns can also be distinguished using a more abstract, mathematical criterion: on the left are accelerations that can be derived from a potential, on the right,

Ref. 166

^{*} Recent research suggest that maybe in certain crystalline systems, such as tungsten bodies on silicon, under ideal conditions gliding friction can be extremely small and possibly even vanish in certain directions of motion. This so-called *superlubrication* is presently a topic of research.

decelerations that can not. As in the case of gravitation, the description of any kind of motion is much simplified by the use of a potential: at every position in space, one needs only the single value of the potential to calculate the trajectory of an object, instead of the three values of the acceleration or the force. Moreover, the magnitude of the velocity of an object at any point can be calculated directly from energy conservation.

The processes from the second column *cannot* be described by a potential. These are the cases where we necessarily have to use force if we want to describe the motion of the system. For example, the *friction* or *drag* force *F* due to wind resistance of a body is *roughly* given by

$$F = \frac{1}{2}c_{\rm w}\rho A v^2 \tag{65}$$

where A is the area of its cross-section and v its velocity relative to the air, ρ is the density of air; the *drag coefficient* c_w is a pure number that depends on the shape of the moving object. (A few examples are given in Figure 145. The formula is valid for all fluids, not only for air, below the speed of sound, as long as the drag is due to turbulence. This is usually the case in air and in water. At low velocities, when the fluid motion is not turbulent but laminar, drag is called *viscous* and follows an (almost) linear relation with speed.) You may check that aerodynamic resistance cannot be derived from a potential.*

The drag coefficient $c_{\rm w}$ is a measured quantity.** An aerodynamic car has a value between 0.25 and 0.3; many sports cars share with vans values of 0.44 and higher, and racing car values can be as high as 1, depending on the amount of the force that is used to keep the car fastened to the ground. The lowest known values are for dolphins and penguins.

Wind resistance is also of importance to humans, in particular in athletics. It is estimated that 100 m sprinters spend between 3 % and 6 % of their power overcoming drag. This leads to varying sprint times $t_{\rm w}$ when wind of speed w is involved, related by the expression

$$\frac{t_0}{t_{\rm w}} = 1.03 - 0.03 \left(1 - \frac{wt_{\rm w}}{100 \,\rm m} \right)^2 \,, \tag{66}$$

where the more conservative estimate of 3 % is used. An opposing wind speed of -2 m/s gives an increase in time of 0.13 s, enough to change a potential world record into an 'only' excellent result. (Are you able to deduce the $c_{\rm w}$ value for running humans from the formula?)

Likewise, parachuting exists due to wind resistance. Can you determine how the speed of a falling body, with or without parachute, changes with time, assuming *constant* shape

The topic of aerodynamic shapes is even more interesting for fluid bodies. They are kept together by *surface tension*. For example, surface tension keeps the hairs of a wet brush together. Surface tension also determines the shape of rain drops. Experiments show that it is spherical for drops smaller than 2 mm diameter, and that larger rain drops are *lens* shaped, with the flat part towards the bottom. The usual tear shape is *not* encountered in nature; something vaguely similar to it appears during drop detachment, but *never* during drop fall.

Challenge 372 nv

Ref. 168

Challenge 374 ny

Challenge 373 s

Vol. V, page 232 Ref. 167

^{*} Such a statement about friction is correct only in three dimensions, as is the case in nature; in the case of a single dimension, a potential can *always* be found.

^{**} Calculating drag coefficients in computers, given the shape of the body and the properties of the fluid, is one of the most difficult tasks of science; the problem is still not solved.

Challenge 375 s

and drag coefficient?

Ref. 169

In contrast, static friction has different properties. It is proportional to the force pressing the two bodies together. Why? Studying the situation in more detail, sticking friction is found to be proportional to the actual contact area. It turns out that putting two solids into contact is rather like turning Switzerland upside down and putting it onto Austria; the area of contact is much smaller than that estimated macroscopically. The important point is that the area of actual contact is proportional to the *normal* force. The study of what happens in that contact area is still a topic of research; researchers are investigating the issues using instruments such as atomic force microscopes, lateral force microscopes and triboscopes. These efforts resulted in computer hard discs which last longer, as the friction between disc and the reading head is a central quantity in determining the lifetime.

All forms of friction are accompanied by an increase in the temperature of the moving body. The reason became clear after the discovery of atoms. Friction is not observed in few – e.g. 2, 3, or 4 – particle systems. Friction only appears in systems with many particles, usually millions or more. Such systems are called *dissipative*. Both the temperature changes and friction itself are due to the motion of large numbers of microscopic particles against each other. This motion is not included in the Galilean description. When it is included, friction and energy loss disappear, and potentials can then be used throughout. Positive accelerations – of microscopic magnitude – then also appear, and motion is found to be conserved. As a result, all motion is conservative on a microscopic scale. Therefore, on a microscopic scale it is possible to describe *all* motion without the concept of force.*

The moral of the story is twofold: First, one should use force and power only in one situation: in the case of friction, and only when one does not want to go into the microscopic details.** Secondly, friction is not an obstacle to predictability.

Et qu'avons-nous besoin de ce moteur, quand l'étude réfléchie de la nature nous prouve que le mouvement perpétuel est la première de ses lois ?***

Donatien de Sade *Justine*, ou les malheurs de la vertu.

Ref. 170

^{*} The first scientist who eliminated force from the description of nature was Heinrich Rudolf Hertz (b. 1857 Hamburg, d. 1894 Bonn), the famous discoverer of electromagnetic waves, in his textbook on mechanics, *Die Prinzipien der Mechanik*, Barth, 1894, republished by Wissenschaftliche Buchgesellschaft, 1963. His idea was strongly criticized at that time; only a generation later, when quantum mechanics quietly got rid of the concept for good, did the idea become commonly accepted. (Many have speculated about the role Hertz would have played in the development of quantum mechanics and general relativity, had he not died so young.) In his book, Hertz also formulated the principle of the straightest path: particles follow geodesics. This same description is one of the pillars of general relativity, as we will see later on.

^{**} In the case of human relations the evaluation should be somewhat more discerning, as the research by James Gilligan shows.

^{*** &#}x27;And whatfor do we need this motor, when the reasoned study of nature proves to us that perpetual motion is the first of its laws?'

COMPLETE STATES - INITIAL CONDITIONS

Quid sit futurum cras, fuge quaerere ...*

Horace, *Odi*, lib. I, ode 9, v. 13.

Let us continue our exploration of the predictability of motion. We often describe the motion of a body by specifying the time dependence of its position, for example as

$$\mathbf{x}(t) = \mathbf{x}_0 + \mathbf{v}_0(t - t_0) + \frac{1}{2}\mathbf{a}_0(t - t_0)^2 + \frac{1}{6}\mathbf{j}_0(t - t_0)^3 + \dots$$
 (67)

The quantities with an index 0, such as the starting position x_0 , the starting velocity v_0 , etc., are called *initial conditions*. Initial conditions are necessary for any description of motion. Different physical systems have different initial conditions. Initial conditions thus specify the *individuality* of a given system. Initial conditions also allow us to distinguish the present situation of a system from that at any previous time: initial conditions specify the *changing aspects* of a system. In other words, they summarize the *past* of a system.

Page 26

Initial conditions are thus precisely the properties we have been seeking for a description of the *state* of a system. To find a complete description of states we thus need only a complete description of initial conditions, which we can thus righty call also *initial states*. It turns out that for gravitation, as for all other microscopic interactions, there is *no* need for initial acceleration a_0 , initial jerk j_0 , or higher-order initial quantities. In nature, acceleration and jerk depend only on the properties of objects and their environment; they do not depend on the past. For example, the expression $a = GM/r^2$ of universal gravity, giving the acceleration of a small body near a large one, does not depend on the past, but only on the environment. The same happens for the other fundamental interactions, as we will find out shortly.

Page 82

The *complete state* of a moving mass point is thus described by specifying its position and its momentum at all instants of time. Thus we have now achieved a *complete* description of the *intrinsic properties* of point objects, namely by their mass, and of their *states of motion*, namely by their momentum, energy, position and time. For *extended rigid* objects we also need orientation, angular velocity and angular momentum. Can you specify the necessary quantities in the cases of extended elastic bodies and of fluids? Can you give an example of an intrinsic property that we have so far missed?

Challenge 376 s Challenge 377 s

The set of all possible states of a system is given a special name: it is called the *phase space*. We will use the concept repeatedly. Like any space, it has a number of dimensions. Can you specify it for a system consisting of *N* point particles?

Challenge 378 s

Given that we have a description of both properties and states of point objects, extended rigid objects and deformable bodies, can we predict all motion? Not yet. There are situations in nature where the motion of an object depends on characteristics other than its mass; motion can depend on its colour (can you find an example?), on its temperature, and on a few other properties that we will soon discover. And for each intrinsic property there are state variables to discover. Each property is the basis of a field of physical enquiry. Speed is the basis for mechanics, temperature is the basis fir thermodynam-

Challenge 379 s

Ref. 78 * 'What future will be tomorrow, never ask ...' Horace is Quintus Horatius Flaccus (65–8 BCE), the great Roman poet.

ics, etc. We must therefore conclude that as yet we do not have a complete description of motion.

Challenge 380 s

It is interesting to recall an older challenge and ask again: does the universe have initial conditions? Does it have a phase space? As a hint, recall that when a stone is thrown, the initial conditions summarize the effects of the thrower, his history, the way he got there etc.; in other words, initial conditions summarize the past of a system, i.e., the effects that the environment had during the history of a system.

(An optimist is somebody who thinks that the future is uncertain.

Anonymous

Do surprises exist? Is the future determined?

Die Ereignisse der Zukunft können wir nicht aus den gegenwärtigen erschließen. Der Glaube an den Kausalnexus ist ein Aberglaube.* Ludwig Wittgenstein, Tractatus, 5.1361

Freedom is the recognition of necessity.

Friedrich Engels (1820–1895)

If, after climbing a tree, we jump down, we cannot halt the jump in the middle of the trajectory; once the jump has begun, it is unavoidable and determined, like all passive motion. However, when we begin to move an arm, we can stop or change its motion from a hit to a caress. Voluntary motion does not seem unavoidable or predetermined. Which of these two cases is the general one?

Challenge 381 e

Let us start with the example that we can describe most precisely so far: the fall of a body. Once the gravitational potential φ acting on a particle is given and taken into account, we can use the expression

$$a(x) = -\nabla \varphi = -GMr/r^3 , \qquad (68)$$

and we can use the state at a given time, given by initial conditions such as

$$\mathbf{x}(t_0)$$
 and $\mathbf{v}(t_0)$, (69)

to determine the motion of the particle in advance. Indeed, with these two pieces of information, we can calculate the complete trajectory x(t).

An equation that has the potential to predict the course of events is called an *evolution* equation. Equation (68), for example, is an evolution equation for the fall of the object. (Note that the term 'evolution' has different meanings in physics and in biology.) An evolution equation embraces the observation that not all types of change are observed in nature, but only certain specific cases. Not all imaginable sequences of events are observed, but only a limited number of them. In particular, equation (68) embraces the

^{* &#}x27;We cannot infer the events of the future from those of the present. Belief in the causal nexus is superstition.' Our adventure, however, will show that this statement is wrong.

idea that from one instant to the next, falling objects change their motion based on the gravitational potential acting on them.

Evolution equations do not exist only for motion due to gravity, but for motion due to all forces in nature. Given an evolution equation and initial state, the whole motion of a system is thus *uniquely fixed*, a property of motion often called *determinism*. For example, astronomers can calculate the position of planets with high precision for thousands of years in advance.

Let us carefully distinguish determinism from several similar concepts, to avoid misunderstandings. Motion can be deterministic and at the same time be *unpredictable in practice*. The unpredictability of motion can have four origins:

- 1. an impracticably large number of particles involved,
- 2. the mathematical complexity of the evolution equations,
- 3. insufficient information about initial conditions, and
- 4. strange shapes of space-time.

For example, in case of the weather the first three conditions are fulfilled at the same time. It is hard to predict the weather over periods longer than about a week or two. (In 1942, Hitler made a fool of himself by ordering a precise weather forecast for the following twelve months.) Despite the difficulty of prediction, weather change is still deterministic. As another example, near black holes all four origins apply together. We will discuss black holes in the section on general relativity. Despite being unpredictable, motion is deterministic near black holes.

Motion can be both deterministic and time *random*, i.e., with different outcomes in similar experiments. A roulette ball's motion is deterministic, but it is also random.* As we will see later, quantum-mechanical situations fall into this category, as do all examples of irreversible motion, such as a drop of ink spreading out in clear water. In all such cases the randomness and the irreproducibility are only apparent; they disappear when the description of states and initial conditions in the microscopic domain are included. In short, determinism does not contradict (*macroscopic*) *irreversibility*. However, on the microscopic scale, deterministic motion is always reversible.

A final concept to be distinguished from determinism is *acausality*. Causality is the requirement that a cause must precede the effect. This is trivial in Galilean physics, but becomes of importance in special relativity, where causality implies that the speed of light is a limit for the spreading of effects. Indeed, it seems impossible to have deterministic motion (of matter and energy) which is *acausal*, in other words, faster than light. Can you confirm this? This topic will be looked at more deeply in the section on special relativity.

Saying that motion is 'deterministic' means that it is fixed in the future *and also in the past*. It is sometimes stated that predictions of *future* observations are the crucial test for a successful description of nature. Owing to our often impressive ability to influence the future, this is not necessarily a good test. Any theory must, first of all, describe *past* observations correctly. It is our lack of freedom to change the past that results in our lack

Challenge 382 s

^{*} Mathematicians have developed a large number of tests to determine whether a collection of numbers may be called *random*; roulette results pass all these tests – in honest casinos only, however. Such tests typically check the equal distribution of numbers, of pairs of numbers, of triples of numbers, etc. Other tests are the χ^2 test, the Monte Carlo test(s), and the gorilla test.

of choice in the description of nature that is so central to physics. In this sense, the term 'initial condition' is an unfortunate choice, because in fact, initial conditions summarize the *past* of a system.* The central ingredient of a deterministic description is that all motion can be reduced to an evolution equation plus one specific state. This state can be either initial, intermediate, or final. Deterministic motion is uniquely specified into the past and into the future.

To get a clear concept of determinism, it is useful to remind ourselves why the concept of 'time' is introduced in our description of the world. We introduce time because we observe first that we are able to define sequences in observations, and second, that unrestricted change is impossible. This is in contrast to films, where one person can walk through a door and exit into another continent or another century. In nature we do not observe metamorphoses, such as people changing into toasters or dogs into toothbrushes. We are able to introduce 'time' only because the sequential changes we observe are extremely restricted. If nature were not reproducible, time could not be used. In short, determinism expresses the observation that *sequential changes are restricted to a single possibility*.

Challenge 383 s

Since determinism is connected to the use of the concept of time, new questions arise whenever the concept of time changes, as happens in special relativity, in general relativity and in theoretical high energy physics. There is a lot of fun ahead.

In summary, every description of nature that uses the concept of time, such as that of everyday life, that of classical physics and that of quantum mechanics, is intrinsically and inescapably deterministic, since it connects observations of the past and the future, *eliminating* alternatives. In short, *the use of time implies determinism, and vice versa.* When drawing metaphysical conclusions, as is so popular nowadays when discussing quantum theory, one should never forget this connection. Whoever uses clocks but denies determinism is nurturing a split personality!** The future is determined.

Vol. V, page 35

FREE WILL

You do have the ability to surprise yourself.

Richard Bandler and John Grinder

The idea that motion is determined often produces fear, because we are taught to associate determinism with lack of freedom. On the other hand, we do experience freedom in our actions and call it *free will*. We know that it is necessary for our creativity and for our happiness. Therefore it seems that determinism is opposed to happiness.

But what precisely is free will? Much ink has been consumed trying to find a precise definition. One can try to define free will as the arbitrariness of the choice of initial conditions. However, initial conditions must themselves result from the evolution equations, so that there is in fact no freedom in their choice. One can try to define free will from the idea of unpredictability, or from similar properties, such as uncomputability. But these definitions face the same simple problem: whatever the definition, there is *no way* to

^{*} The problems with the term 'initial conditions' become clear near the big bang: at the big bang, the universe has no past, but it is often said that it has initial conditions. This contradiction will only be resolved in the last part of our adventure.

^{**} That can be a lot of fun though.

prove experimentally that an action was performed freely. The possible definitions are useless. In short, free will *cannot* be observed. (Psychologists also have a lot of their own data to support this, but that is another topic.)

No process that is *gradual* – in contrast to *sudden* – can be due to free will; gradual processes are described by time and are deterministic. In this sense, the question about free will becomes one about the existence of sudden changes in nature. This will be a recurring topic in the rest of this walk. Can nature surprise us? In everyday life, nature does not. Sudden changes are not observed. Of course, we still have to investigate this question in other domains, in the very small and in the very large. Indeed, we will change our opinion several times. The lack of surprises in everyday life is built deep into our body: the concept of curiosity is based on the idea that everything discovered is useful afterwards. If nature continually surprised us, curiosity would make no sense.

Many observations contradict the existence of surprises: in the beginning of our walk we defined time using the continuity of motion; later on we expressed this by saying that time is a consequence of the conservation of energy. Conservation is the opposite of surprise. By the way, a challenge remains: can you show that time would not be definable even if surprises existed only *rarely*?

Challenge 384 s

In summary, so far we have no evidence that surprises exist in nature. Time exists because nature is deterministic. Free will cannot be defined with the precision required by physics. Given that there are no sudden changes, there is only one consistent definition of free will: it is a *feeling*, in particular of independence of others, of independence from fear and of accepting the consequences of one's actions. Free will is a feeling of satisfaction. This solves the apparent paradox; free will, being a feeling, exists as a human experience, even though all objects move without any possibility of choice. There is no contradiction.*

Ref. 172

Ref. 173

Challenge 386 e

Even if human action is determined, it is still authentic. So why is determinism so frightening? That is a question everybody has to ask themselves. What difference does determinism imply for your life, for the actions, the choices, the responsibilities and the pleasures you encounter?** If you conclude that being determined is different from being free, you should change your life! Fear of determinism usually stems from refusal to take the world the way it is. Paradoxically, it is precisely he who insists on the existence of free will who is running away from responsibility.

SUMMARY ON PREDICTABILITY

Despite difficulties to predict specific cases, all motion we encountered so far is deterministic and predictable. The future is determined. In fact, this is the case for all motion in

Challenge 385 e

Challenge 387 s ** If nature's 'laws' are deterministic, are they in contrast with moral or ethical 'laws'? Can people still be held responsible for their actions?

^{*} That free will is a feeling can also be confirmed by careful introspection. The idea of free will always appears *after* an action has been started. It is a beautiful experiment to sit down in a quiet environment, with the intention to make, within an unspecified number of minutes, a small gesture, such as closing a hand. If you carefully observe, in all detail, what happens inside yourself around the very moment of decision, you find either a mechanism that led to the decision, or a diffuse, unclear mist. You never find free will. Such an experiment is a beautiful way to experience deeply the wonders of the self. Experiences of this kind might also be one of the origins of human spirituality, as they show the connection everybody has with the rest of



FIGURE 146 What shape of rail allows the black stone to glide most rapidly from point A to the lower point B?




FIGURE 147 Can motion be described in a manner common to all observers?

nature, even in the domain of quantum theory. If motion were not predictable, we could not have introduced the concept of 'motion' in the first place.

PREDICTABILITY AND GLOBAL DESCRIPTIONS OF MOTION

Πλεῖν ἀνάγκε, ζῆν οὐκ ἀνάγκη.

Pompeius

Physicists aim to talk about motion with the highest precision possible. Predictability is an aspect of precision. The highest predictability – and thus the highest precision – is possible when motion is described as globally as possible.

All over the Earth – even in Australia – people observe that stones fall 'down'. This ancient observation led to the discovery of the universal 'law' of gravity. To find it, all that was necessary was to look for a description of gravity that was valid *globally*. The only additional observation that needs to be recognized in order to deduce the result $a = GM/r^2$ is the variation of gravity with height.

In short, thinking *globally* helps us to make our description of motion more precise and our predictions more useful. How can we describe motion as globally as possible? It turns out that there are six approaches to this question, each of which will be helpful on our way to the top of Motion Mountain. We first give an overview, and then explore the details of each approach.

1. Action principles or Variational principles, the first global approach to motion, arise when we overcome a limitation of what we have learned so far. When we predict the motion of a particle from its current acceleration with an evolution equation, we are using the most *local* description of motion possible. We use the acceleration of a particle at a certain place and time to determine its position and motion *just after* that moment and *in the immediate neighbourhood* of that place. Evolution equations thus have a mental 'horizon' of radius zero.

The contrast to evolution equations are variational principles. A famous example is illustrated in Figure 146. The challenge is to find the path that allows the fastest

^{*} Navigare necesse, vivere non necesse. 'To navigate is necessary, to live is not.' Gnaeus Pompeius Magnus (106–48 BCE) is cited in this way by Plutarchus (*c.* 45 to *c.* 125).



Page 204




FIGURE 148 What happens when one rope is cut?




FIGURE 149 A famous mechanism, the Peaucellier-Lipkin linkage, that allows drawing a straight line with a compass: fix point F, put a pencil into joint P and move C with a compass along a circle.

Challenge 388 d

possible gliding motion from a high point to a distant low point. The sought path is the brachistochrone, from ancient Greek for 'shortest time', This puzzle asks about a property of motion as a whole, for all times and positions. The global approach required by questions such as this one will lead us to a description of motion which is simple, precise and fascinating: the so-called principle of cosmic laziness, also known as the principle of least action.

- 2. Relativity, the second global approach to motion, emerges when we compare the various descriptions of the same system produced by different observers. For example, the observations by somebody falling from a cliff – as shown in Figure 147 – a passenger in a roller coaster, and an observer on the ground will usually differ. The relationships between these observations, the so-called symmetry transformations, lead us to a global description, valid for everybody. Later, this approach will lead us to Einstein's special and general theory of relativity.
- 3. Mechanics of extended and rigid bodies, rather than mass points, is required to underrotates around the rope.

stand many objects, plants and animals. As an example, the counter-intuitive result of the experiment in Figure 148 shows why this topic is worthwhile. The rapidly rotating wheel suspended on only one end of the axis remains almost horizontal, but slowly

In order to design machines, it is essential to understand how a group of rigid bodies interact with one another. For example, take the Peaucellier-Lipkin linkage shown in Figure 149. A joint F is fixed on a wall. Two movable rods lead to two opposite corners of a movable rhombus, whose rods connect to the other two corners C and P. This mechanism has several astonishing properties. First of all, it implicitly defines a circle of radius R so that one always has the relation $r_{\rm C} = R^2/r_{\rm P}$ between the distances of joints C and P from the centre of this circle. Can you find this special circle? Secondly, if you put a pencil in joint P, and let joint C follow a certain circle, the pencil P draws a straight line. Can you find that circle? The mechanism thus allows drawing a *straight* line with the help of a compass.

Another famous machine challenge is to devise a wooden carriage, with gearwheels that connect the wheels to an arrow, with the property that, whatever path

Challenge 389 ny

Ref. 175

Challenge 390 ny Challenge 391 ny

Ref. 176






FIGURE 150 A south-pointing carriage: whatever the path it follows, the arrow on it always points south.

Challenge 392 d

Vol. II, page 190

Ref. 177

Ref. 178 Challenge 393 s

Challenge 395 s

the carriage takes, the arrow always points south (see Figure 150). The solution to this puzzle will even be useful in helping us to understand general relativity, as we will see. Such a wagon allows measuring the curvature of a surface and of space.

Another interesting example of rigid motion is the way that human movements, such as the general motions of an arm, are composed from a small number of basic motions. All these examples are from the fascinating field of engineering; unfortunately, we will have little time to explore this topic in our hike.

4. The next global approach to motion is the description of *non-rigid extended bodies*. For example, *fluid mechanics* studies the flow of fluids (like honey, water or air) around solid bodies (like spoons, ships, sails or wings). Fluid mechanics thus describes how insects, birds and aeroplanes fly,* why sailing-boats can sail against the wind, what happens when a hard-boiled egg is made to spin on a thin layer of water, or how a bottle full of wine can be emptied in the fastest way possible.

As well as fluids, we can study the behaviour of deformable *solids*. This area of research is called *continuum mechanics*. It deals with deformations and oscillations of extended structures. It seeks to explain, for example, why bells are made in particular shapes; how large bodies – such as the falling chimneys shown in Figure 151 – or small bodies – such as diamonds – break when under stress; and how cats can turn themselves the right way up as they fall. During the course of our journey we will repeatedly encounter issues from this field, which impinges even upon general relativity and the world of elementary particles.

5. *Statistical mechanics* is the study of the motion of huge numbers of particles. Statistical mechanics is yet another global approach to the study of motion. The concepts needed to describe gases, such as temperature, entropy and pressure (see Figure 152),



^{*} The mechanisms of insect flight are still a subject of active research. Traditionally, fluid dynamics has concentrated on large systems, like boats, ships and aeroplanes. Indeed, the smallest human-made object that can fly in a controlled way – say, a radio-controlled plane or helicopter – is much larger and heavier than many flying objects that evolution has engineered. It turns out that controlling the flight of small things requires more knowledge and more tricks than controlling the flight of large things. There is more about this topic on page 206 in Volume V.

FIGURE 151 How and where does a falling Challenge 394 ny brick chimney break?




FIGURE 152 Why do hot-air balloons stay inflated? How can you measure the weight of a bicycle rider using only a ruler?







FIGURE 153 Why do marguerites – or ox-eye daisies, *Leucanthemum vulgare* – usually have around 21 (left and centre) or around 34 (right) petals? (© Anonymous, Giorgio Di Iorio and Thomas Lüthi)

are essential tools of this discipline. They help us to understand why some processes in nature do not occur backwards. These concepts will also help us take our first steps towards the understanding of black holes.

6. The last global approach to motion, *self-organization*, involves all of the above-mentioned viewpoints *at the same time*. Such an approach is needed to understand everyday experience, and *life* itself. Why does a flower form a specific number of petals, as shown in Figure 153? How does an embryo differentiate in the womb? What makes our hearts beat? How do mountains ridges and cloud patterns emerge? How do stars and galaxies evolve? How are sea waves formed by the wind?

All these are examples of *self-organization* processes; life scientists simply speak of *growth* processes. Whatever we call them, all these processes are characterized by the spontaneous appearance of patterns, shapes and cycles. Such processes are a common research theme across many disciplines, including biology, chemistry, medicine, geology and engineering.

We will now explore to these six global approaches to motion. We will begin with the first approach, namely, the global description of motion using a variational principle. This beautiful method was the result of several centuries of collective effort, and is the highlight of particle dynamics. Variational principles also provide the basis for all the other global approaches just mentioned and for all the further descriptions of motion that we will explore afterwards.

CHAPTER 8

MEASURING CHANGE WITH ACTION

OTION can be described by numbers. Take a single particle that oves. The expression (x(t), y(t), z(t)) describes how, during its otion, position changes with time. This description is completed by stating how its speed $(v_x(t), v_y(t), v_z(t))$ changes over time. Indeed, realizing that these two expressions can be used to describe the path and the behaviour of a moving particle was a milestone in the development of modern physics.

The next milestone of modern physics is achieved by answering a short but hard question. If motion is a type of *change*, as the Greek already said, how can we *measure* change? In fact, change can be measured by a single number. Physicists took almost two centuries of attempts to uncover the way to measure change. As a result, the quantity that measures change has a strange name: it is called *(physical) action.** To remember the connection of 'action' with change, just think about a Hollywood film: a lot of action means a large amount of change.

Introducing physical action as a measure of change is important, because it provides the first and the most useful global description of motion. In fact, we are now ready to define action

Imagine taking two snapshots of a system at different times. How could you define the amount of change that occurred in between? When do things change a lot, and when do they change only a little? First of all, a system with many moving parts shows a lot of change. So it makes sense that the action of a system composed of independent subsystems should be the *sum* of the actions of these subsystems.

Secondly, systems with *high energy*, such as the explosions shown in Figure 155, show larger change than systems at lower speed. Indeed, we introduced energy as the quantity

Behind these shifts in terminology is the story of an intense two-centuries-long attempt to describe motion with so-called *extremal* or *variational principles*: the objective was to complete and improve the work initiated by Leibniz. These principles are only of historical interest today, because all are special cases of the principle of least action described here.

Page 20

^{*} Note that this 'action' is not the same as the 'action' appearing in statements such as 'every action has an equal and opposite reaction'. This last usage, coined by Newton for certain forces, has not stuck; therefore the term has been recycled. After Newton, the term 'action' was first used with an intermediate meaning, before it was finally given the modern meaning used here. This modern meaning is the only meaning used in this text.

Another term that has been recycled is the 'principle of least action'. In old books it used to have a different meaning from the one in this chapter. Nowadays, it refers to what used to be called *Hamilton's principle* in the Anglo-Saxon world, even though it is (mostly) due to others, especially Leibniz. The old names and meanings are falling into disuse and are not continued here.

FIGURE 154 Giuseppe Lagrangia/Joseph Lagrange (1736–1813).



FIGURE 155 Physical action measures change: an example of process with large action value (© Christophe Blanc).

Page 102 that measures how much a system changes over time.

Thirdly, change often – but not always – *builds up over time*; in other cases, recent change can compensate for previous change, as in a pendulum. Change can thus increase or decrease with time.

Finally, for a system in which motion is *stored*, transformed or shifted from one subsystem to another, especially when kinetic energy is stored or changed to potential energy, change is smaller than for a system where all systems move freely.

The mentioned properties imply that the natural measure of change is the average difference between kinetic and potential energy multiplied by the elapsed time. This quantity has all the right properties: it is the sum of the corresponding quantities for all sub-



TABLE 31 Some action values for changes and processes either observed or imagined.

SYSTEM AND PROCESS	APPROXIMATE ACTION VALUE
Smallest measurable action	$1.1 \cdot 10^{-34} \text{Js}$
Light	
Smallest blackening of photographic film	$< 10^{-33} \text{Js}$
Photographic flash	$c. 10^{-17} \mathrm{Js}$
Electricity	,
Electron ejected from atom or molecule	$c. 10^{-33} \text{ Js}$
Current flow in lightning bolt	c. 10 ⁴ Js
Mechanics and materials	,
Tearing apart two neighbouring iron atoms	$c. 10^{-33} \mathrm{Js}$
Breaking a steel bar	c. 10 ¹ Js
Tree bent by the wind from one side to the other	c. 500 Js
Making a white rabbit vanish by 'real' magic	c. 100 PJs
Hiding a white rabbit	c. 0.1 Js
Car crash	c. 2 kJs
Driving car stops within the blink of an eye	c. 20 kJs
Levitating yourself within a minute by 1 m	c. 40 kJs
Large earthquake	c. 1 PJs
Driving car disappears within the blink of an eye	c. 1 ZJs
Sunrise	c. 0.1 ZJs
Chemistry	0.0.12)0
Atom collision in liquid at room temperature	$c. 10^{-33} \text{Js}$
Smelling one molecule	$c. 10^{-31} \text{ Js}$
Burning fuel in a cylinder in an average car engine explosion	c. 10 ⁴ Js
Held versus dropped glass	c. 0.8 Js
Life	2. 0.0)0
Air molecule hitting eardrum	$c. 10^{-32} \mathrm{Js}$
Ovule fertilization	c. 10^{-20} Js
Cell division	$c. 10^{-15} \text{Js}$
Fruit fly's wing beat	$c. 10^{-10} \text{Js}$
Flower opening in the morning	c. 1 nJs
Getting a red face	c. 10 mJs
Maximum brain change in a minute	c. 5 Js
Person walking one body length	$c. 10^2 \text{ Js}$
Birth	c. 2 kJs
Change due to a human life	c. 1 EJs
Nuclei and stars	J. 1 11/0
Single nuclear fusion reaction in star	$c. 10^{-15} \mathrm{Js}$
Explosion of gamma-ray burster	c. 10 ⁴⁶ Js
Universe after one second has elapsed	undefined and undefi

Challenge 396 e




FIGURE 156 Defining a total change or action as an accumulation (addition, or integral) of small changes or actions over time.

systems if these are independent; it generally increases with time (unless the evolution compensates for something that happened earlier); and it decreases if the system transforms motion into potential energy.

Thus the (physical) action S, measuring the change in a system, is defined as

$$S = \overline{L} \cdot \Delta t = \overline{T - U} \cdot (t_{\rm f} - t_{\rm i}) = \int_{t_{\rm i}}^{t_{\rm f}} (T - U) \, \mathrm{d}t = \int_{t_{\rm i}}^{t_{\rm f}} L \, \mathrm{d}t \,, \tag{70}$$

where T is the kinetic energy, U the potential energy we already know, L is the difference between these, and the overbar indicates a time average. The quantity L is called the Lagrangian (function) of the system,* describes what is being added over time, whenever things change. The sign \int is a stretched 'S', for 'sum', and is pronounced 'integral of'. In intuitive terms it designates the operation (called integration) of adding up the values of a varying quantity in infinitesimal time steps dt. The initial and the final times are written below and above the integration sign, respectively. Figure 156 illustrates the idea: the integral is simply the size of the dark area below the curve L(t).

Challenge 397 e Mathematically, the integral of the curve L(t) is defined as

$$\int_{t_{i}}^{t_{f}} L(t) dt = \lim_{\Delta t \to 0} \sum_{m=i}^{f} L(t_{m}) \Delta t = \overline{L} \cdot (t_{f} - t_{i}).$$
 (71)

In other words, the *integral* is the limit, as the time slices get smaller, of the sum of the areas of the individual rectangular strips that approximate the function. Since the \sum sign



^{*} It is named after Giuseppe Lodovico Lagrangia (b. 1736 Torino, d. 1813 Paris), better known as Joseph Louis Lagrange. He was the most important mathematician of his time; he started his career in Turin, then worked for 20 years in Berlin, and finally for 26 years in Paris. Among other things he worked on number theory and analytical mechanics, where he developed most of the mathematical tools used nowadays for calculations in classical mechanics and classical gravitation. He applied them successfully to many motions in the solar system.

Motion Mountain – The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christoph Schiller November 1997–Ma

also means a sum, and since an infinitesimal Δt is written $\mathrm{d}t$, we can understand the notation used for integration. Integration is a sum over slices. The notation was developed by Gottfried Wilhelm Leibniz to make exactly this point. Physically speaking, the integral of the Lagrangian measures the total *effect* that L builds up over time. Indeed, action is called 'effect' in some languages, such as German. The effect that builds up is the total change in the system.

In short, the integral of the Lagrangian, the action, measures the total change that occurs in a system. Physical action is total change. Action, or change, is the integral of the Lagrangian over time. The unit of action, and thus of change, is the unit of energy (the Joule) times the unit of time (the second).

▷ Change is measured in Js.

A large value means a big change. Table 31 shows some values of action observed in nature.

To understand the definition of action in more detail, we will start with the simplest case: a system for which the potential energy is zero, such as a particle moving freely. Obviously, the higher the kinetic energy is, the more change there is. Also, if we observe the particle at two instants, the more distant they are the larger the change. This is not surprising.

Next, we explore a single particle moving in a potential. For example, a falling stone loses potential energy in exchange for a gain in kinetic energy. The more kinetic energy is stored into potential energy, the less change there is. Hence the minus sign in the definition of L. If we explore a particle that is first thrown up in the air and then falls, the curve for L(t) first is below the times axis, then above. We note that the definition of integration makes us count the grey surface *below* the time axis *negatively*. Change can thus be negative, and be compensated by subsequent change, as expected.

To measure change for a system made of several independent components, we simply add all the kinetic energies and subtract all the potential energies. This technique allows us to define actions for gases, liquids and solid matter. Even if the components interact, we still get a sensible result. In short, action is an *additive* quantity.

Physical action thus measures, in a single number, the change observed in a system between two instants of time. Physical action quantifies the change of a physical process. This is valid for *all* observations, i.e., for all systems and for all processes: an explosion, a caress or a colour change. Change is measured in Js. We will discover later that describing change with a single number is also possible in relativity and quantum theory. Any change going on in any system of nature, be it transport, transformation or growth, can be measured with a single number.

THE PRINCIPLE OF LEAST ACTION

We now have a precise measure of change, which, as it turns out, allows a simple, global and powerful description of motion. In nature, the change happening between two instants is always the *smallest* possible. *In nature, action is minimal.** Of all possible mo-





^{*} In fact, in some macroscopic situations the action can be a saddle point, so that the snobbish form of the principle is that the action is 'stationary'. In contrast to what is often heard, the action is *never* a maximum.



FIGURE 157 The minimum of a curve has vanishing slope.

tions, nature always chooses for which the change is *minimal*. Let us study a few examples.

In the simple case of a free particle, when no potentials are involved, the principle of minimal action implies that the particle moves in a *straight* line with *constant* velocity. All other paths would lead to larger actions. Can you verify this?

When gravity is present, a thrown stone flies along a parabola (or more precisely, along an ellipse) because any other path, say one in which the stone makes a loop in the air, would imply a *larger* action. Again you might want to verify this for yourself.

All observations support this simple and basic statement: things always move in a way that produces the smallest possible value for the action. This statement applies to the full path and to any of its segments. Bertrand Russell called it the 'law of cosmic laziness'.

It is customary to express the idea of minimal change in a different way. The action varies when the path is varied. The actual path is the one with the smallest action. You will recall from school that at a minimum the derivative of a quantity vanishes: a minimum has a horizontal slope. In the present case, we do not vary a quantity, but a complete path; hence we do not speak of a derivative or slope, but of a variation. It is customary to write the variation of action as δS . The *principle of least action* thus states:

$$\triangleright$$
 The actual trajectory between specified end points satisfies $\delta S = 0$. (72)

Mathematicians call this a *variational principle*. Note that the end points have to be specified: we have to compare motions with the *same* initial and final situations.

Before discussing the principle further, we can check that it is equivalent to the evolution equation.* To do this, we can use a standard procedure, part of the so-called *calculus*

$$\delta W = \sum_{i} m_i \ddot{x}_i \delta x_i \,. \tag{73}$$

Motion is then described in the following way:

$$The actual trajectory satisfies \int_{t_i}^{t_f} (\delta T + \delta W) dt = 0 \quad provided \quad \delta x(t_i) = \delta x(t_f) = 0 \ .$$
 (74)

Challenge 398 e

Challenge 399 e

Page 200



Moreover, for motion on small (infinitesimal) scales, the action is always a minimum. The mathematical condition of vanishing variation, given below, encompasses all these details.

^{*} For those interested, here are a few comments on the equivalence of Lagrangians and evolution equations. First of all, Lagrangians do not exist for non-conservative, or *dissipative* systems. We saw that there is no potential for any motion involving *friction* (and more than one dimension); therefore there is no action in these cases. One approach to overcome this limitation is to use a generalized formulation of the principle of least action. Whenever there is no potential, we can express the *work* variation δW between different trajectories x_i as

Challenge 401 nv

of variations. The condition $\delta S = 0$ implies that the action, i.e., the area under the curve in Figure 156, is a minimum. A little bit of thinking shows that if the Lagrangian is of the form $L(x_n, v_n) = T(v_n) - U(x_n)$, then the minimum area is achieved when

$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\frac{\partial T}{\partial v_{\mathrm{n}}} \right) = -\frac{\partial U}{\partial x_{\mathrm{n}}} \tag{75}$$

where n counts all coordinates of all particles.* For a single particle, these Lagrange's equations of motion reduce to

$$m\mathbf{a} = -\nabla U \ . \tag{77}$$

Challenge 400 ny

Challenge 402 e

The quantity being varied has no name; it represents a generalized notion of change. You might want to check that it leads to the correct evolution equations. Thus, although *proper* Lagrangian descriptions exist only for *conservative* systems, for dissipative systems the principle can be generalized and remains useful.

Many physicists will prefer another approach. What a mathematician calls a generalization is a special case for a physicist: the principle (74) hides the fact that *all* friction results from the usual principle of minimal action, if we include the complete microscopic details. There is no friction in the microscopic domain. Friction is an approximate, macroscopic concept.

Nevertheless, more mathematical viewpoints are useful. For example, they lead to interesting limitations for the use of Lagrangians. These limitations, which apply only if the world is viewed as purely classical – which it isn't – were discovered about a hundred years ago. In those times, computers were not available, and the exploration of new calculation techniques was important. Here is a summary.

The coordinates used in connection with Lagrangians are not necessarily the Cartesian ones. *Generalized* coordinates are especially useful when there are *constraints* on the motion. This is the case for a pendulum, where the weight always has to be at the same distance from the suspension, or for an ice skater, where the skate has to move in the direction in which it is pointing. Generalized coordinates may even be mixtures of positions and momenta. They can be divided into a few general types.

Ref. 181

Generalized coordinates are called *holonomic-scleronomic* if they are related to Cartesian coordinates in a fixed way, independently of time: physical systems described by such coordinates include the pendulum and a particle in a potential. Coordinates are called *holonomic-rheonomic* if the dependence involves time. An example of a rheonomic systems would be a pendulum whose length depends on time. The two terms rheonomic and scleronomic are due to Ludwig Boltzmann. These two cases, which concern systems that are only described by their geometry, are grouped together as *holonomic systems*. The term is due to Heinrich Hertz.

Page 327

Vol. III, page 90

The more general situation is called *anholonomic*, or *nonholonomic*. Lagrangians work well only for holonomic systems. Unfortunately, the meaning of the term 'nonholonomic' has changed. Nowadays, the term is also used for certain rheonomic systems. The modern use calls nonholonomic any system which involves velocities. Therefore, an ice skater or a rolling disc is often called a nonholonomic system. Care is thus necessary to decide what is meant by nonholonomic in any particular context.

Even though the use of Lagrangians, and of action, has its limitations, these need not bother us at microscopic level, since microscopic systems are always conservative, holonomic and scleronomic. At the fundamental level, evolution equations and Lagrangians are indeed equivalent.

* The most general form for a Lagrangian $L(q_n,\dot{q}_n,t)$, using generalized holonomic coordinates q_n , leads to Lagrange equations of the form

$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\frac{\partial L}{\partial \dot{q}_{\mathrm{n}}} \right) = \frac{\partial L}{\partial q_{\mathrm{n}}} \ . \tag{76}$$

In order to deduce these equations, we also need the relation $\delta \dot{q} = d/dt(\delta q)$. This relation is valid only for *holonomic* coordinates introduced in the previous footnote and explains their importance.

It should also be noted that the Lagrangian for a moving system is not unique; however, the study of how Ref. 182 the various Lagrangians for a given moving system are related is not part of this walk.

By the way, the letter q for position and p for momentum were introduced in physics by the mathematician Carl Jacobi (b. 1804 Potsdam, d. 1851 Berlin).

·• {

ion Mountain – The Adventure of Physics | pdf file available free of charge at www.motionmountain.net | Copyright © Christoph Schiller November 1

This is the evolution equation: it says that the force acting on a particle is the gradient of the potential energy U. The principle of least action thus implies the equation of motion. (Can you show the converse, which is also correct?)

Challenge 403 s

In other words, *all systems evolve in such a way that the change or action is as small as possible.* Nature is economical. Nature is thus the opposite of a Hollywood thriller, in which the action is maximized; nature is more like a wise old man who keeps his actions to a minimum.

The principle of minimal action states that the actual trajectory is the one for which the *average* of the Lagrangian over the whole trajectory is minimal (see Figure 156). Nature is a Dr. Dolittle. Can you verify this? This viewpoint allows one to deduce Lagrange's equations (75) directly.

Challenge 404 ny

The principle of least action distinguishes the actual trajectory from all other imaginable ones. This observation lead Leibniz to his famous interpretation that the actual world is the 'best of all possible worlds.'* We may dismiss this as metaphysical speculation, but we should still be able to feel the fascination of the issue. Leibniz was so excited about the principle of least action because it was the first time that actual observations were distinguished from all other imaginable possibilities. For the first time, the search for reasons why things are the way they are became a part of physical investigation. Could the world be different from what it is? In the principle of least action, we have a hint of a negative answer. (What do you think?) The final answer will emerge only in the last part of our adventure.

Challenge 405 s

LAGRANGIANS AND MOTION

Ref. 183

Never confuse movement with action.

Ernest Hemingway

Systems evolve by minimizing change. Change, or action, is the time integral of the Lagrangian. As a way to describe motion, the Lagrangian has several advantages over the evolution equation. First of all, the Lagrangian is usually more *compact* than writing the corresponding evolution equations. For example, only *one* Lagrangian is needed for one system, however many particles it includes. One makes fewer mistakes, especially sign mistakes, as one rapidly learns when performing calculations. Just try to write down the evolution equations for a chain of masses connected by springs; then compare the effort with a derivation using a Lagrangian. (The system is often studied because it behaves like a chain of atoms.) We will encounter another example shortly: David Hilbert took only a few weeks to deduce the equations of motion of general relativity using a Lagrangian, whereas Albert Einstein had worked for ten years searching for them directly.

Challenge 406 ny

In addition, the description with a Lagrangian is valid with *any* set of coordinates describing the objects of investigation. The coordinates do not have to be Cartesian; they can be chosen as one prefers: cylindrical, spherical, hyperbolic, etc. These so-called *generalized coordinates* allow one to rapidly calculate the behaviour of many mechanical systems that are in practice too complicated to be described with Cartesian coordinates.



^{*} This idea was ridiculed by the French philosopher Voltaire (1694–1778) in his lucid writings, notably in the brilliant book *Candide*, written in 1759, and still widely available.

TABLE 32 Some Lagrangians.

System	LAGRANGIAN	QUANTITIES
Free, non-relativistic mass point	$L = \frac{1}{2}mv^2$	mass m, speed v = dx/dt
Particle in potential	$L = \frac{1}{2}mv^2 - m\varphi(x)$	gravitational potential ϕ
Mass on spring	$L = \frac{1}{2}mv^2 - \frac{1}{2}kx^2$	elongation <i>x</i> , spring constant <i>k</i>
Mass on frictionless table attached to spring	$L = \frac{1}{2}mv^2 - k(x^2 + y^2)$	spring constant k , coordinates x , y
Chain of masses and springs (simple model of atoms in a linear crystal)	$L = \frac{1}{2}m\sum v_{i}^{2} - \frac{1}{2}m\omega^{2}\sum_{i,j}(x_{i} - x_{j})^{2}$	coordinates x_i , lattice frequency ω
Free, relativistic mass point	$L = -mc^2 \sqrt{1 - v^2/c^2}$	mass m , speed v , speed of light c

For example, for programming the motion of robot arms, the angles of the joints provide a clearer description than Cartesian coordinates of the ends of the arms. Angles are non-Cartesian coordinates. They simplify calculations considerably: the task of finding the most economical way to move the hand of a robot from one point to another can be solved much more easily with angular variables.

More importantly, the Lagrangian allows one to quickly deduce the essential properties of a system, namely, its *symmetries* and its *conserved quantities*. We will develop this important idea shortly, and use it regularly throughout our walk.

Finally, the Lagrangian formulation can be generalized to encompass *all types of interactions*. Since the concepts of kinetic and potential energy are general, the principle of least action can be used in electricity, magnetism and optics as well as mechanics. The principle of least action is central to general relativity and to quantum theory, and allows one to easily relate both fields to classical mechanics.

As the principle of least action became well known, people applied it to an ever-increasing number of problems. Today, Lagrangians are used in everything from the study of elementary particle collisions to the programming of robot motion in artificial intelligence. (Table 32 shows a few examples.) However, we should not forget that despite its remarkable simplicity and usefulness, the Lagrangian formulation is *equivalent* to the evolution equations. It is neither more general nor more specific. In particular, it is *not an explanation* for any type of motion, but only a different view of it. In fact, the search for a new physical 'law' of motion is *just* the search for a new Lagrangian. This makes sense, as the description of nature always requires the description of change. Change in nature is always described by actions and Lagrangians.

The principle of least action states that the action is minimal when the end points of the motion, and in particular the time between them, are fixed. It is less well known that the reciprocal principle also holds: if the action value is kept fixed, the elapsed time for the actual motion is *maximal*. Can you show this?

Even though the principle of least action is not an explanation of motion, the principle somehow calls for such an explanation. We need some patience, though. *Why* nature fol-

Page 238

Ref. 179

Challenge 407 s

Ref. 184

Challenge 408 ny

lows the principle of least action, and *how* it does so, will become clear when we explore quantum theory.

WHY IS MOTION SO OFTEN BOUNDED?

The optimist thinks this is the best of all possible worlds, and the pessimist knows it.

Robert Oppenheimer

Looking around ourselves on Earth or in the sky, we find that matter is not evenly distributed. Matter tends to be near other matter: it is lumped together in *aggregates*. Some major examples of aggregates are listed in Figure 158 and Table 33. All aggregates have mass and size. In the mass–size diagram of Figure 158, both scales are logarithmic. One notes three straight lines: a line $m \sim l$ extending from the Planck mass* upwards, via black holes, to the universe itself; a line $m \sim 1/l$ extending from the Planck mass downwards, to the lightest possible aggregate; and the usual matter line with $m \sim l^3$, extending from atoms upwards, via everyday objects, the Earth to the Sun. The first of the lines, the black hole limit, is explained by general relativity; the last two, the aggregate limit and the common matter line, by quantum theory.**

The aggregates outside the common matter line also show that the stronger the interaction that keeps the components together, the smaller the aggregate. But why is matter mainly found in lumps?

First of all, aggregates form because of the existence of *attractive* interactions between objects. Secondly, they form because of *friction*: when two components approach, an aggregate can only be formed if the released energy can be changed into heat. Thirdly, aggregates have a finite size because of *repulsive* effects that prevent the components from collapsing completely. Together, these three factors ensure that bound motion is much more common than unbound, 'free' motion.

Only three types of attraction lead to aggregates: gravity, the attraction of electric charges, and the strong nuclear interaction. Similarly, only three types of repulsion are observed: rotation, pressure, and the Pauli exclusion principle (which we will encounter later on). Of the nine possible combinations of attraction and repulsion, not all appear in nature. Can you find out which ones are missing from Figure 158 and Table 33, and why?

Together, attraction, friction and repulsion imply that change and action are minimized when objects come and stay together. The principle of least action thus implies the stability of aggregates. By the way, formation history also explains why so many aggregates *rotate*. Can you tell why?

But why does friction exist at all? And why do attractive and repulsive interactions exist? And why is it – as it would appear from the above – that in some distant past matter was *not* found in lumps? In order to answer these questions, we must first study another global property of motion: symmetry.

Ref. 185

Vol. IV, page 117

Challenge 409 s

Challenge 410 ny



^{*} The Planck mass is given by $m_{\rm Pl} = \sqrt{\hbar c/G} = 21.767(16) \,\mu {\rm g}.$

^{**} Figure 158 suggests that domains beyond physics exist; we will discover later on that this is not the case, as mass and size are not definable in those domains.



universe mass [kg] galaxy 10⁴⁰ black holes star cluster Beyond Takute and science seyond nature and science limit Beyond nature and science: beyond Planck length limit Earth neutron 10²⁰ Beyond nature and science: undefined common matterline mountain human 10⁰ Planck mass o cell **DNA** heavy 10⁻²⁰ nucleus 0 uranium muon proton hydrogen electron thictoscopic aggregate linit **Aggregates** 10⁻⁴⁰ neutrino **Elementary** particles lightest imaginable 10⁻⁶⁰ aggregate 10⁻⁴⁰ 10⁻²⁰ 10²⁰ 10⁰ size [m]

FIGURE 158 Elementary particles and aggregates found in nature.

TABLE 33 Some major aggregates observed in nature.

AGGREGATE	SIZE	Овѕ.	Constituents
	(D I A M E T E R)	NUM.	
Gravitationally bound ag	gregates		
Matter across universe	c. 100 Ym	1	superclusters of galaxies, hydrogen and helium atoms
Quasar	10 ¹² to 10 ¹⁴ m	$20 \cdot 10^6$	baryons and leptons
Supercluster of galaxies	c. 3 Ym	10^{7}	galaxy groups and clusters
Galaxy cluster	c. 60 Zm	$25 \cdot 10^9$	10 to 50 galaxies
Galaxy group or cluster	c. 240 Zm		50 to over 2000 galaxies
Our local galaxy group	50 Zm	1	c. 40 galaxies



AGGREGATE	SIZE	Овѕ.	Constituents
	(D I A M E T E R)	NUM.	
General galaxy	0.5 to 2 Zm	3.5 · 10 ¹²	2 10^{10} to $3 \cdot 10^{11}$ stars, dust and gas clouds, probably solar systems
Our galaxy	1.0(0.1) Zm	1	10 ¹¹ stars, dust and gas clouds, solar systems
Interstellar clouds	up to 15 Em	$\gg 10^5$	hydrogen, ice and dust
Solar system ^a	unknown	> 400	star, planets
Our solar system	30 Pm	1	Sun, planets (Pluto's orbit's diameter 11.8 Tm), moons, planetoids, comets asteroids, dust, gas
Oort cloud	6 to 30 Pm	1	comets, dust
Kuiper belt	60 Tm	1	planetoids, comets, dust
Star ^b	10 km to 100 Gm	10 ^{22±1}	ionized gas: protons, neutrons, electrons, neutrinos, photons
Our star, the Sun	1.39 Gm		
Planet ^a (Jupiter, Earth)	143 Mm, 12.8 Mm	8+ > 400	solids, liquids, gases; in particular, heavy atoms
Planetoids (Varuna, etc)	50 to 1 000 km	> 100 (est. 10^9)	solids)
Moons	10 to 1 000 km	> 50	solids
neutron stars	10 km	> 1000	mainly neutrons
Electromagnetically bound	d aggregates ^c		
Dwarf planets, minor planets, asteroids ^d	1 m to 2400 km	> 10 ⁶	(10 ⁹ estimated) solids, usually monolithic
Comets	10 cm to 50 km	> 109	(10 ¹² possible) ice and dust
Mountains, solids, liquids, gases, cheese	1 nm to > 100 km	n.a.	molecules, atoms
Animals, plants, kefir	5 μm to 1 km	$10^{26\pm2}$	organs, cells
brain, human	0.2 m	10^{10}	neurons and other cell types
Cells:		$10^{31\pm1}$	organelles, membranes, molecules
smallest (Nanoarchaeum equitans)	c. 400 nm		molecules
amoeba	c. 600 µm		molecules
largest (whale nerve, single-celled plants)	c. 30 m		molecules
Molecules:		10 ^{78±2}	atoms
H_2	c. 50 pm	$10^{72\pm2}$	atoms
DNA (human)	2 m (total per cell)	10^{21}	atoms
Atoms, ions	30 pm to 300 pm	$10^{80\pm2}$	electrons and nuclei

Aggregates bound by the weak interaction c None



AGGREGATE	SIZE	Овѕ.	Constituents
	(D I A M E T E R)	NUM.	
Aggregates bound by the st	trong interaction ^c		
Nucleus	0.9 to > 7 fm	$10^{79\pm2}$	nucleons
Nucleon (proton, neutron)	0.9 fm	$10^{80\pm2}$	quarks
Mesons	c. 1 fm	n.a.	quarks
Neutron stars: see above			

Ref. 186

a. Only in 1994 was the first evidence found for objects circling stars other than our Sun; of over 400 extrasolar planets found so far, most are found around F, G and K stars, including neutron stars. For example, three objects circle the pulsar PSR 1257+12, and a matter ring circles the star β Pictoris. The objects seem to be dark stars, brown dwarfs or large gas planets like Jupiter. Due to the limitations of observation systems, none of the systems found so far form solar systems of the type we live in. In fact, only a few Earth-like planets have been found so far.

b. The Sun is among the brightest 7 % of stars. Of all stars, 80 %, are red M dwarfs, 8 % are orange K dwarfs, and 5 % are white D dwarfs: these are all faint. Almost all stars visible in the night sky belong to the bright 7 %. Some of these are from the rare blue O class or blue B class (such as Spica, Regulus and Rigel); 0.7 % consist of the bright, white A class (such as Sirius, Vega and Altair); 2 % are of the yellow–white F class (such as Canopus, Procyon and Polaris); 3.5 % are of the yellow G class (like Alpha Centauri, Capella or the Sun). Exceptions include the few visible K giants, such as Arcturus and Aldebaran, and the rare M supergiants, such as Betelgeuse and Antares. More on stars later on.

Vol. II, page 228 Vol. V, page 264

Ref. 187

- c. For more details on *microscopic* aggregates, see the table of composites..
- d. It is estimated that there are up to 10^{20} *small solar system bodies* (asteroids, planetoids, minor planets, meteroids) that are heavier than 100 kg in the solar system. Incidentally, no asteroids between Mercury and the Sun the hypothetical *Vulcanoids* have been found so far.

CURIOSITIES AND FUN CHALLENGES ABOUT LAGRANGIANS

When Lagrange published his book *Mécanique analytique*, in 1788, it formed one of the high points in the history of mechanics. He was proud of having written a systematic exposition of mechanics without a single figure. Obviously the book was difficult to read and was not a sales success. Therefore his methods took another generation to come into general use.

* *

Given that action is the basic quantity describing motion, we can define energy as action per unit time, and momentum as action per unit distance. The *energy* of a system thus describes how much it changes over time, and the *momentum* how much it changes over distance. What are angular momentum and rotational energy?

Challenge 411 s

* *

'In nature, effects of telekinesis or prayer are impossible, as in most cases the change inside the brain is much smaller than the change claimed in the outside world.' Is this argument correct?

Challenge 412 s






FIGURE 159 Refraction of light is due to travel-time optimization.

In Galilean physics, the Lagrangian is the difference between kinetic and potential energy. Later on, this definition will be generalized in a way that sharpens our understanding of this distinction: the Lagrangian becomes the difference between a term for free particles and a term due to their interactions. In other words, particle motion is a continuous compromise between what the particle would do if it were free and what other particles want it to do. In this respect, particles behave a lot like humans beings.

How is action measured? What is the best device or method to measure action? Challenge 413 ny

Explain: why is T + U constant, whereas T - U is minimal? Challenge 414 s

In nature, the sum T + U of kinetic and potential energy is *constant* during motion (for closed systems), whereas the action is minimal. Is it possible to deduce, by combining these two facts, that systems tend to a state with minimum potential energy?

Challenge 415 s

Another minimization principle can be used to understand the construction of animal bodies, especially their size and the proportions of their inner structures. For example, Ref. 188 the heart pulse and breathing frequency both vary with animal mass m as $m^{-1/4}$, and the dissipated power varies as $m^{3/4}$. It turns out that such exponents result from three properties of living beings. First, they transport energy and material through the organism via a branched network of vessels: a few large ones, and increasingly many smaller ones. Secondly, the vessels all have the same minimum size. And thirdly, the networks are optimized in order to minimize the energy needed for transport. Together, these relations explain many additional scaling rules; they might also explain why animal lifespan scales as $m^{-1/4}$, or why most mammals have roughly the same number of heart beats in a lifetime.

Page 113

Ref. 189

A competing explanation, using a different minimization principle, states that quarter powers arise in any network built in order that the flow arrives to the destination by the most direct path.



* *

The minimization principle for the motion of light is even more beautiful: light always takes the path that requires the shortest travel time. It was known long ago that this idea describes exactly how light changes direction when it moves from air to water. In water, light moves more slowly; the speed ratio between air and water is called the *refractive index* of water. The refractive index, usually abbreviated n, is material-dependent. The value for water is about 1.3. This speed ratio, together with the minimum-time principle, leads to the 'law' of refraction, a simple relation between the sines of the two angles. Can you deduce it?

Challenge 416 s

Challenge 417 s

* *

Can you confirm that all the mentioned minimization principles – that for the growth of trees, that for the networks inside animals, that for the motion of light – are special cases of the principle of least action? In fact, this is the case for *all* known minimization principles in nature. Each of them, like the principle of least action, is a principle of least change.

* *

In Galilean physics, the value of the action depends on the speed of the observer, but not on his position or orientation. But the action, when properly defined, should *not* depend on the observer. All observers should agree on the value of the observed change. Only special relativity will fulfil the requirement that action be independent of the observer's speed. How will the relativistic action be defined?

Challenge 418 s

Challenge 419 s

* *

Measuring all the change that is going on in the universe presupposes that the universe is a physical system. Is this the case?

* *

One motion for which action is particularly well minimized in nature is dear to us: walking. Extensive research efforts try to design robots which copy the energy saving functioning and control of human legs. For an example, see the website by Tao Geng at cswww. essex.ac.uk/tgeng/research.html.

* *

Challenge 420 d Can you prove the following integration challenge?

$$\int_0^{\varphi} \sec t \, \mathrm{d}t = \ln \tan(\frac{\pi}{4} + \frac{\varphi}{2}) \tag{78}$$

What is the shape of the ideal halfpipe for skateboarding? What does 'ideal' imply? Which requirement leads to a *cycloid*? Which requirement speaks against a *cycloid*?

* *

Page 113 As mentioned above, animal death is a physical process and occurs when an animal has



consumed or metabolized around 1 GJ/kg. Show that the total action of an animal scales Challenge 422 e $\,$ as $M^{5/4}$.

SUMMARY ON ACTION

Systems move by minimizing change. Change, or action, is the time average of kinetic energy minus potential energy. The statement 'motion minimizes change' contains motion's predictability, its continuity and its simplicity.

Systems move by minimizing change. Equivalently, systems move by maximizing the elapsed time between two situations.

Systems move by minimizing change. In the next chapters we show that this is a strong statement: change minimization implies the observer-invariance, conservation, mirror-invariance, reversibility and relativity of motion.



(Am Anfang war die Symmetrie.*

Werner Heisenberg

Ref. 191

hat *all* observers agree. Now, whenever an observation stays exactly he *same* when switching from one observer to another, we call the observation *invariant* or *absolute* or *symmetric*. Whenever an observation *changes* when switching from one observer to another, we call it *relative*. To explore relativity thus means to explore symmetry.

Page 208

Symmetry is invariance after change. Change of observer or point of view is one such possible change, as can be some change operated on the observation itself. For example, a forget-me-not flower, shown in Figure 160, is symmetrical because it looks the same after turning it, or after turning around it, by 72 degrees; many fruit tree flowers have the same symmetry. One also says that under certain changes of viewpoint the flower has an *invariant property*, namely its shape. If many such viewpoints are possible, one talks about a *high* symmetry, otherwise a *low* symmetry. For example, a four-leaf clover has a higher symmetry than a usual, three-leaf one. In physics, the viewpoints are often called *frames of reference*.

Ref. 192

When we speak about symmetry in flowers, in everyday life, in architecture or in the arts we usually mean mirror symmetry, rotational symmetry or some combination. These are *geometric symmetries*. Like all symmetries, geometric symmetries imply invariance under specific change operations. The complete list of geometric symmetries is known for a long time. Table 34 gives an overview of the basic types. Figure 161 and Figure 162 give some important examples. Additional geometric symmetries include *colour symmetries*, where colours are exchanged, and *spin groups*, where symmetrical objects do not contain only points but also spins, with their special behaviour under rotations. Also combinations with scale symmetry, as they appear in fractals, and variations on curved backgrounds are extension of the basic table.

A *high* symmetry means that *many* possible changes leave an observation invariant. At first sight, not many objects or observations in nature seem to be symmetrical: after all, geometric symmetry is more the exception than the rule. But this is a fallacy. On the contrary, we can deduce that nature as a whole is symmetric from the simple fact that we have the ability to talk about it! Moreover, the symmetry of nature is considerably higher

Challenge 423 s

^{* &#}x27;In the beginning, there was symmetry.'



FIGURE 160 Forget-me-not, also called *Myosotis* (Boraginaceae) (© Markku Savela).

TABLE 34 The classification and the number of simple geometric symmetries.

DIMENSION	REPETI- TION TYPES	TRANSLA	TRANSLATIONS						
	11120	0	1	2	3				
		POINT	LINE	PLANE	SPACE				
		GROUPS	GROUPS	GROUPS	GROUPS				
1	1 row	2	2	n.a.	n.a.				
2	5 nets	10 crystal groups	7 friezes	17 wall-papers	n.a.				
3	14 lattices	32 crystal groups	75 rods	80 layers	230 crystal structures				

than that of a forget-me-not or of any other symmetry from Table 34. A consequence of this high symmetry is, among others, the famous expression $E_0 = mc^2$.

WHY CAN WE THINK AND TALK ABOUT THE WORLD?

The hidden harmony is stronger than the apparent.

Heraclitus of Ephesus, about 500 BCE

Ref. 193

Why can we understand somebody when he is talking about the world, even though we are not in his shoes? We can for two reasons: because most things look *similar* from different viewpoints, and because most of us have already had similar experiences *beforehand*.

'Similar' means that what we and what others observe somehow correspond. In other words, many aspects of observations do not depend on viewpoint. For example, the number of petals of a flower has the same value for all observers. We can therefore say that this quantity has the highest possible symmetry. We will see below that mass is another such example. Observables with the highest possible symmetry are called *scalars* in physics.




The 17 wallpaper patterns and a way to identify them quickly.

Is the maximum rotation order 1, 2, 3, 4 or 6?

Is there a mirror (m)? Is there an indecomposable glide reflection (g)?

Is there a rotation axis on a mirror? Is there a rotation axis not on a mirror?



Every pattern is identified according to three systems of notation:



- 442 The Conway-Thurston notation.
- p4 The International Union of Crystallography notation.
- S442 The Montesinos notation, as in his book
 - "Classical Tesselations and Three Manifolds"

FIGURE 161 The full list of possible symmetries of wallpaper patterns, the so-called wallpaper groups, their usual names, and a way to distinguish them (© Dror Bar-Natan).



Motion Mountain – The Adventure of Physics pdf file available free of charge at www.motionmountain.net

Crystal system

Crystall class or crystal group

Triclinic system

(three axes, none at right angles)





 C_{i}

 C_1

Monoclinic system

(two axes at right angles, a third not)



 C_2





 C_s or C_{1h}

 C_{2h}

Orthorhombic system (three unequal axes

(three unequal axes at right angles)



 D_2





 C_{2v}

 D_{2h}

Tetragonal system

(three axes at right angles, one unequal)











 C_{4v}





Trigonal system

(three equal axes at 120 degrees, a fourth at right angles with threefold symmetry)



 C_3



 S_6



 D_3



 C_{3v}



D_{3d}

Hexagonal system

(three equal axes at 120 degrees, a fourth at right angles with sixfold symmetry)



 C_6





 C_{6h}









Cubic or isometric system (three equal axes at right angles)











FIGURE 162 The full list of possible symmetries of units cells in crystals, the *crystallographic point* groups or *crystal groups* or *crystal classes* (© Jonathan Goss, after Neil Ashcroft and David Mermin).



Other aspects change from observer to observer. For example, the apparent size varies with the distance of observation. However, the actual size is observer-independent. In general terms, any type of *viewpoint-independence* is a form of symmetry, and the observation that two people looking at the same thing from different viewpoints can understand each other proves that nature is symmetric. We start to explore the details of this symmetry in this section and we will continue during most of the rest of our hike.

In the world around us, we note another general property: not only does the same phenomenon look similar to different observers, but *different* phenomena look similar to the *same* observer. For example, we know that if fire burns the finger in the kitchen, it will do so outside the house as well, and also in other places and at other times. Nature shows *reproducibility*. Nature shows no surprises. In fact, our memory and our thinking are only possible because of this basic property of nature. (Can you confirm this?) As we will see, reproducibility leads to additional strong restrictions on the description of nature.

Challenge 424 s

Without viewpoint-independence and reproducibility, talking to others or to one-self would be impossible. Even more importantly, we will discover that viewpoint-independence and reproducibility do more than determine the possibility of talking to each other: they also fix much (but not all) of the *content* of what we can say to each other. In other words, we will see that most of our description of nature follows logically, almost without choice, from the simple fact that we can talk about nature to our friends.

VIEWPOINTS

Toleranz ... ist der Verdacht der andere könnte Recht haben.*

Kurt Tucholsky (1890-1935), German writer

Toleranz – eine Stärke, die man vor allem dem politischen Gegner wünscht.** Wolfram Weidner (b. 1925) German journalist

When a young human starts to meet other people in childhood, he quickly finds out that certain experiences are shared, while others, such as dreams, are not. Learning to make this distinction is one of the adventures of human life. In these pages, we concentrate on a section of the first type of experiences: *physical* observations. However, even among these, distinctions are to be made. In daily life we are used to assuming that weights, volumes, lengths and time intervals are independent of the viewpoint of the observer. We can talk about these observed quantities to anybody, and there are no disagreements over their values, provided they have been measured correctly. However, other quantities do depend on the observer. Imagine talking to a friend after he jumped from one of the trees along our path, while he is still falling downwards. He will say that the forest floor is approaching with high speed, whereas the observer below will maintain that the floor is stationary. Obviously, the difference between the statements is due to their different viewpoints. The velocity of an object (in this example that of the forest floor or of the friend himself) is thus a less symmetric property than weight or size. Not all observers agree on its value.



^{* &#}x27;Tolerance ... is the suspicion that the other might be right.'

^{** &#}x27;Tolerance - a strength one mainly wishes to political opponents.'

In the case of viewpoint-dependent observations, understanding is still possible with the help of a little effort: each observer can imagine observing from the point of view of the other, and check whether the imagined result agrees with the statement of the other.* If the statement thus imagined and the actual statement of the other observer agree, the observations are consistent, and the difference in statements is due only to the different viewpoints; otherwise, the difference is fundamental, and they cannot agree or talk. Using this approach, you can even argue whether human feelings, judgements, or tastes arise from fundamental differences or not.

Challenge 425 s

The distinction between viewpoint-independent (invariant) and viewpointdependent quantities is an essential one. Invariant quantities, such as mass or shape, describe intrinsic properties, and quantities depending on the observer make up the state of the system. Therefore, we must answer the following questions in order to find a *complete* description of the state of a physical system:

- Which viewpoints are possible?
- How are descriptions transformed from one viewpoint to another?
- Which observables do these symmetries admit?
- What do these results tell us about motion?

In the discussion so far, we have studied viewpoints differing in location, in orientation, in time and, most importantly, in motion. With respect to each other, observers can be at rest, move with constant speed, or accelerate. These 'concrete' changes of viewpoint are those we will study first. In this case the requirement of consistency of observations made by different observers is called the *principle of relativity*. The symmetries associated with this type of invariance are also called *external* symmetries. They are listed in Table 36.

Page 141 Page 243

Vol. III, page 76

A second class of fundamental changes of viewpoint concerns 'abstract' changes. Viewpoints can differ by the mathematical description used: such changes are called changes of gauge. They will be introduced first in the section on electrodynamics. Again, it is required that all statements be consistent across different mathematical descriptions. This requirement of consistency is called the *principle of gauge invariance*. The associated symmetries are called *internal* symmetries.

The third class of changes, whose importance may not be evident from everyday life, is that of the behaviour of a system under exchange of its parts. The associated invariance is called *permutation symmetry*. It is a *discrete* symmetry, and we will encounter it when we explore quantum theory.

Vol. IV, page 97

The three consistency requirements described above are called 'principles' because these basic statements are so strong that they almost completely determine the 'laws' of physics, as we will see shortly. Later on we will discover that looking for a complete description of the state of objects will also yield a complete description of their intrinsic properties. But enough of introduction: let us come to the heart of the topic.

^{*} Humans develop the ability to imagine that others can be in situations different from their own at the age of about four years. Therefore, before the age of four, humans are unable to conceive special relativity; afterwards, they can.





otion Mountain – The Adventure of Physics | pdf file available free of charge at www.motionmountain.net | Copyright © Christoph Schillter November 1997–Marc

SYMMETRIES AND GROUPS

Because we are looking for a description of motion that is complete, we need to understand and describe the *full set of symmetries* of nature. But what is symmetry?

A system is said to be *symmetric* or to possess a *symmetry* if it appears identical when observed from different viewpoints. We also say that the system possesses an *invariance* under change from one viewpoint to the other. Viewpoint changes are called *symmetry operations* or *transformations*. A symmetry is thus a transformation, or more generally, a set of transformations that leaves a system invariant. However, a symmetry is more than a set: the successive application of two symmetry operations is another symmetry operation. In other terms, a symmetry is a set $G = \{a, b, c, ...\}$ of elements, the transformations, together with a binary operation \circ called *concatenation* or *multiplication* and pronounced 'after' or 'times', in which the following properties hold for all elements a, b and c:

associativity, i.e.,
$$(a \circ b) \circ c = a \circ (b \circ c)$$

a neutral element e exists such that $e \circ a = a \circ e = a$
an inverse element a^{-1} exists such that $a^{-1} \circ a = a \circ a^{-1} = e$. (79)

Any set that fulfils these three defining properties, or axioms, is called a *(mathematical) group*. Historically, the notion of group was the first example of a mathematical structure which was defined in a completely abstract manner.* Can you give an example of a group taken from daily life? Groups appear frequently in physics and mathematics, because symmetries are almost everywhere, as we will see.** Can you list the symmetry operations of the pattern of Figure 163?

Challenge 426 s Ref. 195 Challenge 428 s

REPRESENTATIONS

Challenge 429 e

Looking at a symmetric and composed system such as the one shown in Figure 163, we notice that each of its parts, for example each red patch, belongs to a set of similar objects, usually called a *multiplet*. Taken as a whole, the multiplet has (at least) the symmetry properties of the whole system. For some of the coloured patches in Figure 163 we need *four* objects to make up a full multiplet, whereas for others we need *two*, or only *one*, as in the case of the central star. In fact, in any symmetric system each part can be classified according to what type of multiplet it belongs to.

Therefore we have two challenges. Not only do we need to find all symmetries of nature; throughout our mountain ascent we also need to determine the multiplet for every

Challenge 427 e

A subset $G_1 \subset G$ of a group G can itself be a group; one then calls it a *subgroup* and often says sloppily that G is *larger* than G_1 or that G is a *higher* symmetry group than G_1 .



^{*} The term 'group' is due to Evariste Galois (1811–1832), its structure to Augustin-Louis Cauchy (1789–1857) and the axiomatic definition to Arthur Cayley (1821–1895).

^{**} In principle, mathematical groups need not be symmetry groups; but it can be proven that all groups can be seen as transformation groups on some suitably defined mathematical space, so that in mathematics we can use the terms 'symmetry group' and 'group' interchangeably.

A group is called *Abelian* if its concatenation operation is commutative, i.e., if $a \circ b = b \circ a$ for all pairs of elements a and b. In this case the concatenation is sometimes called *addition*. Do rotations form an Abelian group?




FIGURE 163 A Hispano-Arabic ornament from the Governor's Palace in Sevilla (© Christoph Schiller).

part of nature that we observe. Above all, we will need to do this for the smallest parts found in nature, the elementary particles.

A *multiplet* is a set of parts that transform into each other under all symmetry transformations. Mathematicians often call abstract multiplets *representations*. By specifying to which multiplet a component belongs, we describe in which way the component is part of the whole system. Let us see how this classification is achieved.

In mathematical language, symmetry transformations are often described by matrices. For example, in the plane, a reflection along the first diagonal is represented by the matrix

$$D(\text{refl}) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} , \tag{80}$$

since every point (x, y) becomes transformed to (y, x) when multiplied by the matrix

Challenge 430 e

D(refl). Therefore, for a mathematician a *representation* of a symmetry group G is an assignment of a matrix D(a) to each group element a such that the representation of the concatenation of two elements a and b is the product of the representations D of the elements:

$$D(a \circ b) = D(a)D(b) . \tag{81}$$

For example, the matrix of equation (80), together with the corresponding matrices for all the other symmetry operations, have this property.*

For every symmetry group, the construction and classification of all possible representations is an important task. It corresponds to the classification of all possible multiplets a symmetric system can be made of. Therefore, if we understand the classification of all multiplets and parts which can appear in Figure 163, we will also understand how to classify all possible parts of which an object or an example of motion can be composed!

A representation D is called *unitary* if all matrices D(a) are unitary.** Almost all representations appearing in physics, with only a handful of exceptions, are unitary: this term is the most restrictive, since it specifies that the corresponding transformations are one-to-one and invertible, which means that one observer never sees more or less than another. Obviously, if an observer can talk to a second one, the second one can also talk to the first.

The final important property of a multiplet, or representation, concerns its structure. If a multiplet can be seen as composed of sub-multiplets, it is called *reducible*, else *irreducible*; the same is said about representations. The irreducible representations obviously cannot be decomposed any further. For example, the (almost perfect) symmetry group of Figure 163, commonly called D_4 , has eight elements. It has the general, faithful, unitary

In general, if a mapping f from a group G to another G' satisfies

$$f(a \circ_G b) = f(a) \circ_{G'} f(b), \qquad (82)$$

the mapping f is called an *homomorphism*. A homomorphism f that is one-to-one (injective) and onto (surjective) is called a *isomorphism*. If a representation is also injective, it is called *faithful*, *true* or *proper*.

In the same way as groups, more complex mathematical structures such as rings, fields and associative algebras may also be represented by suitable classes of matrices. A representation of the field of complex numbers is given later on.

Vol. IV, page 195

** The transpose A^T of a matrix A is defined element-by-element by $(A^T)_{ik} = A_{ki}$. The complex conjugate A^* of a matrix A is defined by $(A^*)_{ik} = (A_{ik})^*$. The adjoint A^\dagger of a matrix A is defined by $A^\dagger = (A^T)^*$. A matrix is called symmetric if $A^T = A$, orthogonal if $A^T = A^{-1}$, Hermitean or self-adjoint (the two are synonymous in all physical applications) if $A^\dagger = A$ (Hermitean matrices have real eigenvalues), and unitary if $A^\dagger = A^{-1}$. Unitary matrices have eigenvalues of norm one. Multiplication by a unitary matrix is a one-to-one mapping; since the time evolution of physical systems is a mapping from one time to another, evolution is always described by a unitary matrix.

An antisymmetric or skew-symmetric matrix is defined by $A^T = -A$, an anti-Hermitean matrix by $A^{\dagger} = -A$ and an anti-unitary matrix by $A^{\dagger} = -A^{-1}$. All the corresponding mappings are one-to-one.

A matrix is *singular*, and the corresponding vector transformation is not one-to-one, if $\det A = 0$.



^{*} There are some obvious, but important, side conditions for a representation: the matrices D(a) must be invertible, or non-singular, and the identity operation of G must be mapped to the unit matrix. In even more compact language one says that a representation is a *homomorphism* from G into the group of non-singular or invertible matrices. A matrix D is invertible if its determinant $\det D$ is not zero.

Motion Mountain - The Adventure of Physics pdf file available free of charge at www.motionmountain.net | Copyright © Christoph Schiller November 1997-Ma

TABLE 35 Correspondences between the symmetries of an ornament, a flower and nature as a whole.

SYSTEM	HISPANO – ARA - BIC PATTERN	FLOWER	Мотіон
Structure and components	set of ribbons and patches	set of petals, stem	motion path and observables
System symmetry	pattern symmetry	flower symmetry	symmetry of Lagrangian
Mathematical description of the symmetry group	D_4	C ₅	in Galilean relativity: position, orientation, instant and velocity changes
Invariants	number of multiplet elements	petal number	number of coordinates, magnitude of scalars, vectors and tensors
Representations of the components	multiplet types of elements	multiplet types of components	tensors, including scalars and vectors
Most symmetric representation	singlet	part with circular symmetry	scalar
Simplest faithful representation	quartet	quintet	vector
Least symmetric representation	quartet	quintet	no limit (tensor of infinite rank)

Challenge 431 e and irreducible matrix representation

$$\begin{pmatrix} \cos n\pi/2 & -\sin n\pi/2 \\ \sin n\pi/2 & \cos n\pi/2 \end{pmatrix} n = 0..3, \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}.$$
(83)

Challenge 432 ny

The representation is an *octet*. The complete list of possible irreducible representations of the group D_4 also includes *singlets*, *doublets* and *quartets*. Can you find them all? These representations allow the classification of all the white and black ribbons that appear in the figure, as well as all the coloured patches. The most symmetric elements are singlets, the least symmetric ones are members of the quartets. The complete system is always a singlet as well.

With these concepts we are ready to talk about motion with improved precision.

SYMMETRIES, MOTION AND GALILEAN PHYSICS

Every day we experience that we are able to talk to each other about motion. It must therefore be possible to find an *invariant* quantity describing it. We already know it: it is the *action*, the measure of change. For example, lighting a match is a change. The mag-



nitude of the change is the same whether the match is lit here or there, in one direction or another, today or tomorrow. Indeed, the (Galilean) action is a number whose value is the same for each observer *at rest*, independent of his orientation or the time at which he makes his observation.

In the case of the Arabic pattern of Figure 163, the symmetry allows us to deduce the list of multiplets, or representations, that can be its building blocks. This approach must be possible for a moving system as well. Table 35 shows how. In the case of the Arabic pattern, from the various possible observation viewpoints, we deduced the classification of the ribbons into singlets, doublets, etc. For a moving system, the building blocks, corresponding to the ribbons, are the *observables*. Since we observe that nature is symmetric under many different changes of viewpoint, we can classify all observables. To do so, we first need to take the list of all viewpoint transformations and then deduce the list of all their representations.

Our everyday life shows that the world stays unchanged after changes in position, orientation and instant of observation. One also speaks of space translation invariance, rotation invariance and time translation invariance. These transformations are different from those of the Arabic pattern in two respects: they are *continuous* and they are *unbounded*. As a result, their representations will generally be continuously variable and without bounds: they will be *quantities* or *magnitudes*. In other words, observables will be constructed with *numbers*. In this way we have deduced why numbers are *necessary* for any description of motion.*

Since observers can differ in orientation, most representations will be objects possessing a direction. To cut a long story short, the symmetry under change of observation position, orientation or instant leads to the result that all observables are either 'scalars', 'vectors' or higher-order 'tensors.'**

A *scalar* is an observable quantity which stays the same for all observers: it corresponds to a singlet. Examples are the mass or the charge of an object, the distance between two points, the distance of the horizon, and many others. Their possible values are (usually) continuous, unbounded and without direction. Other examples of scalars are the potential at a point and the temperature at a point. Velocity is obviously not a scalar; nor is the coordinate of a point. Can you find more examples and counter-examples?

Challenge 434 s

Energy is a puzzling observable. It is a scalar if only changes of place, orientation and instant of observation are considered. But energy is not a scalar if changes of observer speed are included. Nobody ever searched for a generalization of energy that is a scalar also for moving observers. Only Albert Einstein discovered it, completely by accident. More about this issue shortly.

Any quantity which has a magnitude and a direction and which 'stays the same' with respect to the environment when changing viewpoint is a *vector*. For example, the arrow between two fixed points on the floor is a vector. Its length is the same for all observers; its direction changes from observer to observer, but not with respect to its environment. On the other hand, the arrow between a tree and the place where a rainbow touches the Earth is *not* a vector, since that place does not stay fixed with respect to the environment,



Challenge 433 e

^{*} Only scalars, in contrast to vectors and higher-order tensors, may also be quantities which only take a discrete set of values, such as +1 or -1 only. In short, only scalars may be *discrete* observables.

^{**} Later on, spinors will be added to, and complete, this list.

when the observer changes.

Mathematicians say that vectors are directed entities staying invariant under coordinate transformations. Velocities of objects, accelerations and field strength are examples of vectors. (Can you confirm this?) The magnitude of a vector is a scalar: it is the same for any observer. By the way, a famous and baffling result of nineteenth-century experiments is that the velocity of a light beam is *not* a vector like the velocity of a car; the velocity of a light beam is not a vector for .* This mystery will be solved shortly.

Tensors are generalized vectors. As an example, take the moment of inertia of an object. It specifies the dependence of the angular momentum on the angular velocity. For any object, doubling the magnitude of angular velocity doubles the magnitude of angular momentum; however, the two vectors are not parallel to each other if the object is not a sphere. In general, if any two vector quantities are proportional, in the sense that doubling the magnitude of one vector doubles the magnitude of the other, but without the two vectors being parallel to each other, then the proportionality 'factor' is a (second order) *tensor*. Like all proportionality factors, tensors have a magnitude. In addition, tensors have a direction and a *shape*: they describe the connection between the vectors they relate. Just as vectors are the simplest quantities with a magnitude and a direction, so tensors are the simplest quantities with a magnitude and with a direction depending on a second, chosen direction. Vectors can be visualized as oriented arrows; tensors can be

Let us get back to the description of motion. Table 35 shows that in physical systems we always have to distinguish between the symmetry of the whole Lagrangian – corresponding to the symmetry of the complete pattern – and the representation of the observables – corresponding to the ribbon multiplets. Since the action must be a scalar, and since all observables must be tensors, Lagrangians contain sums and products of tensors only in combinations forming scalars. Lagrangians thus contain only scalar products or generalizations thereof. In short, Lagrangians always look like

visualized as oriented ellipsoids.** Can you name another example of tensor?

$$L = \alpha \, a_i b^i + \beta \, c_{jk} d^{jk} + \gamma \, e_{lmn} \, f^{lmn} + \dots \tag{84}$$

where the indices attached to the variables a, b, c etc. always come in matching pairs to be summed over. (Therefore summation signs are usually simply left out.) The Greek letters represent constants. For example, the action of a free point particle in Galilean

A vector has the same length and direction for every observer; a tensor (of rank 2) has the same determinant, the same trace, and the same sum of diagonal subdeterminants for all observers.

A vector is described mathematically by a *list* of components; a tensor (of rank 2) is described by a *matrix* of components. The rank or order of a tensor thus gives the number of indices the observable has. Can you show this?

Challenge 435 e

Page 108

Page 145

Challenge 437 s



^{*} Galilean transformations are changes of viewpoints from one observer to a second one, moving with respect to the first. 'Galilean transformation' is just a term for what happens in everyday life, where velocities add and time is the same for everybody. The term, introduced in 1908 by Philipp Frank, is mostly used as a contrast to the Lorentz transformation that is so common in special relativity.

^{**} A rank-n tensor is the proportionality factor between a rank-1 tensor, i.e., between a vector, and an rank-(n-1) tensor. Vectors and scalars are rank 1 and rank 0 tensors. Scalars can be pictured as spheres, vectors as arrows, and rank-2 tensors as ellipsoids. Tensors of higher rank correspond to more and more complex shapes.

physics was given as

$$S = \int L \, \mathrm{d}t = \frac{m}{2} \int v^2 \, \mathrm{d}t \tag{85}$$

which is indeed of the form just mentioned. We will encounter many other cases during our study of motion. *

Galileo already understood that motion is also invariant under change of viewpoints with different velocity. However, the action just given does not reflect this. It took some years to find out the correct generalization: it is given by the theory of special relativity. But before we study it, we need to finish the present topic.

REPRODUCIBILITY, CONSERVATION AND NOETHER'S THEOREM

I will leave my mass, charge and momentum to science.

Graffito

Challenge 439 ny

Page 141

The reproducibility of observations, i.e., the symmetry under change of instant of time or 'time translation invariance', is a case of viewpoint-independence. (That is not obvious; can you find its irreducible representations?) The connection has several important consequences. We have seen that symmetry implies invariance. It turns out that for *continuous* symmetries, such as time translation symmetry, this statement can be made more precise: for any continuous symmetry of the Lagrangian there is an associated conserved constant of motion and vice versa. The exact formulation of this connection is the theorem of Emmy Noether.** She found the result in 1915 when helping Albert Einstein and

Ref. 196

Challenge 438 ny

* By the way, is the usual list of possible observation viewpoints – namely different positions, different observation instants, different orientations, and different velocities – also *complete* for the action (85)? Surprisingly, the answer is no. One of the first who noted this fact was Niederer, in 1972. Studying the quantum theory of point particles, he found that even the action of a Galilean free point particle is invariant under some additional transformations. If the two observers use the coordinates (t, x) and (τ, ξ) , the action (85) is invariant under the transformations

$$\xi = \frac{rx + x_0 + vt}{\gamma t + \delta}$$
 and $\tau = \frac{\alpha t + \beta}{\gamma t + \delta}$ with $r^T r = 1$ and $\alpha \delta - \beta \gamma = 1$. (86)

where r describes the rotation from the orientation of one observer to the other, v the velocity between the two observers, and x_0 the vector between the two origins at time zero. This group contains two important special cases of transformations:

The connected, static Galilei group
$$\xi = rx + x_0 + vt$$
 and $\tau = t$

The transformation group SL(2,R)
$$\xi = \frac{x}{\gamma t + \delta}$$
 and $\tau = \frac{\alpha t + \beta}{\gamma t + \delta}$ (87)

The latter, three-parameter group includes *spatial inversion*, *dilations*, *time translation* and a set of time-dependent transformations such as $\xi = x/t$, $\tau = 1/t$ called *expansions*. Dilations and expansions are rarely mentioned, as they are symmetries of point particles only, and do not apply to everyday objects and systems. They will return to be of importance later on, however.

** Emmy Noether (b. 1882 Erlangen, d. 1935 Bryn Mawr), German mathematician. The theorem is only a sideline in her career which she dedicated mostly to number theory. The theorem also applies to gauge symmetries, where it states that to every gauge symmetry corresponds an identity of the equation of motion, and vice versa.



David Hilbert, who were both struggling and competing at constructing general relativity. However, the result applies to any type of Lagrangian.

Noether investigated continuous symmetries depending on a continuous parameter b. A viewpoint transformation is a symmetry if the action S does not depend on the value of b. For example, changing position as

$$x \mapsto x + b$$
 (88)

leaves the action

$$S_0 = \int T(v) - U(x) dt$$
 (89)

invariant, since $S(b) = S_0$. This situation implies that

$$\frac{\partial T}{\partial v} = p = \text{const} \,. \tag{90}$$

In short, symmetry under change of position implies conservation of momentum. The converse is also true.

Challenge 440 ny

In the case of symmetry under shift of observation instant, we find

$$T + U = \text{const} . (91)$$

In other words, time translation invariance implies constant energy. Again, the converse is also correct. One also says that energy and momentum are the *generators* of time and space translations.

The conserved quantity for a continuous symmetry is sometimes called the *Noether charge*, because the term *charge* is used in theoretical physics to designate conserved extensive observables. So, energy and momentum are Noether charges. 'Electric charge', 'gravitational charge' (i.e., mass) and 'topological charge' are other common examples. What is the conserved charge for rotation invariance?

Challenge 441 s

We note that the expression 'energy is conserved' has several meanings. First of all, it means that the energy of a *single* free particle is constant in time. Secondly, it means that the total energy of any number of independent particles is constant. Finally, it means that the energy of a *system* of particles, i.e., including their interactions, is constant in time. Collisions are examples of the latter case. Noether's theorem makes all of these points at the same time, as you can verify using the corresponding Lagrangians.

Challenge 442 e

But Noether's theorem also makes, or rather repeats, an even stronger statement: if energy were not conserved, time could not be defined. The whole description of nature requires the existence of conserved quantities, as we noticed when we introduced the concepts of object, state and environment. For example, we defined objects as *permanent* entities, that is, as entities characterized by conserved quantities. We also saw that the introduction of time is possible only because in nature there are 'no surprises'. Noether's theorem describes exactly what such a 'surprise' would have to be: the non-conservation of energy. However, energy jumps have never been observed – not even at the quantum level.

Page 26

Page 204

Түре

Symmetry

invariance

Vac-

SERVED UUM/

Main

EFFECT

Since symmetries are so important for the description of nature, Table 36 gives an overview of all the symmetries of nature we will encounter. Their main properties are also listed. Except for those marked as 'approximate' or 'speculative', an experimental proof of incorrectness of any of them would be a big surprise indeed.

SPACE GROUP POSSI - CON-

TABLE 36 The known symmetries of nature, with their properties; also the complete list of logical *inductions* used in the two fields.

NUM- OF AC-TOPOL-BLE

	[1, 0 1,1	0 1 11 0	10102	222	OLKILD	0 0 1.1 ,	211201
	BER OF	TION	O G Y	REPRE-	QUAN-	M A T -	
	PARAM-			SENTA-	TITY	TERIS	
	ETERS			TIONS		S Y M -	
	,					METRIC	
Geometric or spa	ace-time, ex	ternal, sy	mmetries				
Time and space translation	$R \times R^3$ [4 par.]	space, time	not compact	scalars, vectors,	momentum and energy	yes/yes	allow everyday
Rotation	SO(3) [3 par.]	space	S^2	tensors	angular momentum	yes/yes	communi- cation
Galilei boost	R ³ [3 par.]	space, time	not compact	scalars, vectors, tensors	velocity of centre of mass	yes/for low speeds	relativity of motion
Lorentz	homogeneous Lie SO(3,1) [6 par.]	_	not compact	tensors, spinors	energy- momentum $T^{\mu\nu}$	yes/yes	constant light speed
Poincaré ISL(2,C)	inhomogeneous Lie [10 par.]		not compact	tensors, spinors	energy- momentum $T^{\mu\nu}$	yes/yes	
Dilation invariance	R ⁺ [1 par.]	space- time	ray	<i>n</i> -dimen. continuum	none	yes/no	massless particles
Special conformal invariance	R ⁴ [4 par.]	space- time	R^4	<i>n</i> -dimen. continuum	none	yes/no	massless particles
Conformal invariance	[15 par.]	space- time	involved	massless tensors, spinors	none	yes/no	light cone invariance
Dynamic, intera	ction-deper	ndent sym	nmetries: g	gravity			
$1/r^2$ gravity	SO(4) [6 par.]	•	-	•	perihelion direction	yes/yes	closed orbits
Diffeomorphism	$[\infty \text{ par.}]$	space-	involved	space-	local	yes/no	perihelion

times

energymomentum

time



shift

TABLE 36 (Continued) The known symmetries of nature, with their properties; also the complete list of logical *inductions* used in the two fields.

Symmetry	Түре	SPACE	Group	Possi-	Con-	Vac-	MAIN
	[N U M -	OF AC	- T O P O L	-B L E	SERVED	uum/	EFFECT
	BER OF	TION	O G Y	REPRE-	QUAN-	M A T -	
	PARAM	-		SENTA-	TITY	TERIS	
	ETERS]			TIONS		S Y M -	
						METRIC	C

Dynamic, classical and quantum-mechanical motion symmetries

	1			•			
Motion('time') inversion T	discrete	Hilbert or phase space	discrete	even, odd	T-parity	yes/no	reversibil- ity
Parity('spatial') inversion P	discrete	Hilbert or phase space	discrete	even, odd	P-parity	yes/no	mirror world exists
Charge conjugation C	global, antilinear, anti- Hermitean	or phase space	discrete	even, odd	C-parity	yes/no	antiparti- cles exist
CPT	discrete	Hilbert or phase space	discrete	even	CPT-parity	yes/yes	makes field theory possible
Dynamic, intera	ction-deper	ndent, gau	ige symme	etries			
Electromagnetic classical gauge invariance	[∞ par.]	space of fields	un- im- portant	un- important	electric charge	yes/yes	massless light
Electromagnetic q.m. gauge inv.	Abelian Lie U(1) [1 par.]	Hilbert space	circle S ¹	fields	electric charge	yes/yes	massless photon
Electromagnetic duality	Abelian Lie U(1) [1 par.]	space of fields	circle S ¹	abstract	abstract	yes/no	none
Weak gauge	non- Abelian Lie SU(2) [3 par.]	Hilbert space	as <i>SU</i> (3)	particles	weak charge	no/ approx.	
Colour gauge	non- Abelian Lie SU(3) [8 par.]	Hilbert space	as SU(3)	coloured quarks	colour	yes/yes	massless gluons
Chiral symmetry	discrete	fermions	discrete	left, right	helicity	approxi- mately	'massless' fermions ^a



TABLE 36 (Continued) The known symmetries of nature, with their properties; also the complete list of logical *inductions* used in the two fields.

Symmetry	Түре	SPACE	GROUP	Possi-	Con-	Vac-	Main
	[N U M -	OF AC	- T O P O L	-B L E	SERVED	uum/	EFFECT
	BER OF	TION	OGY	REPRE-	QUAN-	M A T -	
	PARAM	-		SENTA-	TITY	TERIS	
	ETERS]			TIONS		S Y M -	
						METRIC	

Permutation symmetries

Particle	discrete	Fock	discrete	fermions	none	n.a./yes	Gibbs'
exchange		space		and			paradox
		etc.		bosons			

Vol. III, page 256 For details about the connection between symmetry and induction, see later on. The explanation of the terms in the table will be completed in the rest of the walk. The real numbers are denoted as *R*.

a. Only approximate; 'massless' means that $m \ll m_{\rm pl}$, i.e., that $m \ll 22 \, \mu \rm g$.

CURIOSITIES AND FUN CHALLENGES ABOUT SYMMETRY

What is the path followed by four turtles starting on the four angles of a square, if each of them continuously walks at the same speed towards the next one?

* *

Challenge 444 s What is the symmetry of a simple oscillation? And of a wave?

* *

Challenge 445 s For what systems is motion reversal a symmetry transformation?

* *

Challenge 446 s What is the symmetry of a continuous rotation?

* *

A sphere has a tensor for the moment of inertia that is diagonal with three equal numbers. The same is true for a cube. Can you distinguish spheres and cubes by their rotation behaviour?

Challenge 447 s behaviour?

Challenge 443 ny

* *

Challenge 448 s Is there a motion in nature whose symmetry is perfect?

* *

Can you show that in two dimensions, *finite* objects can have only rotation and reflection symmetry, in contrast to *infinite* objects, which can have also translation and glidereflection symmetry? Can you prove that for finite objects in two dimensions, if no ro-

Mountain - The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christoph Schiller November 1997-March 2012



tation symmetry is present, there is only one reflection symmetry? And that all possible rotations are always about the same centre? Can you deduce from this that at least one point is unchanged in all symmetrical finite two-dimensional objects?

Challenge 449 e

* *

Can you show that in three dimensions, *finite* objects can have only rotation, reflection, inversion and rotatory inversion symmetry, in contrast to *infinite* objects, which can have also translation, glide-reflection, and screw rotation symmetry? Can you prove that for finite objects in three dimensions, if no rotation symmetry is present, there is only one reflection plane? And that for all inversions or rotatory inversions the centre must lie on a rotation axis or on a reflection plane? Can you deduce from this that at least one point is unchanged in all symmetrical finite three-dimensional objects?

Challenge 450 e

PARITY AND TIME INVARIANCE

The table of symmetries also list two so-called *discrete* symmetries that are important for the discussion of motion.

The first is parity inversion, or right-left symmetry. How far can you throw a stone with your other hand? Most people have a preferred hand, and the differences are quite pronounced. Does nature have such a right-left preference? In everyday life, the answer is clear: everything that happens one way can also happen in its mirrored way. This has also been tested in precision experiments; it was found that everything due to gravitation, electricity or magnetism can also happen in a mirrored way. There are no exceptions. For example, there are people with the heart on the right side; there are snails with left-handed houses; there are planets that rotate the other way. Astronomy and everyday life are *mirror-invariant*. One also says that gravitation and electromagnetism are *parity invariant*. (Later we will discover that certain rare processes not due to gravity or electromagnetism, but to the weak nuclear interaction, violate parity.)

Vol. V, page 182

The other discrete symmetry is *motion reversal*. Can things happen backwards? This question is not easy. A study of motion due to gravitation shows that such motion can always also happen in the reverse direction. In case of motion due to electricity and magnetism, such as the behaviour of atoms in gases and liquids, the question is more involved. We will discuss it in the section of thermodynamics, but we will reach the same conclusion: motion inversion is a symmetry for all processes due to gravitation and the electromagnetic interaction. Everyday motion is *reversible*. And again, certain even rarer nuclear processes will provide exceptions.

SUMMARY ON SYMMETRY

Symmetry is partial invariance to change. The simplest symmetries are geometrical: the point symmetries of flowers or translation symmetry of infinite crystals are examples.

All possible changes that leave a system invariant – i.e., all possible symmetry transformations of a system – form a mathematical group. Apart from geometrical symmetry groups, several additional symmetry groups appear in nature.

The reproducibility and predictability of nature implies several fundamental continuous symmetries: since we can *talk* about nature we can deduce that above all, nature is symmetrical under time and space translations. Motion is universal. Any universal-

re of Physics 🏻 pdf file available free of charge at www.motionmountain.net 🖰 Copyright © Christoph Schiller November 1997–March 2013



Motion Mountain – The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christoph Schiller November 1997–1


ity statement implies a symmetry. As further examples, everyday observations are found to be mirror symmetric, and simple motions are found to be symmetric under motion reversal. These are fundamental discrete symmetries.

From nature's continuous symmetries, using Noether's theorem, we can deduce conserved 'charges'. These are energy, linear momentum and angular momentum. In other words, the definition of mass, space and time, together with their symmetry properties, is *equivalent* to the conservation of energy and momenta. Conservation and symmetry are two ways to express the same property of nature. To put it simply, our ability to talk about nature means that energy, linear momentum and angular momentum are conserved.

The isolability of systems from their surroundings implies a symmetry valid for all interactions: since the behaviour of isolated systems is independent from what happens in their surroundings, interactions must have no effect at large distances.

An elegant way to uncover the 'laws' of nature is to search for nature's symmetries. In many historical cases, once this connection had been understood, physics made rapid progress. For example, Albert Einstein discovered the theory of relativity in this way, and Paul Dirac started off quantum electrodynamics. We will use the same method throughout our walk; in our final leg we will uncover some symmetries which are even more mind-boggling than those of relativity. Now, though, we will move on to the next approach to a global description of motion.





CHAPTER 10

SIMPLE MOTIONS OF EXTENDED BODIES – OSCILLATIONS AND WAVES

HE observation of change is a fundamental aspect of nature. Among all hese observations, *periodic* change is frequent aorund us. Indeed, hroughout everyday life be observe oscillations and waves: Talking, singing, hearing and seeing would be impossible without them. Exploring oscillations and waves, the next global approach to motion, is both useful and beautiful.

OSCILLATIONS

idea of how harmonic motion 'sounds'.)

Page 213 Oscillations are recurring changes, i.e., cyclic or periodic changes. Above, we defined action, and thus change, as the integral of the Lagrangian, and we defined the Lagrangian as the difference between kinetic and potential energy. One of the simplest oscillating systems in nature is a mass *m* attached to a (linear) *spring*. The Lagrangian for the mass position *x* is given by

$$L = \frac{1}{2}mv^2 - \frac{1}{2}kx^2 \,, (92)$$

where k is a quantity characterizing the spring, the so-called spring constant. The Lagrangian is due to Robert Hooke, in the seventeenth century. Can you confirm the expression?

The motion that results from this Lagrangian is periodic, and shown in Figure 164. The Lagrangian (92) thus describes the *oscillation* of the spring length over time. The motion is exactly the same as that of a long pendulum at small amplitude. The motion is called *harmonic motion*, because an object vibrating rapidly in this way produces a completely pure – or harmonic – musical sound. (The musical instrument producing the purest harmonic waves is the transverse flute. This instrument thus gives the best

The graph of this harmonic or linear oscillation, shown in Figure 164, is called a *sine curve*; it can be seen as the basic building block of all oscillations. All other, anharmonic oscillations in nature can be composed from harmonic ones, i.e., from sine curves, as we shall see shortly. Any quantity x(t) that oscillates harmonically is described by its amplitude A, its angular frequency ω and its phase φ :

$$x(t) = A\sin(\omega t + \varphi). \tag{93}$$

The amplitude and the phase depend on the way the oscillation is started. In contrast,

Challenge 451 e

Page 252




FIGURE 164 Above: the simplest oscillation, the *linear* or *harmonic* oscillation: how position changes over time, and how it is related to rotation. Below: an example of anharmonic oscillation.

the angular frequency is an intrinsic property of the system. Can you show that for the mass attached to the spring, we have $\omega = 2\pi f = 2\pi/T = \sqrt{k/m}$?

Every harmonic oscillation is described by three quantities: the *amplitude*, the *period* (the inverse of the frequency) and the *phase*. The phase distinguishes oscillations of the same amplitude and period; it defines at what time the oscillation starts. Some observed oscillation frequencies are listed in Table 37. Figure 164 shows how a harmonic oscillation is related to an imaginary rotation. As a result, the phase is best described by an angle between 0 and 2π .

Every oscillating motion continuously transforms kinetic energy into potential energy and vice versa. This is the case for the tides, the pendulum, or any radio receiver. But many oscillations also diminish in time: they are damped. Systems with large damping, such as the shock absorbers in cars, are used to avoid oscillations. Systems with *small* damping are useful for making precise and long-running clocks. The simplest measure of damping is the number of oscillations a system takes to reduce its amplitude to $1/e \approx 1/2.718$ times the original value. This characteristic number is the so-called *Q-factor*, named after the abbreviation of 'quality factor'. A poor Q-factor is 1 or less, an extremely good one is 100 000 or more. (Can you write down a simple Lagrangian for a damped oscillation with a given Q-factor?) In nature, damped oscillations do not usually keep

Challenge 453 ny

Challenge 452 s



TABLE 37 Some frequency values found in nature.

OBSERVATION	FREQUENCY
Sound frequencies in gas emitted by black holes	c. 1 fHz
Precision in measured vibration frequencies of the Sun	down to 2 nHz
Vibration frequencies of the Sun	down to c. 300 nHz
Vibration frequencies that disturb gravitational radiation detection	down to 3 μHz
Lowest vibration frequency of the Earth Ref. 198	$309\mu Hz$
Resonance frequency of stomach and internal organs (giving the 'sound in the belly' experience)	1 to 10 Hz
Common music tempo	2 Hz
Frequency used for communication by farting fish	c. 10 Hz
Sound audible to young humans	20 Hz to 20 kHz
Sound produced by loudspeaker sets (horn, electromagetic, piezoelectric, electret, plasma, laser)	<i>c.</i> 18 Hz to over 150 kHz
Adult male speaking voice, fundamental	85 to 180 Hz
Adult female speaking voice, fundamental	165 to 255 Hz
Official value, or <i>standard pitch</i> , of musical note 'A' or 'la', following ISO 16 (and of the telephone line signal in many countries)	440 Hz
Common values of musical note 'A' or 'la' used by orchestras	442 to 451 Hz
Wing beat of tiniest flying insects	c. 1000 Hz
Feather oscillation during wing beat of male club-winged manakin, a bird	c. 1000 Hz
Sound produced by various loudspeakers	30 Hz to 100 kHz
Sonar used by bats	up to over 100 kHz
Sonar used by dolphins	up to 150 kHz
Sound frequency used in ultrasound imaging	2 to 20 MHz
Quartz oscillator frequencies	20 kHz up to 350 MHz
Radio emission of atomic hydrogen, esp. in the universe	1420.4057518(1) MHz
Highest electronically generated frequency (with CMOS, in 2007)	324 GHz
Phonon (sound) frequencies measured in single crystals	up to 20 THz and more




FIGURE 165 The interior of a commercial quartz oscillator, a few millimetre in size, driven at high amplitude (QuickTime film © Microcrystal)

constant frequency; however, for the simple pendulum this remains the case to a high degree of accuracy. The reason is that for a pendulum, the frequency does not depend significantly on the amplitude (as long as the amplitude is smaller than about 20°). This is one reason why pendulums are used as oscillators in mechanical clocks.

Obviously, for a good clock, the driving oscillation must not only show small damping, but must also be independent of temperature and be insensitive to other external influences. An important development of the twentieth century was the introduction of quartz crystals as oscillators. Technical quartzes are crystals of the size of a few grains of sand; they can be made to oscillate by applying an electric signal. They have little temperature dependence and a large Q-factor, and therefore low energy consumption, so that precise clocks can now run on small batteries. The inside of a quartz oscillator is shown in Figure 165.

RESONANCE


In most physical systems that are brought to oscillate, the amplitude depends on the frequency. The selected frequencies for which the amplitude is maximal are called *resonance frequencies* or simply *resonances*. For example, the quartz oscillator of Figure 165, or the usual vibration frequencies of guitar strings are resonance frequencies.

Usually, the oscillations at which a system will oscillate when triggered by a short hit will occur at resonance frequencies. Most musical instruments are examples. Most systems have several resonance frequencies; flutes and strings are well-known examples.

In contrast to music or electronics, resonance often needs to be avoided in other situations. In buildings, earthquakes can trigger resonances; in bridges, the wind can trigger resonant oscillations; similarly, in many machines resonances need to be dampened or blocked in order to avoid that the large amplitude of a resonance destroys the system.

All systems that oscillate also emit *waves*. In fact, resonance only occurs because all oscillations are in fact *localized waves*. Indeed, oscillations only appear in *extended* systems; and oscillations are only simplified descriptions of the repetitive motion of any extended system. The complete and general repetitive motion of an extended system is the *wave*.





An example of *anharmonic* signal




FIGURE 166 Top: the main properties of a harmonic wave. Bottom: A general *periodic* signal, or anharmonic wave – here a black square wave – can be decomposed uniquely into simplest, or *harmonic* waves. The first three components (green, blue and red) and also their intermediate sum (black dotted line) are shown. This is called a *Fourier decomposition* and the general method to do this *Fourier analysis*. (© Wikimedia) Not shown: the unique decomposition into harmonic waves is even possible for *non-periodic* signals.



FIGURE 167 The measured fundamental vibration patterns of a bell. Bells – like every other source of oscillations, be it an atom, a molecule, a music instrument or the human voice – show that all oscillations in nature are due to waves. (© H. Spiess & al.).



Aotion Mountain – The Adventure of Physics 🛮 pdf file available free of charge at www.motionmountain.net 🖯 Copyright © Christoph Schiller November 1997–Mar

TABLE 38 Some wave velocities.

Wave	VELOCITY
Tsunami	around 0.2 km/s
Sound in most gases	$0.3 \pm 0.1 \text{km/s}$
Sound in air at 273 K	0.331 km/s
Sound in air at 293 K	0.343 km/s
Sound in helium at 293 K	0.983 km/s
Sound in most liquids	$1.2 \pm 0.2\mathrm{km/s}$
Seismic waves	1 to 14 km/s
Sound in water at 273 K	$1.402\mathrm{km/s}$
Sound in water at 293 K	1.482 km/s
Sound in sea water at 298 K	1.531 km/s
Sound in gold	4.5 km/s
Sound in steel	5.8 to 5.960 km/s
Sound in granite	5.8 km/s
Sound in glass (longitudinal)	4 to 5.9 km/s
Sound in beryllium (longitudinal)	12.8 km/s
Sound in boron	up to 15 km/s
Sound in diamond	up to 18 km/s
Sound in fullerene (C ₆₀)	up to 26 km/s
Plasma wave velocity in InGaAs	600 km/s
Light in vacuum	$2.998 \cdot 10^8 \text{m/s}$

WAVES: GENERAL AND HARMONIC

Waves are travelling imbalances, or, equivalently, travelling oscillations. Waves move, even though the substrate does not move. Every wave can be seen as a superposition of *harmonic* waves. Every sound effect can be thought of as being composed of harmonic waves. Harmonic waves, also called *sine waves* or *linear waves*, are the building blocks of which all internal motions of an extended body are constructed, as shown in Figure 166. Can you describe the difference in wave shape between a pure harmonic tone, a musical sound, a noise and an explosion?

Challenge 454 e

Every harmonic wave is characterized by an oscillation frequency f, a propagation (or phase) velocity c, a wavelength λ , an amplitude A and a phase φ , as can be deduced from Figure 166. Low-amplitude water waves show this most clearly; they are harmonic. In a harmonic wave, every position by itself performs a harmonic oscillation. The phase of a wave specifies the position of the wave (or a crest) at a given time. It is an angle between 0 and 2π .

The phase velocity c is the speed with which a wave maximum moves. A few examples are listed in Table 38. Can you show that frequency and wavelength in a wave are related by $f\lambda = c$?

Challenge 455 e

Waves appear inside all *extended* bodies, be they solids, liquids, gases or plasmas. Inside fluid bodies, waves are *longitudinal*, meaning that the wave motion is in the same




FIGURE 168 The formation of the shape of deep gravity waves, on and under water, from the circular motion of the water particles. Note the non-sinusoidal shape of the wave.

direction as the wave oscillation. Sound in air is an example of a longitudinal wave. Inside solid bodies, waves can also be *transverse*; in that case the wave oscillation is perpendicular to the travelling direction.

Waves appear also on *interfaces* between bodies: water-air interfaces are a well-known case. Even a saltwater-freshwater interface, so-called *dead water*, shows waves: they can appear even if the upper surface of the water is immobile. Any flight in an aeroplane provides an opportunity to study the regular cloud arrangements on the interface between warm and cold air layers in the atmosphere. Seismic waves travelling along the boundary between the sea floor and the sea water are also well-known. General surface waves are usually neither longitudinal nor transverse, but of a mixed type.

To get a first idea about waves, we have a look at water waves.

WATER WAVES

Water waves on water surfaces show a large range of fascinating phenomena. First of all, there are two different types of water waves. In the first type, the force that restores the plane surface is the surface tension of the wave. These so-called *surface tension waves* play a role on scales up to a few centimetres. In the second, larger type of waves, the restoring force is gravity, and one speaks of *gravity waves*.* The difference is easily noted by watching them: surface tension waves have a sinusoidal shape, whereas gravity waves have a shape with sharper maxima and broader troughs. This occurs because of the special way the water moves in such a wave. As shown in Figure 168, the surface water for a (short) water wave moves in circles; this leads to the typical, asymmetrical wave shape with short sharp crests and long shallow troughs. Under the crests, the water particles move in the direction of the wave motion; under the troughs, the water particles move against the wave motion. As long as there is no wind and the floor below the water is horizontal, gravity waves are also symmetric under front-to-back reflection. If the amplitude is very



^{*} Meteorologists also know of a third type: there are large wavelength waves whose restoring force is the .

FIGURE 169 The four types of water waves: a deep water gravity wave, a shallow water gravity wave, a deep water ripple and a shallow water ripple.

high, or if the wind is too strong, waves break, because a cusp angle of more than 120° is not possible.

In addition, water waves need to be distinguished according to the depth of the water, when compared to their wavelength. One speaks of *short* (or *deep water*) waves, when the depth of the water is so high that the floor below plays no role; in the opposite case one speaks of *long* (or *shallow water*) waves. It turns out that deep water waves are *dispersive*, i.e., their speed depends on their frequency, whereas shallow water waves are *non-dispersive*. The transitional region between the two cases are waves whose wavelength is between twice and twenty times the water depth.

The two classifications of water waves give *four* limit cases; they are shown in Figure 170. (The figure also shows where capillary waves, long period waves, tides and transtidal waves are located.) It is interesting to explore each of these four limit cases.

Experiments and theory show that the phase speed of *gravity waves*, the lower two cases in Figure 170, depends on the wavelength λ and on the depth of the water d in the following way:

$$c = \sqrt{\frac{g\lambda}{2\pi}} \tanh \frac{2\pi d}{\lambda} , \qquad (94)$$

where g is the acceleration due to gravity (and an amplitude much smaller than the wavelength is assumed*). The formula shows two limiting regimes. First, so-called *deep water* or *short* gravity waves appear when the water depth is larger than about half the wavelength; for deep water waves, the phase velocity is $c \approx \sqrt{g\lambda/2\pi}$, thus wavelength dependent – (all) deep waves are dispersive. Shorter deep waves are thus slower. The group velocity is *half* the phase velocity. The general effects of dispersion on wave groups are shown in Figure 171.

The usual sea wave is a deep water gravity wave, so are wakes generated by ships. Short gravity waves generated by wind are called *sea* they are generated by local winds,





^{*} The expression for the phase velocity can be derived by solving for the motion of the liquid in the linear regime, but this leads us too far from our walk.




FIGURE 170 The different types of water waves, visualized in two different diagrams using the depth d, the wave number k and the surface tension $\gamma = 72$ mPa.






FIGURE 171 A visualisation of group velocity (blue) and phase velocity (red) for different types of wave dispersion (QuickTime film © ISVR, University of Southhampton).

and *swell* if they are generated by distant winds. The typical phase speed of a gravity wave is of the order of the wind speed that generates it. Therefore, as surfers know, waves on a shore that are due to a distant storm arrive separately: first the long period waves, then the short period waves.

The typical *wake* generated by a ship is made of waves that have the phase velocity of the ship. These waves form a wave group, and it travels with half that speed. Therefore, from a ship's point of view, the wake trails the ship. Wakes are behind the ship, because the group velocity is lower than the phase velocity. (For more about wakes, see below.)

The second limiting regime are *shallow water* or *long* gravity waves. They appear when the depth is less than 1/20th or 5% of the wavelength; in this case, the phase velocity is $c \approx \sqrt{gd}$, there is no dispersion, and the group velocity is the *same* as the phase velocity. In shallow water waves, water particles move on very flat elliptic paths.

For example, the tide is a shallow gravity wave. Apart from tides, the most impressive shallow gravity waves are *tsunamis*, the large waves triggered by submarine earthquakes. (The Japanese name is composed of *tsu*, meaning harbour, and *nami*, meaning wave.) Since tsunamis are shallow waves, they show no (or little) dispersion and thus travel over long distances; they can go round the Earth several times. Typical oscillation times of tsunamis are between 6 and 60 minutes, giving wavelengths between 70 and 700 km and speeds in the open sea of 200 to 250 m/s, similar to that of a jet plane. Their amplitude on the open sea is often of the order of 10 cm; however, the amplitude scales with depth d as $1/d^4$ and heights up to 40 m have been measured at the shore. This was the order of magnitude of the large and disastrous tsunami observed in the Indian Ocean on 26 December 2004 and the one in Japan in 2011 that destroyed several nuclear power plants. Tsunamis can also be used to determine the ocean depth by measuring the speed of tsunamis. This allowed to deduce, long before sonar and other high-tech systems were available, that the North Pacific has a depth of around 4 to 4.5 km.

The upper two cases in Figure 170 are the surface tension waves. (The surface tension of water is 72 mPa.) The first of these limit regimes are ripples on deep water. The phase velocity is $c = \sqrt{\gamma k/\rho}$. As mentioned above, all deep waves are dispersive. Indeed, the group velocity of ripples on deep water is 3/2 times the phase velocity. Therefore ripples

Page 279

Challenge 456 e

Challenge 457 e

steam ahead of a boat, whereas wakes trail behind.

The minimum speed is the reason for what we see when throw a *pebble* in a lake. A typical pebble creates ripples with a wavelength of about 1 cm. For waves in this region, there is a minimum a group velocity of 17.7 cm/s and a minimum phase velocity of around 23 cm/s. When a pebble falls into the water, it creates waves of various wavelengths; those with a wavelength of about 1 cm are the slowest ones are seen most clearly. The minimum phase velocity for ripples also means that insects walking on water generate no waves if they move more slowly than the minimum phase velocity; thus they feel little drag and can walk easily.

The final case of waves are ripples on shallow water. An example are the waves emitted by raindrops falling in a shallow puddle, say with a depth of 1 mm or less. The phase velocity is $c = \sqrt{\gamma dk^2/\rho}$; the group velocity has twice that value. Ripples on shallow waves are dispersive.

Figure 170 shows the four types of water waves. The general dispersion relation for water waves is $\omega^2 = (gk + \gamma k^3/\rho) \tanh kd$. Several other types of water waves also exist, such as seiches, internal waves and solitons of various kinds. We will only explore the last case in some detail, later on.

Waves and their motion

Waves move. Therefore, any study of motion must include the study of wave motion. We know from experience that waves can hit or even damage targets; thus every wave carries energy and momentum, even though (on average) no matter moves along the wave propagation direction. The *energy E* of a wave is the sum of its kinetic and potential energy. The kinetic energy (density) depends on the temporal change of the displacement u at a given spot: rapidly changing waves carry a larger kinetic energy. The potential energy (density) depends on the gradient of the displacement, i.e., on its spatial change: steep waves carry a larger potential energy than shallow ones. (Can you explain why the potential energy does not depend on the displacement itself?) For harmonic waves, i.e., sinusoidal waves, propagating along the direction z, each type of energy is proportional to the square of its respective displacement change:

 $E \sim \left(\frac{\partial u}{\partial t}\right)^2 + v^2 \left(\frac{\partial u}{\partial z}\right)^2. \tag{95}$

How is the energy density related to the frequency?

The *momentum* of a wave is directed along the direction of war.

The *momentum* of a wave is directed along the direction of wave propagation. The momentum value depends on both the temporal and the spatial change of displacement u. For harmonic waves, the momentum (density) P is proportional to the product of these two quantities:

$$P_z \sim \frac{\partial u}{\partial t} \frac{\partial u}{\partial z} \,. \tag{96}$$

When two linear wave trains collide or interfere, the total momentum is conserved throughout the collision. An important consequence of momentum conservation is that waves that are reflected by an obstacle do so with an outgoing angle equal to minus the

Challenge 458 e

Challenge 459 e

Page 268

Challenge 460 s

Challenge 461 ny

Ref. 199

f file available free of charge at www.motionmountain.net 🛮 Copyright © Christoph Schiller November 1997–March 2012




FIGURE 172 The six main properties of the motion of waves.

Challenge 462 s infalling angle. What happens to the phase?

In summary, waves, like moving bodies, carry energy and momentum. In simple terms, if you shout against a wall, the wall is hit. This hit, for example, can start avalanches on snowy mountain slopes. In the same way, waves, like bodies, can carry also angular momentum. (What type of wave is necessary for this to be possible?) However, the motion of waves also differs from the motion of bodies. Six main properties distinguish the motion of waves from the motion of bodies.

Challenge 463 ny

- 1. Waves can add up or cancel each other out; thus they can interpenetrate each other. These effects, called *superposition* and *interference*, are strongly tied to the linearity of most waves.
- 2. Waves, such as sound, can go around corners. This is called *diffraction*.
- 3. Waves change direction when they change medium. This is called *refraction*.
- 4. Waves can have a frequency-dependent propagation speed. This is called *dispersion*.
- 5. Often, the wave amplitude decreases over time: waves show *damping*.
- 6. Transverse waves in three dimensions can oscillate in different directions: they show *polarization*.

Material bodies in everyday life do not behave in these ways when they move. The six






FIGURE 173 Interference of two circular or spherical waves emitted in phase: a snapshot of the amplitude (left), most useful to describe observations of water waves, and the distribution of the time-averaged intensity (right), most useful to describe interference of light waves (© Rüdiger Paschotta).

wave effects appear because wave motion is the motion of *extended* entities. The famous debate whether electrons or light are waves or particles thus requires us to check whether these effects specific to waves can be observed or not. This is one topic of quantum theory. Before we study it, can you give an example of an observation that automatically implies that the specific motion *cannot* be a wave?

Challenge 464 s

As a result of having a frequency f and a propagation or phase velocity c, all sine waves are characterized by the distance λ between two neighbouring wave crests: this distance is called the wavelength λ . All waves obey the basic relation

$$\lambda f = c . (97)$$

In many cases the phase velocity c depends on the wavelength of the wave. For example, this is the case for many water waves. This change of speed with wavelength is called *dispersion*. In contrast, the speed of sound in air does *not* depend on the wavelength (to a high degree of accuracy). Sound in air shows (almost) no dispersion. Indeed, if there were dispersion for sound, we could not understand each other's speech at larger distances.

Now comes a surprise. Waves can also exist in empty space. Both light and gravity waves are examples. The exploration of electromagnetism and relativity will tell us more about their specific properties. Here is an appetizer. Light is a wave. In everyday life we do not experience light as a wave, because the wavelength is only around one two-thousandth of a millimetre. But light shows all six effects typical of wave motion. A rainbow, for example, can only be understood fully when the last five wave effects are taken into account. Diffraction and interference can even be observed with your fingers only. Can you tell how?

Vol. III, page 88

Challenge 465 s

Like every anharmonic oscillation, every anharmonic wave can be decomposed into

Page 252 sine waves. Figure 166 gives examples. If the various sine waves contained in a disturbance propagate differently, the original wave will change in shape while it travels. That is the reason why an echo does not sound exactly like the original sound; for the same reason, a nearby thunder and a far-away one sound different. These are effects of the weak dispersion of sound waves.

All systems which oscillate also emit waves. Any radio or TV receiver contains oscillators. As a result, any such receiver is also a (weak) transmitter; indeed, in some countries the authorities search for people who listen to radio without permission listening to the radio waves emitted by these devices. Also, inside the human ear, numerous tiny structures, the hair cells, oscillate. As a result, the ear must also emit sound. This prediction, made in 1948 by Tommy Gold, was finally confirmed in 1979 by David Kemp. These so-called *otoacoustic emissions* can be detected with sensitive microphones; they are presently being studied in order to unravel the still unknown workings of the ear and in order to diagnose various ear illnesses without the need for surgery.

Ref. 200

Since any travelling disturbance can be decomposed into sine waves, the term 'wave' is used by physicists for all travelling disturbances, whether they look like sine waves or not. In fact, the disturbances do not even have to be travelling. Take a standing wave: is it a wave or an oscillation? Standing waves do not travel; they are oscillations. But a standing wave can be seen as the superposition of two waves travelling in opposite directions. In fact, in nature, any object that we call 'oscillating' or 'vibrating' is extended, and its oscillation or vibration is always a standing wave (can you confirm this?); so we can say that in nature, *all oscillations are special forms of waves*.

Challenge 466 e

Page 252

The most important travelling disturbances are those that are localized. Figure 166 shows an example of a localized wave, also called a *wave group* or *pulse*, together with its decomposition into harmonic waves. Wave groups are used to talk and as signals for communication.

WHY CAN WE TALK TO EACH OTHER? - HUYGENS' PRINCIPLE

The properties of our environment often disclose their full importance only when we ask simple questions. Why can we use the radio? Why can we talk on mobile phones? Why can we listen to each other? It turns out that a central part of the answer to these questions is that the space we live has an *odd* numbers of dimensions.

In spaces of *even* dimension, it is impossible to talk, because messages do not stop. This is an important result which is easily checked by throwing a stone into a lake: even after the stone has disappeared, waves are still emitted from the point at which it entered the water. Yet, when we stop talking, no waves are emitted any more. Waves in two and three dimensions thus behave differently.

In three dimensions, it is possible to say that the propagation of a wave happens in the following way: Every point on a wave front (of light or of sound) can be regarded as the source of secondary waves; the surface that is formed by the envelope of all the secondary waves determines the future position of the wave front. The idea is illustrated in Figure 174. It can be used to describe, without mathematics, the propagation of waves, their reflection, their refraction, and, with an extension due to Augustin Fresnel, their diffraction. (Try!)

Challenge 467 e

This idea was first proposed by Christiaan Huygens in 1678 and is called Huygens'




FIGURE 174 Wave propagation as a consequence of Huygens' principle.




FIGURE 175 An impossible gravity water wave: the centre is never completely flat.

principle. Almost two hundred years later, Gustav Kirchhoff showed that the principle is a consequence of the wave equation in three dimensions, and thus, in the case of light, a consequence of Maxwell's field equations.

But the description of wave fronts as envelopes of secondary waves has an important limitation. It is not correct in two dimensions (even though Figure 174 is two-dimensional!). In particular, it does not apply to water waves. Water wave propagation *cannot* be calculated in this way in an exact manner. (It is only possible if the situation is limited to a wave of a single frequency.) It turns out that for water waves, secondary waves do not only depend on the wave front of the primary waves, but depend also on their interior. The reason is that in two (and other even) dimensions, waves of different frequency necessarily have different speeds. (Shallow waves are not counterexamples; they are intrinsically three-dimensional.) And a stone falling into water generates waves of many frequencies. In contrast, in three (and larger odd) dimensions, waves of all frequencies have the same speed.

We can also say that Huygens' principle holds if the wave equation is solved by a circular wave leaving no amplitude behind it. Mathematicians translate this by requiring that the evolving delta function $\delta(c^2t^2-r^2)$ satisfies the wave equation, i.e., that $\partial_t^2\delta=c^2\Delta\delta$. The delta function is that strange 'function' which is zero everywhere except at the origin, where it is infinite. A few more properties describe the precise way in which this happens.* It turns out that the delta function is a solution of the wave equation only if the space dimension is odd and at least three. In other words, while a spherical wave pulse is possible, a circular pulse is not: there is no way to keep the centre of an expanding wave quiet. (See Figure 175.) That is exactly what the stone experiment shows. You

^{*} The main property is $\int \delta(x) dx = 1$. In mathematically precise terms, the delta 'function' is a *distribution*.

can try to produce a circular pulse (a wave that has only a few crests) the next time you are in the bathroom or near a lake: you will not succeed.

In summary, the reason a room gets dark when we switch off the light, is that we live in a space with a number of dimensions which is odd and larger than one.

WHY IS MUSIC SO BEAUTIFUL?

Music works because it connects to emotions. And it does so, among others, by reminding us of the sounds (and emotions connected to them) that we experienced before birth. Percussion instruments remind us of the heart beat of our mother and ourselves, cord and wind instruments remind us of all the voices we heard back then. Musical instruments are especially beautiful if they are driven and modulated by the body of the player. All classical instruments are optimized to allow this modulation and the ability to express emotions in this way.

The connection between the musician and the instrument is most intense for the human voice; the next approximation are the wind instruments. In all these cases, the breath of the singer or player does two things: it provides the energy for the sound and it gives an input for the feedback loop that sets the pitch. While singing, the air passes the vocal cords. The rapid air flow reduces the air pressure, which attracts the cords to each other and thus reduces the cross section for the air flow. (This pressure reduction is described by the Bernoulli equation, as explained below.) As a result of the smaller cross section, the airflow is reduced, the pressure rises again, and the vocal cords open up again. This leads to larger airflow, and the circle starts again. The change between larger and smaller cord distance repeats so rapidly that sound is produced; the sound is then amplified in the mouth by the resonances that depend on the shape of the oral cavity.

In *reed instruments*, such as the clarinet, the reed has the role of the vocal cords, and the pipe and the mechanisms have the role of the mouth. In *brass instruments*, such as the trombone, the lips play the role of the reed. In *airflow instruments*, such as the flute, the feedback loop is due to another effect: at the sound-producing edge, the airflow is deflected by the sound itself.

The second reason that music is beautiful is due to the way the frequencies of the notes are selected. Certain frequencies sound agreeable to the ear when they are played together or closely after each other; other produce a sense of tension. Already the ancient Greek had discovered that these sensations depend exclusively on the *ratio* of the frequencies, or as musician say, on the *interval* between the pitches.

More specifically, a frequency ratio of 2 – musicians call the interval an *octave* – is the most agreeable consonance. A ratio of 3/2 (called perfect fifth) is the next most agreeable, followed by the ratio 4/3 (a perfect fourth), the ratio 5/4 (a major third) and the ratio 6/5 (a [third, minor]minor third). The choice of the first third in a scale has an important effect on the average emotions expressed by the music and is therefore also taken over in the name of the scale. Songs in C major generally have a more happy tune, whereas songs in A minor tend to sound sadder.

The least agreeable frequency ratios, the dissonances, are the *tritone* (7/5, also called *augmented fourth* or *diminished fifth* or *false quint*) and, to a lesser extent, the major and minor seventh (15/8 and 9/5). The tritone is used for the siren in German red cross vans. Long sequences of dissonances have the effect to induce trance; they are common in

Page 302

Ref. 201

TABLE 39 Some signals.

SYSTEM	SIGNAL	SPEED	SENSOR
-	OTGNAL	OFEED	OLNSOR
Matter signals			
Human	voltage pulses in nerves	up to 120 m/s	brain, muscles
	hormones in blood stream	up to 0.3 m/s	molecules on cell membranes
	immune system signals	up to 0.3 m/s	molecules on cell membranes
	singing	340 m/s	ear
Elephant, insects	soil trembling	c. 2 km/s	feet
Whale	singing, sonar	1500 m/s	ear
Dog	chemical tracks	1 m/s	nose
Butterfly	chemical mating signal carried by the wind	up to 10 m/s	antennae
Tree	chemical signal of attack carried by the air from one tree to the next	up to 10 m/s	leaves
Erratic block	carried by glacier	up to 0.1 μm/s	foot
Post	paper letters transported by trucks, ships and planes	up to 300 m/s	mail box
Electromagnetic fields			
Humans	yawning	300 Mm/s	eye
Electric eel	voltage pulse	up to 300 Mm/s	nerves
Insects, fish, molluscs	light pulse sequence	up to 300 Mm/s	eye
Flag signalling	orientation of flags	300 Mm/s	eye
Radio transmissions	electromagnetic field strength	up to 300 Mm/s	radio
Nuclear signals			
Supernovas	neutrino pulses	close to 300 Mm/s	specific chemical and radiation detectors
Nuclear reactions	glueballs, if they exist	close to 300 Mm/s	custom particle detectors

Balinese music and in jazz.

After centuries of experimenting, these results lead to a standardized arrangement of the notes and their frequencies that is shown in Figure 176. The arrangement, called the




FIGURE 176 The twelve notes used in music and their frequency ratios.




FIGURE 177 A schematic animation of a vibrating membrane (drawn in red), from a drum or a loudspeaker, generating a sound wave in air, and showing the molecular motion (Wikicommons)

equal intonation or well-tempered intonation, contains approximations to all the mentioned intervals; the approximations have the advantage that they allow the *transposition* of music to lower or higher notes. This is not possible with the ideal, so-called *just intonation*.

Ref. 202

Challenge 468 e

The next time you sing a song that you like, you might try to determine whether you use just or equal intonation – or a different intonation altogether. Different people have different tastes and habits.

TABLE 40 Selected sound intensities.

OBSERVATION	SOUND INTENSITY
Sound threshold at 1 kHz	0 dB or 1 pW
Human speech	25 to 35 dB
Subway entering a subway station	100 dB
Ultrasound imaging of babies	over 100 dB
Conventional pain threshold	120 dB or 1 W
Rock concert with 400 000 W loudspeakers	135 to 145 dB
Fireworks	up to 150 dB
Gunfire	up to 155 dB
Missile launch	up to 170 dB
Blue whale singing	up to 175 dB
Volcanic eruptions, earthquakes, conventional bomb	up to 210 dB
Large meteoroid impact, large nuclear bomb	over 300 dB





FIGURE 178 A modern ultrasound imaging system, and a common, but harmful ultrasound image of a foetus (© General Electric, Wikimedia).

IS ULTRASOUND IMAGING SAFE FOR BABIES?

Ultrasound is used in medicine to explore the interior of human bodies. The technique, called ultrasound imaging, is helpful, convenient and widespread, as shown in Figure 178. However, it has a disadvantage. Studies at the Mayo Clinic in Minnesota have found that *pulsed* ultrasound, in contrast to *continuous* ultrasound, produces extremely high levels of audible sound inside the body. (Some sound intensities are listed in Table 40.)

Ref. 203

Pulsed ultrasound is used in ultrasound imaging, and in some, but not all, foetal heart-beat monitors. Such machines thus produce high levels of sound in the *audible* range. This seems paradoxical; if you go to a gynaecologist and put the ultrasound head on your ear or head, you will only hear a very faint noise. In fact, it is this low intensity that tricks everybody to think that the noise level is low. The noise level is only low because the human ear is full of air. In contrast, in a foetus, the ear is filled with liquid. This fact changes the propagation of sound completely: the sound generated by imaging machines is now

fully focused and directly stimulates the inner ear. The total effect is similar to what happens if you put your finger in you ear: this can be very loud to yourself, but nobody else can hear what happens.

Ref. 203

Recent research has shown that sound levels of over 100 dB, corresponding to a sub-way train entering the station, are generated by ultrasound imaging systems. Indeed, every gynecologist will confirm that imaging disturbs the foetus. Questioned about this issue, several makers of ultrasound imaging devices confirmed that "a sound output of *only* 5 mW is used". That is 'only' the acoustic power of an oboe at full power! Since many ultrasound examinations take ten minutes and more, a damage to the ear of the foetus cannot be excluded. It is not sensible to expose a baby to this level of noise without good reason

Ref. 204

In short, ultrasound should be used for pregnant mothers only in case of necessity. Ultrasound is *not safe* for the ears of foetuses. In all other situations, ultrasound imaging seems safe. It should be noted however, that another potential problem of ultrasound imaging, the issue of tissue damage through cavitation, has not been explored in detail yet.

Ref. 205

SIGNALS

A signal is the transport of information. Every signal, including those from Table 39, is motion of energy. Signals can be either objects or waves. A thrown stone can be a signal, as can a whistle. Waves are a more practical form of communication because they do not require transport of matter: it is easier to use electricity in a telephone wire to transport a statement than to send a messenger. Indeed, most modern technological advances can be traced to the separation between signal and matter transport. Instead of transporting an orchestra to transmit music, we can send radio signals. Instead of sending paper letters we write email messages. Instead of going to the library we browse the internet.

The greatest advances in communication have resulted from the use of signals to transport large amounts of energy. That is what electric cables do: they transport energy without transporting any (noticeable) matter. We do not need to attach our kitchen machines to the power station: we can get the energy via a copper wire.

For all these reasons, the term 'signal' is often meant to imply waves only. Voice, sound, electric signals, radio and light signals are the most common examples of wave signals.

Signals are characterized by their speed and their information content. Both quantities turn out to be limited. The limit on speed is the central topic of the theory of special relativity.

Vol. II, page 14

A simple limit on information content can be expressed when noting that the information flow is given by the detailed shape of the signal. The shape is characterized by a frequency (or wavelength) and a position in time (or space). For every signal – and every wave – there is a relation between the time-of-arrival error Δt and the angular frequency error $\Delta \omega$:

$$\Delta t \ \Delta \omega \geqslant \frac{1}{2} \ . \tag{98}$$

This time–frequency indeterminacy relation expresses that, in a signal, it is impossible to specify both the time of arrival and the frequency with full precision. The two errors are (within a numerical factor) the inverse of each other. (One also says that the time-

bandwidth product is always larger than $1/4\pi$.) The limitation appears because on the one hand one needs a wave as similar as possible to a sine wave in order to precisely determine the frequency, but on the other hand one needs a signal as narrow as possible to precisely determine its time of arrival. The contrast in the two requirements leads to the limit. The indeterminacy relation is thus a feature of every wave phenomenon. You might want to test this relation with any wave in your environment.

Challenge 469 e

Similarly, there is a relation between the position error Δx and the wave vector error $\Delta k = 2\pi/\Delta \lambda$ of a signal:

$$\Delta x \ \Delta k \geqslant \frac{1}{2} \ . \tag{99}$$

Like the previous case, also this indeterminacy relation expresses that it is impossible to specify both the position of a signal and its wavelength with full precision. Also this position—wave-vector indeterminacy relation is a feature of any wave phenomenon.

Every indeterminacy relation is the consequence of a smallest entity. In the case of waves, the smallest entity of the phenomenon is the period (or cycle, as it used to be called). Whenever there is a smallest unit in a natural phenomenon, an indeterminacy relation results. We will encounter other indeterminacy relations both in relativity and in quantum theory. As we will find out, they are due to smallest entities as well.

Whenever signals are sent, their content can be lost. Each of the six characteristics of waves listed on page 259 can lead to content degradation. Can you provide an example for each case? The energy, the momentum and all other conserved properties of signals are never lost, of course. The disappearance of signals is akin to the disappearance of motion. When motion disappears by friction, it only seems to disappear, and is in fact transformed into heat. Similarly, when a signal disappears, it only seems to disappear, and is in fact transformed into noise. (*Physical*) noise is a collection of numerous disordered

signals, in the same way that heat is a collection of numerous disordered movements.

Ref. 206

Challenge 470 ny

All signal propagation is described by a wave equation. A famous example is the set of equations found by Hodgkin and Huxley. It is a realistic approximation for the behaviour of electrical potential in nerves. Using facts about the behaviour of potassium and sodium ions, they found an elaborate wave equation that describes the voltage V in nerves, and thus the way the signals are propagated. The equation describes the characteristic voltage spikes measured in nerves, shown in Figure 179. The figure clearly shows that these waves differ from sine waves: they are not harmonic. Anharmonicity is one result of nonlinearity. But nonlinearity can lead to even stronger effects.

SOLITARY WAVES AND SOLITONS

In August 1834, the Scottish engineer John Scott Russell (1808–1882) recorded a strange observation in a water canal in the countryside near Edinburgh. When a boat pulled through the channel was suddenly stopped, a strange water wave departed from it. It consisted of a *single* crest, about 10 m long and 0.5 m high, moving at about 4 m/s. He followed that crest, shown in a reconstruction in Figure 180, with his horse for several kilometres: the wave died out only very slowly. Russell did not observe any dispersion, as is usual in deep water waves: the width of the crest remained constant. Russell then started producing such waves in his laboratory, and extensively studied their properties.

Ref. 207




Fig. 13. Upper curve: solution of eqn. (26) for initial depolarization of 15 mV, calculated for 6° C. Lower curve: tracing of membrane action potential recorded at 9·1° C (axon 14). The vertical scales are the same in both curves (apart from curvature in the lower record). The horizontal scales differ by a factor appropriate to the temperature difference.

FIGURE 179 The electrical signals calculated (above) and measured (below) in a nerve, following Hodgkin and Huxley.



FIGURE 180 A solitary water wave followed by a motor boat, reconstructing the discovery by Scott Russel (© Dugald Duncan).

He showed that the speed depended on the amplitude, in contrast to linear, harmonic waves. He also found that the depth d of the water canal was an important parameter. In fact, the speed v, the amplitude A and the width L of these single-crested waves are






FIGURE 181 Solitons are stable against encounters (QuickTime film © Jarmo Hietarinta)

related by

$$v = \sqrt{gd} \left(1 + \frac{A}{2d} \right)$$
 and $L = \sqrt{\frac{4d^3}{3A}}$. (100)

As shown by these expressions, and noted by Russell, high waves are narrow and fast, whereas shallow waves are slow and wide. The shape of the waves is fixed during their motion. Today, these and all other stable waves with a single crest are called *solitary waves*. They appear only where the dispersion and the nonlinearity of the system exactly compensate for each other. Russell also noted that the solitary waves in water channels can cross each other unchanged, even when travelling in opposite directions; solitary waves with this property are called *solitons*. In short, solitons are stable against encounters, as shown in Figure 181, whereas solitary waves in general are not.

Only sixty years later, in 1895, Korteweg and de Vries found out that solitary waves in water channels have a shape described by

$$u(x,t) = A \operatorname{sech}^{2} \frac{x - vt}{L} \quad \text{where} \quad \operatorname{sech} x = \frac{2}{e^{x} + e^{-x}},$$
 (101)

and that the relation found by Russell was due to the wave equation

$$\frac{1}{\sqrt{ad'}}\frac{\partial u}{\partial t} + \left(1 + \frac{3}{2d'}u\right)\frac{\partial u}{\partial x} + \frac{d^2}{6}\frac{\partial^3 u}{\partial x^3} = 0.$$
 (102)

This equation for the elongation u is called the *Korteweg–de Vries equation* in their honour.* The surprising stability of the solitary solutions is due to the opposite effect of the

^{*} The equation can be simplified by transforming the variable u; most concisely, it can be rewritten as $u_t + u_{xxx} = 6uu_x$. As long as the solutions are sech functions, this and other transformed versions of the equation

two terms that distinguish the equation from linear wave equations: for the solitary solutions, the nonlinear term precisely compensates for the dispersion induced by the thirdderivative term.

For many decades such solitary waves were seen as mathematical and physical curiosities. The reason was simple:nobody could solve the equations. All this changed almost a hundred years later, when it became clear that the Korteweg-de Vries equation is a universal model for weakly nonlinear waves in the weak dispersion regime, and thus of basic importance. This conclusion was triggered by Kruskal and Zabusky, who in 1965 proved mathematically that the solutions (101) are unchanged in collisions. This discovery prompted them to introduce the term soliton. These solutions do indeed interpenetrate one another without changing velocity or shape: a collision only produces a small positional shift for each pulse.

Solitary waves play a role in many examples of fluid flows. They are found in ocean currents; and even the red spot on Jupiter, which was a steady feature of Jupiter photographs for many centuries, is an example.

Solitary waves also appear when extremely high-intensity sound is generated in solids. In these cases, they can lead to sound pulses of only a few nanometres in length. Solitary Ref. 211 light pulses are also used inside certain optical communication fibres, where the lack of dispersion allows higher data transmission rates than are achievable with usual light pulses.

Towards the end of the twentieth century, mathematicians discovered that solitons obey a nonlinear generalization of the superposition principle. (It is due to the Darboux-Backlund transformations and the structure of the Sato Grassmannian.) The mathematics of solitons is extremely interesting. The progress in mathematics triggered a second wave of interest in the mathematics of solitons arose, when quantum theorists became interested in them. The reason is simple: a soliton is a 'middle thing' between a particle and a wave; it has features of both concepts. For this reason, solitons were often seen incorrectly though – as candidates for the description of elementary particles.

CURIOSITIES AND FUN CHALLENGES ABOUT WAVES AND EXTENDED BODIES

Society is a wave. The wave moves onward, but the water of which it is composed does not. Ralph Waldo Emerson, Self-Reliance.

When the frequency of a tone is doubled, one says that the tone is higher by an octave. Two tones that differ by an octave, when played together, sound pleasant to the ear. Two other agreeable frequency ratios – or 'intervals', as musicians say – are quarts and quints. What are the corresponding frequency ratios? (Note: the answer was one of the oldest discoveries in physics and perception research; it is attributed to Pythagoras, around 500 BCE.)

* *

Also the bumps of skiing slopes, the so-called ski moguls, are waves. Ski moguls are essential in many winter Olympic disciplines. Observation shows that ski moguls have a Ref. 209

are known by the same name.

Challenge 471 e

Ref. 210

Ref. 207

Ref. 208

Challenge 472 s Page 348

Challenge 473 s

Challenge 476 e

Challenge 477 e

wavelength of typically 5 to 6 m and that they move with an average speed of 8 cm/day. Surprisingly, the speed is directed *upwards*, towards the top of the skiing slope. Can you explain why this is so? In fact, ski moguls are also an example of self-organization; this topic will be covered in more detail below.

* *

An orchestra is playing music in a large hall. At a distance of 30 m, somebody is listening to the music. At a distance of 3000 km, another person is listening to the music via the radio. Who hears the music first?

* *

What is the period of a simple pendulum, i.e., a mass m attached to a massless string of length l? What is the period if the string is much longer than the radius of the Earth?

* *

What path is followed by a body that moves without friction on a plane, but that is at-Challenge 475 s tached by a spring to a fixed point on the plane?

* *

The blue whale, *Balaenoptera musculus*, is the *loudest* animal found in nature: its voice can be heard at a distance of hundreds of kilometres.

* *

The exploraation of sound in the sea, from the communication of whales to the sonar of dolphins, is a world of its own. As a start, explore the excellent www.dosits.org website.

* *

A device that shows how rotation and oscillation are linked is the alarm *siren*. Find out how it works, and build one yourself.

* *

Light is a wave, as we will discover later on. As a result, light reaching the Earth from space is refracted when it enters the atmosphere. Can you confirm that as a result, stars appear somewhat higher in the night sky than they really are?

* *

What are the highest sea waves? This question has been researched systematically only recently, using satellites. The surprising result is that sea waves with a height of 25 m and more are *common*: there are a few such waves on the oceans at any given time. This result confirms the rare stories of experienced ship captains and explains many otherwise unexplained ship sinkings.

Surfers may thus get many chances to ride 30 m waves. (The present record is just below this height.) But maybe the most impressive waves to surf are those of the Pororoca, a series of 4 m waves that move from the sea into the Amazon River every spring, against the flow of the river. These waves can be surfed for tens of kilometres.




FIGURE 182 Shadows show the refraction of light.

Interestingly, every water surface has waves, even if it seems completely flat. As a consequence of the finite temperature of water, its surface always has some roughness: there are thermal capillary waves. For water, with a surface tension of 72 mPa, the typical roughness at usual conditions is 0.2 nm. These thermal capillary waves, predicted since many

centuries, have been observed only recently. Ref. 213

All waves are damped, eventually. This effect is often frequency-dependent. Can you provide a confirmation of this dependence in the case of sound in air? Challenge 478 s

Challenge 479 e

When you make a hole with a needle in black paper, the hole can be used as a magnifying lens. (Try it.) Diffraction is responsible for the lens effect. By the way, the diffraction of light by holes was noted already by Francesco Grimaldi in the seventeenth century; he correctly deduced that light is a wave. His observations were later discussed by Newton, who wrongly dismissed them.

Put an empty cup near a lamp, in such a way that the bottom of the cup remains in the shadow. When you fill the cup with water, some of the bottom will be lit, because of the refraction of the light from the lamp. The same effect allows us to build lenses. The same effect is at the basis of instruments such as the telescope.

Vol. III, page 136

Are water waves transverse or longitudinal? Challenge 480 s

The speed of water waves limits the speeds of ships. A surface ship cannot travel (much) faster than about $v_{\text{crit}} = \sqrt{0.16gl}$, where $g = 9.8 \,\text{m/s}^2$, l is the boat's length, and 0.16 is a number determined experimentally, called the critical Froude number. This relation is valid for all vessels, from large tankers (l = 100 m gives $v_{crit} = 13 \text{ m/s}$) down to ducks (l = 100 m gives $v_{crit} = 13 \text{ m/s}$) 0.3 m gives $v_{\rm crit} = 0.7$ m/s). The critical speed is that of a wave with the same wavelength as the ship. In fact, moving a ship at higher speeds than the critical value is possible, but requires much more energy. (A higher speed is also possible if the ship surfs on a wave.) How far away is the crawl *olympic swimming* record from the critical value?

Challenge 481 s

Most water animals and ships are faster when they swim below the surface - where

the limit due to surface waves does not exist – than when they swim on the surface. For example, ducks can swim three times as fast under water than on the surface.

* *

The group velocity of water waves (in deep water) is less than the velocity of the individual wave crests, the so-called *phase velocity*. As a result, when a group of wave crests travels, within the group the crests move from the back to the front: they appear at the back, travel forward and then die out at the front. The group velocity of water waves is lower than its phase velocity.

* *

One can hear the distant sea or a distant highway more clearly in the evening than in the morning. This is an effect of refraction. Sound speed increases with temperature. In the evening, the ground cools more quickly than the air above. As a result, sound leaving the ground and travelling upwards is refracted downwards, leading to the long hearing distance typical of evenings. In the morning, usually the air is cold above and warm below. Sound is refracted upwards, and distant sound does not reach a listener on the ground. Refraction thus implies that mornings are quiet, and that we can hear more distant sounds in the evenings. Elephants use the sound situation during evenings to communicate over distances of more than 10 km. (They also use sound waves in the ground to communicate, but that is another story.)

* *

Refraction also implies that there is a sound channel in the ocean, and in the atmosphere. Sound speed increases with temperature, and increases with pressure. At an ocean depth of 1 km, or at an atmospheric height of 13 to 17 km (that is at the top of the tallest cumulonimbus clouds or equivalently, at the middle of the ozone layer) sound has *minimal* speed. As a result, sound that starts from that level and tries to leave is channelled back to it. Whales use this *sound channel* to communicate with each other with beautiful songs; one can find recordings of these songs on the internet. The military successfully uses microphones placed at the sound channel in the ocean to locate submarines, and microphones on balloons in the atmospheric channel to listen for nuclear explosions. (In fact, sound experiments conducted by the military are the main reason why whales are deafened and lose their orientation, stranding on the shores. Similar experiments in the air with high-altitude balloons are often mistaken for flying saucers, as in the famous Roswell incident.)

* :

Ref. 215 Also small animals communicate by sound waves. In 2003, it was found that herring communicate using noises they produce when farting. When they pass wind, the gas creates a ticking sound whose frequency spectrum reaches up to 20 kHz. One can even listen to recordings of this sound on the internet. The details of the communication, such as the differences between males and females, are still being investigated. It is possible that the sounds may also be used by predators to detect herring, and they might even be used by future fishing vessels.

Challenge 482 e

Ref. 214




FIGURE 183 Rubik's cube: the complexity of simple three-dimensional motion (© Wikimedia).

* *

On windy seas, the white wave crests have several important effects. The noise stems from tiny exploding and imploding water bubbles. The noise of waves on the open sea is thus the superposition of many small explosions. At the same time, white crests are the events where the seas absorb carbon dioxide from the atmosphere, and thus reduce global warming.

* *

Challenge 483 s Why are there many small holes in the ceilings of many office rooms?

* *

Which quantity determines the wavelength of water waves emitted when a stone is thrown into a pond?

Challenge 484 ny

Ref. 2

* *

- Yakov Perelman lists the following four problems in his delightful physics problem book.
- (1) A stone falling into a lake produces circular waves. What is the shape of waves produced by a stone falling into a river, where the water flows?
- (2) It is possible to build a lens for sound, in the same way as it is possible to build lenses for light. What would such a lens look like?
 - (3) What is the sound heard inside a shell?
- (4) Light takes about eight minutes to travel from the Sun to the Earth. What consequence does this have for the timing of sunrise?

Challenge 488 s

Challenge 485 s

Challenge 486 s

Challenge 487 s




FIGURE 184 The human ear (© Northwestern University).

* *

Every student probably knows *Rubik's cube*. Can you or did you deduce how Rubik built the cube without looking at its interior? Also for cubes with other numbers of segments? Is there a limit to the number of segments? These puzzles are even tougher than the search for the rearrangement of the cube.

* *

Typically, sound of a talking person produces a pressure variation of 20 mPa on the ear.

Challenge 490 ny

How is this value determined?

The ear is indeed a sensitive device. It is now known that most cases of sea mammals, like whales, swimming onto the shore are due to ear problems: usually some military device (either sonar signals or explosions) has destroyed their ear so that they became deaf and lose orientation.

* *

Why is the human ear, shown in Figure 184, so complex? The outer part, the *pinna* or *auricola*, concentrates the sound pressure at the tympanic membrane; it produces a gain of 3 dB. The tympanic membrane, or eardrum, is made in such a way as to always oscillate in fundamental mode, thus without any nodes. The tympanic membrane has a (very wide) resonance at 3 kHz, in the region where the ear is most sensitive. The eardrum transmits its motion, using the ossicles, into the inner ear. This mechanism thus transforms air waves into water waves in the inner ear, where they are detected. (The efficiency with which this transformation takes place is almost ideal; using the language of wave theory, ossicles are thus above all impedance transformers.) Why does the ear transform

Challenge 489 s

air waves to water waves? Because the sound wavelength in liquids is much shorter, and allows a small detector; otherwise, the ear would not fit inside the head!

* *

Infrasound, inaudible sound below 20 Hz, is a modern topic of research. In nature, infrasound is emitted by earthquakes, volcanic eruptions, wind, thunder, waterfalls, falling meteorites and the surf. Glacier motion, seaquakes, avalanches and geomagnetic storms also emit infrasound. Human sources include missile launches, traffic, fuel engines and air compressors.

It is known that high intensities of infrasound lead to vomiting or disturbances of the sense of equilibrium (140 dB or more for 2 minutes), and even to death (170 dB for 10 minutes). The effects of lower intensities on human health are not yet known.

Infrasound can travel several times around the world before dying down, as the explosion of the Krakatoa volcano showed in 1883. With modern infrasound detectors, sea surf can be detected hundreds of kilometres away. Sea surf leads to a constant 'hum' of the Earth's crust at frequencies between 3 and 7 mHz. The *Global infrasound Network* uses infrasound to detect nuclear weapon tests, earthquakes and volcanic eruptions, and can count meteorites. Only very rarely can meteorites be heard with the human ear.

* *

The method used to deduce the sine waves contained in a signal, as shown in Figure 166, is called the Fourier transformation. It is of importance throughout science and technology. In the 1980s, an interesting generalization became popular, called the *wavelet transformation*. In contrast to Fourier transformations, wavelet transformations allow us to localize signals in time. Wavelet transformations are used to compress digitally stored images in an efficient way, to diagnose aeroplane turbine problems, and in many other applications.

Ref. 217

Challenge 491 r

* *

If you like engineering challenges, here is one that is still open. How can one make a robust and efficient system that transforms the energy of sea waves into electricity?

* *

If you are interested in ocean waves, you might also enjoy the science of *oceanography*. For an introduction, see the open source textbooks at oceanworld.tamu.edu.

* *

In our description of extended bodies, we assumed that each spot of a body can be followed separately throughout its motion. Is this assumption justified? What would happen if it were not?

Challenge 492 r

* *

A special type of waves appears in explosions and supersonic flight: *shock waves*. In a shock wave, the density or pressure of a gas changes abruptly, on distances of a few micrometers. Studying shock waves is a research field in itself; shock waves determine the flight of bullets, the snapping of whips and the effects of detonations.

sound

waves

moving shock wave: moving 'sonic boom'


sound source

medium at supersonic speed

moving through

supersonic sound source

moving on ground



Around a body moving with supersonic speed, the sound waves form a cone, as shown in Figure 185. When the cone passes an observer on the ground, the cone leads to a *sonic boom*. What is less well known is that the boom can be amplified. If an aeroplane *accelerates* through the sound barrier, certain observers at the ground will hear two booms or even a so-called superboom, because cones from various speeds can superpose at certain spots on the ground. A plane that performs certain manoeuvres, such as a curve at high speed, can even produce a superboom at a predefined spot on the ground. In contrast to normal sonic booms, superbooms can destroy windows, eardrums and lead to trauma, especially in children. Unfortunately, they are regularly produced on purpose by frustrated military pilots in various places of the world.

* *

What have swimming swans and ships have in common? The *wake* behind them. Despite the similarity, this phenomenon has no relation to the sonic boom. In fact, the angle of the wake is the *same* for ducks and ships, and is *independent* of the speed they travel or of the size of the moving body, provided the water is deep enough.

As explained above, water waves in *deep* water differ from sound waves: their group velocity is *one half* the phase velocity. (Can you deduce this from the dispersion relation

Ref. 2

Page 254

pdf file available

FIGURE 186 The wakes behind a ship and behind a swan, and the way to deduce the shape (photos © Wikimedia, Christopher Thorn).

Challenge 493 e

 $\omega = \sqrt{gk}$ between angular frequency and wave vector, valid for deep water waves?) Water waves will interfere where most of the energy is transported, thus around the group velocity. For this reason, in the graph shown in Figure 186, the diameter of each wave circle is always half the distance of their leftmost point O to the apex A. As a result, the half angle of the wake apex obeys

$$\sin \alpha = \frac{1}{3}$$
 giving a wake angle $2\alpha = 38.942^{\circ}$. (103)

Ref. 219

Figure 186 also allows deducing the curves that make up the wave pattern of the wake, using simple geometry.

It is essential to note that the fixed wake angle is valid only in deep water, i.e., only in water that is much deeper than the wavelength of the involved waves. In other words, for a given depth, the wake has the fixed shape only up to a maximum source speed. For high speeds, the wake angle narrows, and the pattern inside the wake changes.




FIGURE 187 The calculated motion of a dromion across a two-dimensional substrate (QuickTime film © Jarmo Hietarinta)

* *

Bats fly at night using *echolocation*. Dolphins also use it. Sonar, used by fishing vessels to look for fish, copies the system of dolphins. Less well known is that humans have the same ability. Have you ever tried to echolocate a wall in a completely dark room? You will be surprised at how easily this is possible. Just make a loud hissing or whistling noise that stops abruptly, and listen to the echo. You will be able to locate walls reliably.

Challenge 494 e

Ref. 220

Birds sing. If you want to explore how this happens, look at the impressive X-ray film

found at the www.indiana.edu/~songbird/multi/cineradiography_index.html website.

* *

Every soliton is a one-dimensional structure. Do two-dimensional analogues exist? This issue was open for many years. Finally, in 1988, Boiti, Leon, Martina and Pempinelli found that a certain evolution equation, the so-called Davey–Stewartson equation, can have solutions that are localized in two dimensions. These results were generalized by Fokas and Santini and further generalized by Hietarinta and Hirota. Such a solution is today called a *dromion*. Dromions are bumps that are localized in two dimensions and can move, without disappearing through diffusion, in non-linear systems. An example is shown in Figure 187. However, so far, no such solution has be observed in experiments; this is one of the most important experimental challenges left open in non-linear science.

* *

Water waves have not lost their interest to this day. Most of all, two-dimensional solutions of solitonic water waves remain a topic of research. The mathematics is complicated, the experiments simple, and the issue is fascinating. In two dimensions, soliton crests can form hexagonal patterns! The relevant equation for shallow waves, the generalization of

the Korteweg-de Vries equation to two dimensions, is called the *Kadomtsev-Petviashvili* equation. For pictures of such extremely rare waves, see the site by Bernard Deconinck at www.amath.washington.edu/~bernard/kp/waterwaves.html. The issue of whether rectangular patterns exist is still open, as are the equations and solutions for deep water waves. Water waves have not yet yielded all their secrets.

* *

How does the tone produced by blowing over a bottle depend on the dimension? For bottles that are bulky, the frequency f, the so-called *cavity resonance*, is found to depend on the volume V of the bottle:

ChallengeRef 5222

$$f = \frac{c}{2\pi} \sqrt{\frac{A}{VL}}$$
 or $f \sim \frac{1}{\sqrt{V}}$ (104)

Challenge 496 e

Ref. 223

where c is the speed of sound, A is the area of the opening, and L is the length of the neck of the bottle. Does the formula agree with your observations?

In fact, tone production is a complicated issue, and specialized books exist on the topic. For example, when overblowing, a saxophone produces a second harmonic, an octave, whereas a clarinet produces a third harmonic, a quint (more precisely, a twelfth). Why is this the case? The theory is complex, but the result simple: instruments whose cross-section increases along the tube, such as horns, trumpets, oboes or saxophones, overblow to octaves. For air instruments that have a (mostly) cylindrical tube, the effect of overblowing depends on the tone generation mechanism. Flutes overblow to the

octave, but clarinets to the twelfth.

* *

Many acoustical systems do not only produce harmonics, but also subharmonics. There is a simple way to observe production of subharmonics: sing with your ears below water, in the bathtub. Depending on the air left in your ears, you can hear subharmonics of your own voice. The effect is quite special.

* *

The origin of the sound of cracking joints, for example in the hand, is a well-known puzzle. How would you test the conjecture that it is due to cavitation? What would you do to find out definitively?

Challenge 497 e

* *

Among the most impressive sound experiences are the singing performances of *countertenors* and of the even higher singing *male sopranos*. If you ever have the chance to hear one, do not miss the occasion.

SUMMARY ON WAVES AND OSCILLATIONS

In nature, apart from the motion of bodies, we observe also the motion of waves and wave groups, or signals. Waves have energy, momentum and angular momentum. Waves can interfere, diffract, refract, disperse, dampen out and, if transverse, can be polarized.

Oscillations are a special case of waves; usually they are standing waves. Solitary waves, i.e., waves with only one crest, are a special case of waves.



CHAPTER 11

DO EXTENDED BODIES EXIST? - LIMITS OF CONTINUITY

E have just discussed the motion of bodies that are extended. e have found that all extended bodies, be they solid of fluid, show ave motion. But are extended bodies actually found in nature? Strangely enough, this question has been one of the most intensely discussed questions in physics. Over the centuries, it has reappeared again and again, at each improvement of the description of motion; the answer has alternated between the affirmative and the negative. Many thinkers have been imprisoned, and many still are being persecuted, for giving answers that are not politically correct! In fact, the issue already arises in everyday life.

MOUNTAINS AND FRACTALS

Whenever we climb a mountain, we follow the outline of its shape. We usually describe this outline as a curved two-dimensional surface. But is this correct? There are alternative possibilities. The most popular is the idea that mountains are fractal surfaces. A *fractal* was defined by Benoît Mandelbrot as a set that is self-similar under a countable but infinite number of magnification values. We have already encountered fractal lines. An example of an algorithm for building a (random) fractal *surface* is shown on the right side of Figure 188. It produces shapes which look remarkably similar to real mountains. The results are so realistic that they are used in Hollywood films. If this description were correct, mountains would be extended, but not continuous.

But mountains could also be fractals of a different sort, as shown in the left side of Figure 188. Mountain surfaces could have an infinity of small and smaller holes. In fact, we could also imagine that mountains are described as three-dimensional versions of the left side of the figure. Mountains would then be some sort of mathematical Swiss cheese. Can you devise an experiment to decide whether fractals provide the correct description for mountains? To settle the issue, we study chocolate bars, bones, trees and soap bubbles.

Challenge 498 s

Page 54

Ref. 224

CAN A CHOCOLATE BAR LAST FOREVER?

From a drop of water a logician could predict an Atlantic or a Niagara. Arthur Conan Doyle, A Study in Scarlet

Any child knows how to make a chocolate bar last forever: eat half the remainder every day. However, this method only works if matter is scale-invariant. In other words, the method only works if matter is either *fractal*, as it then would be scale-invariant for a

284 11 EXTENDED BODIES




FIGURE 188 Floors (left) and mountains (right) could be fractals; for mountains this approximation is often used in computer graphics (image © Paul Martz).

discrete set of zoom factors, or *continuous*, in which case it would be scale-invariant for any zoom factor. Which case, if either, applies to nature?

Page 53

We have already encountered a fact making continuity a questionable assumption: continuity would allow us, as Banach and Tarski showed, to multiply food and any other matter by clever cutting and reassembling. Continuity would allow children to eat the *same* amount of chocolate every day, without ever buying a new bar. Matter is thus not continuous. Now, fractal chocolate is not ruled out in this way; but other experiments settle the question. Indeed, we note that melted materials do not take up much smaller volumes than solid ones. We also find that even under the highest pressures, materials do not shrink. Thus we conclude again that matter is not a fractal. What then is its structure?

To get an idea of the structure of matter we can take fluid chocolate, or even just some oil – which is the main ingredient of chocolate anyway – and spread it out over a large surface. For example, we can spread a drop of oil onto a pond on a day without rain or wind; it is not difficult to observe which parts of the water are covered by the oil and which are not. A small droplet of oil cannot cover a surface larger than – can you guess the value? The oil-covered water and the uncovered water have different colours. Trying to spread the oil film further inevitably *rips it apart*. The child's method of prolonging chocolate thus does not work for ever: it comes to a sudden end. The oil experiment, which can even be conducted at home, shows that there is a *minimum* thickness of oil films, with a value of about 2 nm. The experiment shows* that there is a smallest size in oil. Oil is made of tiny components. Is this valid for *all* matter?

Challenge 499 s

^{*} The oil experiment was popularized a few decades after Loschmidt's determination of the size of molecules, by Thomson-Kelvin. It is often claimed that Benjamin Franklin was the first to conduct the oil experiment; that is wrong. Franklin did not measure the thickness, and did not even consider the question of the thickness. He did pour oil on water, but missed the most important conclusion that could be drawn from it. Even geniuses do not discover everything.

THE CASE OF GALILEO GALILEI

After the middle ages, Galileo (1564–1642) was the first to state that *all* matter was made of smallest parts, which he called *piccolissimi quanti*, i.e., smallest quanta. Today, they are called *atoms*. However, Galileo paid dearly for this statement.

Indeed, during his life, Galileo was under attack for two reasons: because of his ideas on the motion of the Earth, and because of his ideas about atoms.* The discovery of the importance of both issues is the merit of the great historian Pietro Redondi, a collaborator of another great historian, Pierre Costabel. One of Redondi's research topics is the history of the dispute between the Jesuits, who at the time defended orthodox theology, and Galileo and the other scientists. In the 1980s, Redondi discovered a document of that time, an anonymous denunciation called G3, that allowed him to show that the condemnation of Galileo to life imprisonment for his views on the Earth's motion was organized by his friend the Pope to *protect* him from a sure condemnation to death over a different issue: atoms.

Page 287

Galileo defended the view, explained in detail shortly, that since matter is not scale invariant, it must be made of 'atoms' or, as he called them, *piccolissimi quanti*. This was and still is a heresy, because atoms of matter contradict the central Catholic idea that in the Eucharist the *sensible qualities* of bread and wine exist independently of their *substance*. The distinction between substance and sensible qualities, introduced by Thomas Aquinas, is essential to make sense of *transubstantiation*, the change of bread and wine into human flesh and blood, which is a central tenet of the Catholic faith. Indeed, a true Catholic is still not allowed to believe in atoms to the present day, because the idea that matter is made of atoms contradicts transubstantiation. (Protestant faith usually does not support transubstantiation, by the way.)

Ref. 225

In Galileo's days, church tribunals punished heresy, i.e., personal opinions deviating from orthodox theology, by the death sentence. But Galileo was not sentenced to death. Galileo's life was saved by the Pope by making sure that the issue of transubstantiation would not be a topic of the trial, and by ensuring that the trial at the Inquisition be organized by a papal commission led by his nephew, Francesco Barberini. But the Pope also wanted Galileo to be punished, because he felt that his own ideas had been mocked in Galileo's book *Il Dialogo* and also because, under attack for his foreign policy, he was not able to ignore or suppress the issue.

ef. 226

As a result, in 1633 the seventy-year-old Galileo was condemned to a prison sentence, 'after invoking the name of Jesus Christ', for 'suspicion of heresy' (and thus not for heresy), because he did not comply with an earlier promise not to teach that the Earth moves. Indeed, the motion of the Earth contradicts what the Christian bible states. Galileo was convinced that *truth* was determined by observation, the Inquisition that it was determined by a book – and by itself. In many letters that Galileo wrote throughout his life he expressed his conviction that observational truth could never be a heresy. The trial showed him the opposite: he was forced to state that he erred in teaching that the Earth moves. After a while, the Pope reduced the prison sentence to house arrest.

Ref. 227

^{*} To get a clear view of the matters of dispute in the case of Galileo, especially those of interest to physicists, the best text is the excellent book by Pietro Redondi, *Galileo eretico*, Einaudi, 1983, translated into English as *Galileo Heretic*, Princeton University Press, 1987. It is also available in many other languages; an updated edition that includes the newest discoveries appeared in 2004.

286 11 EXTENDED BODIES

Galileo's condemnation on the motion of the Earth was not the end of the story. In the years after Galileo's death, also atomism was condemned in several trials against Galileo's ideas and his followers. But the effects of these trials were not those planned by the Inquisition. Only twenty years after the famous trial, around 1650, every astronomer in the world was convinced of the motion of the Earth. And the result of the trials against atomism was that at the end of the 17th century, practically every scientist in the world was convinced that atoms exist. The trials accelerated an additional effect: after Galileo and Descartes, the centre of scientific research and innovation shifted from Catholic countries, like Italy or France, to protestant countries. In these, such as the Netherlands, England, Germany or the Scandinavian countries, the Inquisition had no power. This shift is still felt today.

It is a sad story that in 1992, the Catholic church did *not* revoke Galileo's condemnation. In that year, Pope John Paul II gave a speech on the Galileo case. Many years before, he had asked a study commission to re-evaluate the trial, because he wanted to express his regrets for what had happened and wanted to rehabilitate Galileo. The commission worked for twelve years. But the bishop that presented the final report was a crook: he avoided citing the results of the study commission, falsely stated the position of both parties on the subject of truth, falsely stated that Galileo's arguments on the motion of the Earth were weaker than those of the church, falsely summarized the past positions of the church on the motion of the Earth, avoided stating that prison sentences are not good arguments in issues of opinion or of heresy, made sure that rehabilitation was not even discussed, and of course, avoided any mention of transubstantiation. At the end of this power struggle, Galileo was thus *not* rehabilitated, in contrast to what the Pope wanted and in contrast to what most press releases of the time said; the Pope only stated that 'errors were made on both sides', and the crook behind all this was rewarded with a promotion.*

But that is not the end of the story. The documents of the trial, which were kept locked when Redondi made his discovery, were later made accessible to scholars by Pope John Paul II. In 1999, this led to the discovery of a new document, called EE 291, an internal expert opinion on the atom issue that was written for the trial in 1632, a few months before the start of the procedure. The author of the document comes to the conclusion that Galileo was indeed a heretic in the matter of atoms. The document thus proves that the cover-up of the transubstantiation issue during the trial of Galileo must have been systematic and thorough, as Redondi had deduced. Indeed, church officials and the Catholic catechism carefully avoid the subject of atoms even today; you can search the Vatican website www.vatican.va for any mention of them.

But Galileo did not want to attack transubstantiation; he wanted to advance the idea of atoms. And he did. Despite being condemned to prison in his trial, Galileo's last book, the *Discorsi*, written as a blind old man under house arrest, includes the discussion of atoms, or *piccolissimi quanti*. It is an irony of history that today, quantum theory, named by Max Born after the term used by Galileo for atoms, has become the most precise description of nature yet. Let us explore how Galileo concluded that all matter is made of atoms.

Vol. IV, page 13

D (22)

Ref. 229

^{*} We should not be too indignant: the same situation happens in many commercial companies every day; most industrial employees can tell similar stories.

HOW HIGH CAN ANIMALS JUMP?

Fleas can jump to heights a hundred times their size, humans only to heights about their own size. In fact, biological studies yield a simple observation: most animals, regardless of their size, achieve about the same jumping height, namely between 0.8 and 2.2 m, whether they are humans, cats, grasshoppers, apes, horses or leopards. We have explained this fact earlier on.

At first sight, the observation of constant jumping height seems to be a simple example of scale invariance. But let us look more closely. There are some interesting exceptions at both ends of the mass range. At the small end, mites and other small insects do not achieve such heights because, like all small objects, they encounter the problem of air resistance. At the large end, elephants do not jump that high, because doing so would break their bones. But why do bones break at all?

Why are all humans of about the same size? Why are there no giant adults with a height of ten metres? Why aren't there any land animals larger than elephants? Galileo already gave the answer. The bones of which people and animals are made would not allow such changes of scale, as the bones of giants would collapse under the weight they have to sustain. (A human scaled up by a factor 10 would weigh 1000 times as much, but its bones would only be 100 times as wide.) But why do bones have a finite strength at all? There is only one explanation: because the constituents of bones stick to each other with a finite attraction. In contrast to bones, *continuous* matter – which exists only in cartoons – could not break at all, and *fractal* matter would be infinitely fragile. Galileo concluded that matter breaks under finite loads because it is composed of small basic constituents.

FELLING TREES

The gentle lower slopes of Motion Mountain are covered by trees. Trees are fascinating structures. Take their size. Why do trees have limited size? Already in the sixteenth century, Galileo knew that it is not possible to increase tree height without limits: at some point a tree would not have the strength to support its own weight. He estimated the maximum height to be around 90 m; the actual record, unknown to him at the time, seems to be 150 m, for the Australian tree *Eucalyptus regnans*. But why does a limit exist at all? The answer is the same as for bones: wood has a finite strength because it is not scale invariant; and it is not scale invariant because it is made of small constituents, namely *atoms*.*

In fact, the derivation of the precise value of the height limit is more involved. Trees must not break under strong winds. Wind resistance limits the height-to-thickness ratio h/d to about 50 for normal-sized trees (for $0.2\,\mathrm{m} < d < 2\,\mathrm{m}$). Can you say why? Thinner trees are limited in height to less than 10 m by the requirement that they return to the vertical after being bent by the wind.

Such studies of natural constraints also answer the question of why trees are made from wood and not, for example, from steel. You could check for yourself that the maximum height of a column of a given mass is determined by the ratio E/ρ^2 between the elastic module and the square of the mass density. For a long time, wood was actually the

Challenge 500 ny

Ref. 232

Challenge 501 s

Ref. 231 * There is another important limiting factor: the water columns inside trees must not break. Both factors seem to yield similar limiting heights.

Ref. 233 material for which this ratio was highest. Only recently have material scientists managed to engineer slightly better ratios: the fibre composites.

Why do materials break at all? All observations yield the same answer and confirm Galileo's reasoning: because there is a smallest size in materials. For example, bodies under stress are torn apart at the position at which their strength is minimal. If a body were completely homogeneous or continuous, it could not be torn apart; a crack could not start anywhere. If a body had a fractal, Swiss-cheese structure, cracks would have places to start, but they would need only an infinitesimal shock to do so.

LITTLE HARD BALLS

I prefer knowing the cause of a single thing to being king of Persia.

Democritus

Precise observations show that matter is neither continuous nor a fractal: all matter is made of smallest basic particles. Galileo, who deduced their existence by thinking about giants and trees, called them *smallest quanta*. Today they are called *atoms*, in honour of a famous argument of the ancient Greeks. Indeed, 2500 years ago, the Greeks asked the following question. If motion and matter are conserved, how can change and transformation exist? The philosophical school of Leucippus and Democritus of Abdera* studied two particular observations in special detail. They noted that *salt dissolves in water*. They also noted that *fish can swim in water*. In the first case, the volume of water does not increase when the salt is dissolved. In the second case, when fish advance, they must push water aside. Leucippus and Democritus deduced that there is only one possible explanation that satisfies these two observations and also reconciles conservation with transformation: nature is made of void and of small, indivisible and conserved particles.**

In short, since matter is hard, has a shape and is divisible, Leucippus and Democritus imagined it as being *made of atoms*. Atoms are particles which are hard, have a shape, but are indivisible. The Greek thus deduced that *every* example of motion, change and

Especially if we imagine particles as little balls, we cannot avoid calling this a typically male idea. (What would be the female approach?)

Challenge 502 ny

^{*} Leucippus of Elea (Λευκιππος) (c. 490 to c. 430 bce), Greek philosopher; Elea was a small town south of Naples. It lies in Italy, but used to belong to the Magna Graecia. Democritus (Δεμοκριτος) of Abdera (c. 460 to c. 356 or 370 bce), also a Greek philosopher, was arguably the greatest philosopher who ever lived. Together with his teacher Leucippus, he was the founder of the atomic theory; Democritus was a much admired thinker, and a contemporary of Socrates. The vain Plato never even mentions him, as Democritus was a danger to his own fame. Democritus wrote many books which all have been lost; they were not copied during the Middle Ages because of his scientific and rational world view, which was felt to be a danger by religious zealots who had the monopoly on the copying industry. Nowadays, it has become common to claim – incorrectly – that Democritus had no proof for the existence of atoms. That is a typical example of disinformation with the aim of making us feel superior to the ancients.

^{**} The story is told by Lucretius, in full Titus Lucretius Carus, in his famous text *De rerum natura*, around 60 BCE. (An English translation can be found on www.perseus.tufts.edu/hopper/text?doc=Lucr.+1.1.) Lucretius relates many other proofs; in Book 1, he shows that there is vacuum in solids – as proven by porosity and by density differences – and in gases – as proven by wind. He shows that smells are due to particles, and that so is evaporation. (Can you find more proofs?) He also explains that the particles cannot be seen due to their small size, but that their effects can be felt and that they allow explaining all observations consistently.

transformation is due to rearrangements of these particles; change and conservation are thus reconciled.

In other words, the Greeks imagined nature as a big Lego set. Lego pieces are first of all hard or *impenetrable*, i.e., repulsive at very small distances. Atoms thus explain why solids cannot be compressed much. Lego pieces are also *attractive* at small distances: they remain stuck together. Atoms this explain that solids exist. Finally, lego bricks have *no interaction* at large distances. Atoms thus explain the existence of gases. (Actually, what the Greeks called 'atoms' partly corresponds to what today we call 'molecules'. The latter term was introduced in 1811 by Amedeo Avogadro* in order to clarify the distinction. But we can forget this detail for the moment.)

Since atoms are invisible, it took many years before all scientists were convinced by the experiments showing their existence. In the nineteenth century, the idea of atoms was beautifully verified by a large number of experiments, such as the discovery of the 'laws' of chemistry and those of gas behaviour. We briefly explore the most interesting ones

THE SOUND OF SILENCE

Climbing the slopes of Motion Mountain, we arrive in a region of the forest covered with deep snow. We stop for a minute and look around. It is already dark; all the animals are asleep; there is no wind and there are no sources of sound. We stand still, without breathing, and listen to the silence. (You can have the same experience also in a sound studio such as those used for musical recordings, or in a quiet bedroom at night.) In situations of complete silence, the ear automatically becomes more sensitive**; we then have a strange experience. We hear two noises, a lower- and a higher-pitched one, which are obviously generated inside the ear. Experiments show that the higher note is due to the activity of the nerve cells in the inner ear. The lower note is due to pulsating blood streaming through the head. But why do we hear a noise at all?

Many similar experiments confirm that whatever we do, we can never eliminate noise, i.e., random fluctuations, from measurements. This unavoidable type of random fluctuations is called *shot noise* in physics. The statistical properties of this type of noise actually correspond precisely to what would be expected if flows, instead of being motions of continuous matter, were transportation of a large number of equal, small and discrete entities. Therefore, the precise measurement of noise can be used to prove that air and liquids are made of molecules, that electric current is made of electrons, and that light is made of photons. In a sense, the sound of silence is the sound of atoms. Shot noise would not exist if matter were continuous or fractal.

How to count what cannot be seen

Ref. 234

In everyday life, atoms cannot be counted, mainly because they cannot be seen. Interestingly, the progress of physics allowed to count atoms nevertheless. As mentioned, many

^{*} Amedeo Avogadro (b. 1776 Turin, d. 1856 Turin) Piedmontese physicist and chemist. Avogadro's number is named for him.

^{**} The human ear can detect, in its most sensitive mode, pressure variations at least as small as $20\,\mu\text{Pa}$ and ear drum motions as small as $11\,\text{pm}$.

of these methods use the measurement of noise.*

Page 326

In physics, the term *noise* is not only used for the acoustical effect; it is used for *any* process that is random. The most famous kind of noise is Brownian motion, the motion of small particles, such as dust or pollen, floating in liquids. But small particles falling in air, such as mercury globules or selenium particles, and these fluctuations can be observed, for example with help of air flows.

A mirror glued on a quartz fibre hanging in the air, especially at low pressure, changes orientation randomly, by small amounts, due to the collision of the air molecules. The random orientation changes, again a kind of noise, can be followed by reflecting a beam of light on the mirror and watching the light spot at a large distance.

Also density fluctuations, critical opalescence, and critical miscibility of liquids are forms of noise. In fact, density fluctuations are important for the formation of the colour of the sky, because the density fluctuations of air molecules are the actual source of the scattering of light. The colour of the sky is a noise effect.

It turns out that *every* kind of noise can be used to count atoms. The reason is that all noise in nature is related to the particle nature of matter or radiation. Indeed, all the mentioned methods have been use to count atoms and molecules, and to determine their sizes. Since the colour of the sky is a noise effect, one can indeed count air molecules by looking at the sky!

The result of all these measurements is that a *mol* of matter – for any gas, that is the amount of matter contained in 22.41 of that gas at standard pressure – always contains the same number of atoms.

\triangleright A mol contains $6.0 \cdot 10^{23}$ particles.

The number is called *Avogadro's number*, after the first man who understood that volumes of gases have equal number of molecules, or *Loschmidt's number*, after the first man who measured it.** All the methods to determine Avogadro's number also allow us to deduce that most atoms have a size in the range between 0.1 and 0.3 nm. Molecules are composed of several or many atoms and are correspondingly larger.

How did Joseph Loschmidt*** manage to be the first to determine his and Avogadro's number, and be the first to determine reliably the size of the components of matter?

Loschmidt knew that the dynamic viscosity μ of a gas was given by $\mu = \rho lv/3$, where ρ is the density of the gas, v the average speed of the components and l their mean free path. With Avogadro's prediction (made in 1811 without specifying any value) that a volume V of any gas always contains the same number N of components, one also has $l = V/\sqrt{2\pi N\sigma^2}$, where σ is the cross-section of the components. (The cross-section is roughly the area of the shadow of an object.) Loschmidt then assumed that when the gas is liquefied, the volume of the liquid is the sum of the volumes of the particles. He

on Mountain – The Adventure of Physics | pdf file available free of charge at www.motionmountain.net | Copyright © Christoph Schiller November 1997–March 201

^{*} There are also various methods to count atoms by using electrolysis and determining the electron charge, by using radioactivity, X-ray scattering or by determining Planck's constant \hbar . We leave them aside here, because these methods actually *count* atoms. They are more precise, but also less interesting.

^{**} The term 'Loschmidt's number' is sometimes also used to designate the number of molecules in one cubic centimetre of gas.

^{***} Joseph Loschmidt (b. 1821 Putschirn, d. 1895 Vienna) Austrian chemist and physicist.



FIGURE 189 Soap bubbles show visible effects of molecular size: before bursting, soap bubbles show small transparent spots; they appear black in this picture due to a black background. These spots are regions where the bubble has a thickness of only two molecules, with no liquid in between (© LordV).

then measured all the involved quantities, for mercury, and determined N. He thus determined the number of particles in one mole of matter, in one cubic centimetre of matter, and also the size of these particles.

EXPERIENCING ATOMS

Matter is not continuous nor fractal. Matter contains smallest components with a characteristic size. Can we see effects of single atoms or molecules in everyday life? Yes, we can. We just need to watch *soap bubbles*. Soap bubbles have colours. But just before they burst, on the upper side of the bubble, the colours are interrupted by small transparent spots, as shown in Figure 189. Why? Inside a bubble, the liquid flows downwards, so that over time, the bubble gets thicker at the bottom and thinner at the top. After a while, in some regions all the liquid is gone, and in these regions, the bubble consists only of two molecular layers of soap molecules.

In fact, the arrangement of soap or oil molecules on water surfaces can be used to measure Avogadro's number. This has been done in various ingenious ways, and yields an extremely precise value with very simple means.

A simple experiment showing that solids have smallest components is shown in Figure 190. A cylindrical rod of pure, single crystal aluminium shows a surprising behaviour when it is illuminated from the side: its brightness depends on how the rod is oriented, even though it is completely round. This angular dependence is due to the atomic arrangement of the aluminium atoms in the rod.

Ref. 234

292 11 EXTENDED BODIES




FIGURE 190 Atoms exist: rotating an illuminated, perfectly round single crystal aluminium rod leads to brightness oscillations because of the atoms that make it up




FIGURE 191 Atomic steps in broken gallium arsenide crystals (wafers) can be seen under a light microscope.





FIGURE 192 An effect of atoms: steps on single crystal surfaces – here silicon carbide grown on a carbon-terminated substrate (left) and on a silicon terminated substrate (right) observed in a simple light microscope (© Dietmar Siche).

Challenge 504 ny

It is not difficult to confirm experimentally the existence of smallest size in crystals. It is sufficient to break a single crystal, such as a gallium arsenide wafer, in two. The breaking surface is either completely flat or shows extremely small steps, as shown in Figure 191. These steps are visible under a normal light microscope. (Why?) Similarly, Figure 192 shows a defect that appeared in crystal growth. It turns out that all such step heights are multiples of a smallest height: its value is about 0.2 nm. The existence of a smallest height, corresponding to the height of an atom, contradicts all possibilities of scale invariance in matter.

SEEING ATOMS

Ref. 235, Ref. 236

Ref. 237

Nowadays, with advances in technology, single atoms can be seen, photographed, hologrammed, counted, touched, moved, lifted, levitated, and thrown around. And all these manipulations confirm that like everyday matter, atoms have mass, size, shape and colour. Single atoms have even been used as lamps and as lasers. Some experimental results are

tain – The Adventure of Physics 🏻 pdf file available free of charge at www.motionmountain.net 🔻 Copyright © Christoph Schiller November 1997–March 2012

FIGURE 193 A single barium ion levitated in a Paul trap (image size around 2 mm) at the centre of the picture, visible also to the naked eye in the original experiment, performed in 1985 (© Werner Neuhauser).




FIGURE 194 The atoms on the surface of a silicon crystal, mapped with an atomic force microscope (© Universität Augsburg)




FIGURE 195 The result of moving helium atoms on a metallic surface. Both the moving and the imaging was performed with an atomic force microscope (© IBM).






FIGURE 196 The principle and a realization of an atomic force microscope (photograph © Nanosurf).

shown in Figure 193, Figure 194, Figure 194 and Figure 195.

Ref. 238

Ref. 239

Ref. 240

The Greek imagined nature as a Lego set. And indeed, many modern researchers in several fields have fun playing with atoms in the same way that children play with Lego. A beautiful demonstration of these possibilities is provided by the many applications of the atomic force microscope. If you ever have the opportunity to use one, do not miss it! An atomic force microscope is a simple table-top device which follows the surface of an object with an atomically sharp needle;* such needles, usually of tungsten, are easily manufactured with a simple etching method. The changes in the height of the needle along its path over the surface are recorded with the help of a deflected light ray, as shown in Figure 196. With a little care, the atoms of the object can be felt and made visible on a computer screen. With special types of such microscopes, the needle can be used to move atoms one by one to specified places on the surface. It is also possible to scan a surface, pick up a given atom and throw it towards a mass spectrometer to determine what sort of atom it is.

Incidentally, the construction of atomic force microscopes is only a small improvement on what nature is building already by the millions; when we use our ears to listen, we are actually detecting changes in eardrum position of down to 11 pm. In other words, we all have two 'atomic force microscopes' built into our heads.

Why is it useful to know that matter is made of atoms? Given only the *size* of atoms, it is possible to deduce many material properties. The mass density, the elastic modulus, the surface tension, the thermal expansion coefficient, the heat of vaporization, the heat of fusion, the viscosity, the specific heat, the thermal diffusivity and the thermal conductivity. Just try.

Challenge 505 s

CURIOSITIES AND FUN CHALLENGES ABOUT SOLIDS

Glass is a solid. Nevertheless, many textbooks say that glass is a liquid. This error has been propagated for about a hundred years, probably originating from a mistranslation of a sentence in a German textbook published in 1933 by Gustav Tamman, *Der Glaszustand*. Can you give at least three reasons why glass is a solid and not a liquid?

Challenge 506 s

* *

Challenge 507 s

What is the maximum length of a vertically hanging wire? Could a wire be lowered from a suspended geostationary satellite down to the Earth? This would mean we could realize a space 'lift'. How long would the cable have to be? How heavy would it be? How would you build such a system? What dangers would it face?

* *

Physics is often good to win bets. See Figure 213 for a way to do so, due to Wolfgang Stalla.

* *

Matter is made of atoms. Over the centuries the stubborn resistance of many people to this idea has lead to the loss of many treasures. For over a thousand years, people thought that genuine pearls could be distinguished from false ones by hitting them with a

 $^{^*}$ A cheap version costs only a few thousand euro, and will allow you to study the difference between a silicon wafer – crystalline – a flour wafer – granular-amorphous – and a consecrated wafer.

hammer: only false pearls would break. However, all pearls break. (Also diamonds break in this situation.) Due to this belief, over the past centuries, all the most beautiful pearls in the world have been smashed to pieces.

Comic books have difficulties with the concept of atoms. Could Asterix really throw Romans into the air using his fist? Are Lucky Luke's precise revolver shots possible? Can Spiderman's silk support him in his swings from building to building? Can the Roadrunner stop running in three steps? Can the Sun be made to stop in the sky by command? Can space-ships hover using fuel? Take any comic-book hero and ask yourself whether matter made of atoms would allow him the feats he seems capable of. You will find that most cartoons are comic precisely because they assume that matter is not made of atoms, but continuous! In a sense, atoms make life a serious adventure.

Can humans start earthquakes? What would happen if 1000 million Indians were to jump at the same time from the kitchen table to the floor?

In fact, several strong earthquakes *have* been triggered by humans. This has happened when water dams have been filled, or when water has been injected into drilling holes. It has also been suggested that the extraction of deep underground water also causes earthquakes. If this is confirmed by future research, a sizeable proportion of all earthquakes could be human-triggered.

How can a tip of a stalactite be distinguished from a tip of a stalagmite? Does the difference exist also for icicles?

Fractals do not exist. Which structures approximate them most closely? One candidate is the lung. Its bronchi divide over and over, between 26 and 28 times. Each end then arrives at one of the 300 million alveoli, the 0.25 mm cavities in which oxygen is absorbed into the blood and carbon dioxide is expelled in to the air.

How much more weight would your bathroom scales show if you stood on them in a vacuum?

One of the most complex extended bodies is the human body. In modern simulations of the behaviour of humans in car accidents, the most advanced models include ribs, vertebrae, all other bones and the various organs. For each part, its specific deformation properties are taken into account. With such models and simulations, the protection of passengers and drivers in cars can be optimized.

The human body is a remarkable structure. It is stiff and flexible, as the situation demands.

Challenge 508 e

Challenge 509 s

Challenge 510 s

Challenge 511 s




FIGURE 197 A stiff structure in which no rigid piece is attached to any other one (© Kenneth Snelson).

Additionally, most stiff parts, the bones, are not attached to other stiff parts. Since a few years, artists and architects have started exploring such structures. An example of such a structure, a tower, is shown in Figure 197. It turns out that similar structures – sometimes called *tensegrity* structures – are good models for the human spine, for example. Just search the internet for more examples.

* *

The deepest hole ever drilled into the Earth is 12 km deep. In 2003, somebody proposed to enlarge such a hole and then to pour millions of tons of liquid iron into it. He claimed that the iron would sink towards the centre of the Earth. If a measurement device communication were dropped into the iron, it could send its observations to the surface using sound waves. Can you give some reasons why this would not work?

Challenge 512 s

* *

The economic power of a nation has long been associated with its capacity to produce high-quality steel. Indeed, the Industrial Revolution started with the mass production of steel. Every scientist should know the basics facts about steel. *Steel* is a combination of

Motion Mountain – The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christoph Schiller November 1997–Ma

TABLE 41 Steel types, properties and uses.

FERRITIC STEEL	AUSTENITIC STEEL	MARTENSITIC STEEL	
Definition			
'usual' steel	'soft' steel	hardened steel, brittle	
body centred cubic (bcc)	face centred cubic (fcc)	body centred tetragonal (bct)	
iron and carbon	iron, chromium, nickel, manganese, carbon	carbon steel and alloys	
Examples			
construction steel	most stainless (18/8 Cr/Ni) steels	knife edges	
car sheet steel	kitchenware	drill surfaces	
ship steel	food industry	spring steel, crankshafts	
12 % Cr stainless ferrite	Cr/V steels for nuclear reactors		
Properties			
phases described by the iron-carbon phase diagram	phases described by the Schaeffler diagram	phases described by the iron-carbon diagram and the TTT (time-temperature transformation) diagram	
in equilibrium at RT	some alloys in equilibrium at RT	not in equilibrium at RT, but stable	
mechanical properties and grain size depend on heat treatment	mechanical properties and grain size depend on thermo-mechanical pre-treatment	mechanical properties and grain size strongly depend on heat treatment	
hardened by reducing grain size, by forging, by increasing carbon content or by nitration	hardened by cold working only	hard anyway – made by laser irradiation, induction heating, etc.	
grains of ferrite and paerlite, with cementite (Fe ₃ C)	grains of austenite	grains of martensite	
ferromagnetic	not magnetic or weakly magnetic	ferromagnetic	

iron and carbon to which other elements, mostly metals, may be added as well. One can distinguish three main types of steel, depending on the crystalline structure. *Ferritic steels* have a body-centred cubic structure, as shown in Figure 198, *austenitic steels* have a face-centred cubic structure, and *martensitic steels* have a body-centred tetragonal structure. Table 41 gives further details.

* *

A simple phenomenon which requires a complex explanation is the cracking of a whip. Since the experimental work of Peter Krehl it has been known that the whip cracks when the tip reaches a velocity of *twice* the speed of sound. Can you imagine why?

298 11 EXTENDED BODIES



FIGURE 198 Ferritic steels are bcc (body centred cubic), as shown by the famous *Atomium* in Brussels, a section of an iron crystal magnified to a height of over 100 m (photo and building are © Asbl Atomium Vzw – SABAM Belgium 2007).

* *

A bicycle chain is an extended object with no stiffness. However, if it is made to rotate rapidly, it acquires dynamical stiffness, and can roll down an inclined plane or along the floor. This surprising effect can be watched at the www.iwf.de/iwf/medien/infothek? Signatur=C+14825 website.

* *

Mechanical devices are not covered in this text. There is a lot of progress in the area even at present. For example, people have built robots that are able to ride a unicycle. But even the physics of human unicycling is not simple. Try it; it is an excellent exercise to stay young.

* *

There are many arguments against the existence of atoms as hard balls. Thomson-Kelvin put it in writing: "the monstrous assumption of infinitely strong and infinitely rigid pieces of matter." Even though Thomson was right in his comment, atoms do exist. Why?

Ref. 242

Ref. 243

* *

Sand has many surprising ways to move, and new discoveries are still made regularly. In 2001, Sigurdur Thoroddsen and Amy Shen discovered that a steel ball falling on a bed of sand produces, after the ball has sunk in, a *granular jet* that jumps out upwards from the sand. Figure 199 shows a sequence of photographs of the effect. The discovery has led to a stream of subsequent research.




FIGURE 199 An example of motion of sand: granular jets (© Amy Shen).





FIGURE 200 Modern engineering highlights: a lithography machine for the production of integrated circuits and a paper machine (© ASML, Voith).

* *

Engineering is not a part of this text. Nevertheless, it is an interesting topic. A few examples of what engineers do are shown in Figure 200 and Figure 201.

300 11 EXTENDED BODIES



FIGURE 201 Sometimes unusual moving objects cross German roads (© RWE).

SUMMARY ON ATOMS

In summary, matter is not scale invariant: in particular, it is neither smooth (continuous) nor fractal. There are no arbitrary small parts in matter. Everyday matter is made of countable components: everyday matter is made of atoms. This has been confirmed for all solids, liquids and gases. Pictures from atomic force microscopes show that the size and arrangement of atoms produce the shape and the extension of objects, confirming the Lego model of matter due to the ancient Greek. Different types of atoms, as well as their various combinations, produce different types of substances.

Studying matter in even more detail - as will be done later on - yields the now well-known idea that matter, at higher and higher magnifications, is made of molecules, atoms, nuclei, protons and neutrons, and finally, quarks. Atoms also contain electrons. A final type of matter, neutrinos, is observed coming from the Sun and from certain types of radioactive materials. Even though the fundamental bricks have become somewhat smaller in the twentieth century, this will *not* happen in the future. The basic idea of the ancient Greek remains: matter is made of smallest entities, nowadays called elementary particles. In the parts on quantum theory of our mountain ascent we will explore the consequences in detail. We will discover later on that the discreteness of matter is itself a consequence of the existence of a smallest change in nature.

Vol. V, page 193

Due to the existence of atoms, the description of everyday motion of extended objects can be reduced to the description of the motion of their atoms. Atomic motion will be

a major theme in the following pages. Two of its consequences are especially important: pressure and heat. We study them now.



CHAPTER 12

FLUIDS AND THEIR MOTION

Ref. 246

LUIDS can be liquids or gases, including plasmas. And the motion of luids can be exceedingly intricate, as Figure 202 shows. In fact, luid motion is so common – think about breathing, blood circulation or the weather – that exploring is worthwhile.

THE STATE OF A FLUID

To describe motion means to describe the state of a system. For most fluids, the state at every point in space is described by composition, velocity, temperature and pressure. We will explore temperature below. We thus have one new observable:

Page 319

▶ The *pressure* at a point in a fluid is the force per area that a body of negligible size feels at that point.

Pressure is measured with the help of barometers or similar instruments. The unit of pressure is the pascal: 1 Pa is 1 N/m^2 . A selection of pressure values found in nature is given in Table 42.

Pressure is not a simple property. Can you explain the observations of Figure 204? If the hydrostatic paradox would not be true, it would be easy to make perpetuum mobiles. Can you think about an example?

Challenge 515 e

Air has a considerable pressure, of the order of 100 kPa. As a result, human forces are often to weak to overcome it, as Figure 205 shows.

THE STATE OF A FLUID

Like all motion, fluid motion obeys energy conservation. In the case that no energy is transformed into heat, the conservation of energy is particularly simple. Motion that does not generate heat implies the lack of vortices; such fluid motion is called *laminar*. If the speed of the fluid does not depend on time at all positions, it is called *stationary*. For motion that is both laminar and stationary, energy conservation can be expressed with speed v and pressure p:

$$\frac{1}{2}\rho v^2 + p + \rho g h = \text{const} \tag{105}$$












FIGURE 202 Examples of fluid motion: a vertical water jet striking a horizontal impactor, two jets of a glycerol–water mixture colliding at an oblique angle, a water jet impinging on a reservoir, a glass of wine showing tears (all © John Bush, MIT) and a dripping water tap (© Andrew Davidhazy).

TABLE 42 Some measured pressure values.

OBSERVATION	Pressure
Record negative pressure (tension) measured in water, after careful purification Ref. 247	−140 MPa = −1400 bar
Negative pressure measured in tree sap (xylem) Ref. 248, Ref. 232	up to −10 MPa = −100 bar
Negative pressure in gases	does not exist
Negative pressure in solids	is called tension
Record vacuum pressure achieved in laboratory	10 pPa (10 ⁻¹³ torr)
Pressure variation at hearing threshold	20 μΡα
Pressure variation at hearing pain	100 Pa
Atmospheric pressure in La Paz, Bolivia	51 kPa
Atmospheric pressure in cruising passenger aircraft	75 kPa
Time-averaged pressure in pleural cavity in human thorax	0.5 kPa 5 mbar below atmospheric pressure
Standard sea-level atmospheric pressure	101.325 kPa or 1013.25 mbar or 760 torr
Healthy human arterial blood pressure at height of the heart: systolic, diastolic	17 kPa,11 kPa above atmospheric pressure
Record pressure produced in laboratory, using a diamond anvil	c. 200 GPa
Pressure at the centre of the Earth	c. 370(20) GPa
Pressure at the centre of the Sun	c. 24 PPa
Pressure at the centre of a neutron star	$c. 4 \cdot 10^{33} \text{ Pa}$
Planck pressure (maximum pressure possible in nature)	$4.6 \cdot 10^{113} \text{Pa}$

where h is the height above ground. This is called *Bernoulli's equation*.* In this equation, the last term is only important if the fluid rises against ground. The first term is the

^{*} Daniel Bernoulli (b. 1700 Bâle, d. 1782 Bâle), important Swiss mathematician and physicist. His father Johann and his uncle Jakob were famous mathematicians, as were his brothers and some of his nephews. Daniel Bernoulli published many mathematical and physical results. In physics, he studied the separation of compound motion into translation and rotation. In 1738 he published the *Hydrodynamique*, in which he deduced all results from a single principle, namely the conservation of energy. The so-called *Bernoulli equation* states that (and how) the pressure of a fluid decreases when its speed increases. He studied the tides and many complex mechanical problems, and explained the Boyle–Mariotte gas 'law'. For his publications



FIGURE 203 Daniel Bernoulli (1700-1782)

FIGURE 204 The hydrostatic and the hydrodynamic paradox.

kinetic energy (per volume) of the fluid, and the other two terms are potential energies (per volume). Indeed, the second term is the potential energy (per volume) resulting from the compression of the fluid. This is due to a second way to define pressure:

Challenge 516 e

▶ Pressure is potential energy per volume

Energy conservation implies that the lower the pressure is, the larger the speed of a fluid becomes. We can use this relation to measure the speed of a stationary water flow in a tube. We just have to narrow the tube somewhat at one location along the tube, and measure the pressure difference before and at the tube restriction. The speed v far from the constriction is then given as $v = k\sqrt{p_1 - p_2}$. (What is the constant k?) A device using this method is called a Venturi gauge.

Challenge 517 s

If the geometry of a system is kept fixed and the fluid speed is increased, at a certain speed one observes a transition: the liquid loses its clarity, the flow is not stationary any more. This is seen whenever a water tap is opened. At a certain speed, the flow changes from laminar to turbulent. At this point, Bernoulli's equation is *not* valid any more.

The description of turbulence might be the toughest of all problems in physics. When the young Werner Heisenberg was asked to continue research on turbulence, he refused

he won the prestigious prize of the French Academy of Sciences – a forerunner of the Nobel Prize – ten times.



FIGURE 205 The pressure of air is stronger than humans for sufficiently large objects that enclose a vacuum (© Kornelia Möller).

– rightly so – saying it was too difficult; he turned to something easier and he discovered and developed quantum mechanics instead. Turbulence is such a vast topic, with many of its concepts still not settled, that despite the number and importance of its applications, only now, at the beginning of the twenty-first century, are its secrets beginning to be unravelled.

Ref. 249

It is thought that the equations of motion describing fluids, the so-called *Navier–Stokes equations*, are sufficient to understand turbulence.* But the mathematics behind them is mind-boggling. There is even a prize of one million dollars offered by the Clay Mathematics Institute for the completion of certain steps on the way to solving the equations.

Important systems which show laminar flow, vortices and turbulence at the same time are wings and sails. (See Figure 206.) All wings work best in laminar mode. The essence of a wing is that it imparts air a downward velocity with as little turbulence as possible. (The aim to minimize turbulence is the reason that wings are curved. If the engine is very powerful, a flat wing at an angle also works. Strong turbulence is also of advantage for landing safely.) The downward velocity of the trailing air leads to a centrifugal force acting on the air that passes above the wing. This leads to a lower pressure, and thus to lift. (Wings thus do *not* rely on the Bernoulli equation, where lower pressure *along* the

Ref. 250

^{*} They are named after Claude Navier (b. 1785 Dijon, d. 1836 Paris), important French engineer and bridge builder, and Georges Gabriel Stokes (b. 1819 Skreen, d. 1903 Cambridge), important Irish physicist and mathematician.





FIGURE 206 The moth sailing class: a 30 kg boat that sails above the water using hydrofoils, i.e., underwater wings (© Bladerider International).




FIGURE 207 Wasting money because of lack of knowledge about fluids.

flow leads to higher air speed, as unfortunately, many books used to say. Above a wing, the higher speed is related to lower pressure *across* the flow.)

The different speeds of the air above and below the wing lead to vortices at the end of every wing. These vortices are especially important for the take-off of any insect, bird and aeroplane. More details on wings are discussed later on.

Vol. V, page 206

CURIOSITIES AND FUN CHALLENGES ABOUT FLUIDS

What happens if people do not know the rules of nature? The answer is the same since 2000 years ago: taxpayer's money is wasted or health is in danger. One of the oldest examples, the aqueducts from Roman time, is shown in Figure 207. They only exist because Romans did not know how fluids move. Now you know why there are no aqueducts any more. But using a 1 or 2 m water hose in this way to transport gasoline can be dangerous.

Challenge 518 s Why?

* *

Your bathtub is full of water. You have an unmarked 3-litre container and an unmarked 5-litre container. How can you get 4 litres of water from the bathtub?

* *

The easiest way to create a supersonic jet of air is to drop a billiard ball into a bucket full Ref. 251 of water.

* *

Where did the water in the oceans come from? Interestingly enough, this question is not fully settled! In the early age of the Earth, the high temperatures made all water evaporate and escape into space. So where did today's water come from? (For example, could the hydrogen come from the radioactivity of the Earth's core?) The most plausible proposal is that the water comes from comets. Comets are made, to a large degree, of ice. Comets hitting the Earth in the distant past seem have formed the oceans. In 2011, it was shown for the first time, by the Herschel infrared space telescope of the European Space Agency, that comets from the Kuiper belt – in contrast to comets from the inner solar system – have ice of the same isotope composition as the Earth's oceans. The comet origin of oceans seems near final confirmation.

* *

The physics of under water diving, in particular of apnoea diving, is full of wonders and of effects that are not yet understood. For example, every apnoea champion known that it is quite hard to hold the breath for five or six minutes while sitting in a chair. But if the same is done in a swimming pool, the feat becomes readily achievable for modern apnoea champions. (It is not fully clear why this is the case.) There are many apnoea diving disciplines. In 2009,the no-limit diving record is at the incredible depth of 214 m, achieved by Herbert Nitsch.

When an apnoea diver reaches a depth of 100 m, the water pressure corresponds to a weight of over 11 kg on each square centimetre of his skin. To avoid the problems of ear pressure compensation at great depths, a diver has to flood the mouth and the trachea with water. His lungs have shrunk to one eleventh of their original size, to the size of apples. The water pressure shifts almost all blood from the legs and arms into the thorax and the brain. At 150 m, there is no light, and no sound – only the heart beat. And the heart beat is slow: there is only a beat every seven or eight seconds. He becomes relaxed and euphoric at the same time. None of these fascinating observations is fully understood.

* *

What is the speed record for motion under water? Probably nobody knows: it is a military secret. In fact, the answer needs to be split. The fastest published speed for a projectile under water, almost fully enclosed in a gas bubble, is 1550 m/s, faster than the speed of sound in water, achieved over a distance of a few metres in a military laboratory in the 1990s. The fastest system with an engine seems to be a torpedo, also moving mainly in a gas bubble, that reaches over 120 m/s, thus faster than any formula 1 racing car. The

Ref. 25

exact speed achieved is higher and secret. (The method of enclosing objects under water in gas bubbles, called *supercavitation*, is a research topic of military engineers all over the world.) The fastest fish, the sailfish *Istiophorus platypterus*, reaches 22 m/s, but speeds up to 30 m are suspected. The fastest manned objects are military submarines, whose speeds are secret, but believed to be around 21 m/s. (All military naval engineers in this world, with the enormous budgets they have, are not able to make submarines that are faster than fish. The reason that aeroplanes are faster than birds is evident: aeroplanes were not developed by military engineers.) The fastest human-powered submarines reach around 4 m/s. One can guess that if human-powered submarine developers had the same development budget as military engineers, their machines would probably be faster than nuclear submarines.

There are no record lists for swimming under water. Underwater swimming is known to be faster than above-water breast stroke, back stroke or dolphin stroke: that is the reason that swimming underwater over long distances is forbidden in competitions in these styles. However, it is not known whether crawl-style records are faster or slower than records for the fastest swimming style below water. Which one is faster in your own case?

Challenge 520 e

Challenge 521 e

* *

How much water is necessary to moisten the air in a room in winter? At 0° C, the vapour pressure of water is 6 mbar, 20° C it is 23 mbar. As a result, heating air in the winter gives at most a humidity of 25%. To increase the humidity by 50%, one thus needs about 1 litre of water per $100 \, \text{m}^3$.

* *

Fluid motion is of vital importance. There are at least four fluid circulation systems inside the human body. First, *air* is circulated inside the lungs by the diaphragm and other chest muscles. Second, *blood* flows through the blood system by the heart. Third, *lymph* flows through the lymphatic vessels, moved passively by body muscles. Fourth, the *cerebrospinal fluid* circulates around the brain and the spine, moved by motions of the head. For this reason, medical doctors like the simple statement: every illness is ultimately due to bad circulation.

Why do animals have blood and other circulation systems? Because fluid diffusion is too slow. Can you detail the argument?

Challenge 522 e

* *

All animals have similar blood circulation speeds, namely between 0.2 m/s and 0.4 m/s. Why?

Challenge 523 ny

•

Liquid pressure depends on height. If the average human blood pressure at the height of the heart is 13.3 kPa, can you guess what it is inside the feet when standing?

* *

The human heart pumps blood at a rate of about 0.1 l/s. A typical capillary has the di-

Challenge 526 s

Challenge 531 s

ameter of a red blood cell, around $7 \mu m$, and in it the blood moves at a speed of half a millimetre per second. How many capillaries are there in a human?

* *

You are in a boat on a pond with a stone, a bucket of water and a piece of wood. What happens to the water level of the pond after you throw the stone in it? After you throw the water into the pond? After you throw the piece of wood?

* *

Challenge 527 s A ship leaves a river and enters the sea. What happens?

* *

Put a rubber air balloon over the end of a bottle and let it hang inside the bottle. How much can you blow up the balloon inside the bottle?

* *

Put a small paper ball into the neck of a horizontal bottle and try to blow it into the bottle.

Challenge 529 e The paper will fly *towards* you. Why?

* *

It is possible to blow an egg from one egg-cup to a second one just behind it. Can you perform this trick?

* *

In the seventeenth century, engineers who needed to pump water faced a challenge. To pump water from mine shafts to the surface, no water pump managed more than 10 m of height difference. For twice that height, one always needed two pumps in series, connected by an intermediate reservoir. Why? How then do trees manage to pump water upwards for larger heights?

* *

When hydrogen and oxygen are combined to form water, the amount of hydrogen needed is exactly twice the amount of oxygen, if no gas is to be left over after the reaction.

Challenge 532 s How does this observation confirm the existence of atoms?

* *

Challenge 533 s How are alcohol-filled chocolate pralines made? Note that the alcohol is not injected into them afterwards, because there would be no way to keep the result tight enough.

* *

How often can a stone jump when it is thrown over the surface of water? The present world record was achieved in 2002: 40 jumps. More information is known about the previous world record, achieved in 1992: a palm-sized, triangular and flat stone was thrown with a speed of 12 m/s (others say 20 m/s) and a rotation speed of about 14 revolutions per second along a river, covering about 100 m with 38 jumps. (The sequence was filmed with a video recorder from a bridge.)





FIGURE 208 What is your personal stone-skipping record?



Challenge 537 s

FIGURE 209 Heron's fountain in operation.

What would be necessary to increase the number of jumps? Can you build a machine Challenge 534 r that is a better thrower than yourself?

* *

The most abundant component of air is nitrogen (about 78 %). The second component is oxygen (about 21 %). What is the third one?

* *

Which everyday system has a pressure lower than that of the atmosphere and usually Challenge 536 s kills a person if the pressure is raised to the usual atmospheric value?

* *

Water can flow uphill: Heron's fountain shows this most clearly. Heron of Alexandria (*c*. 10 to *c*. 70) described it 2000 years ago; it is easily built at home, using some plastic bottles and a little tubing. How does it work? How is it started?

* *

A light bulb is placed, underwater, in a stable steel cylinder with a diameter of 16 cm.

An original Fiat Cinquecento car (500 kg) is placed on a piston pushing onto the water surface. Will the bulb resist?

Challenge 538 s

* *

Challenge 539 s What is the most dense gas? The most dense vapour?

* *

Every year, the Institute of Maritime Systems of the University of Rostock organizes a contest. The challenge is to build a paper boat with the highest carrying capacity. The paper boat must weigh at most 10 g and fulfil a few additional conditions; the carrying capacity is measured by pouring small lead shot onto it, until the boat sinks. The 2008 record stands at 5.1 kg. Can you achieve this value? (For more information, see the www.paperboat.de website.)

Challenge 540 e

* *

A modern version of an old question – already posed by Daniel Colladon (1802-1893) – is the following. A ship of mass m in a river is pulled by horses walking along the river bank attached by ropes. If the river is of superfluid helium, meaning that there is no friction between ship and river, what energy is necessary to pull the ship upstream along the river until a height h has been gained?

Challenge 541 s

Challenge 542 e

* *

An urban legend pretends that at the bottom of large waterfalls there is not enough air to breathe. Why is this wrong?

* *

The Swiss physicist and inventor Auguste Piccard (1884–1962) was a famous explorer. Among others, he explored the stratosphere: he reached the record height of 16 km in his *aerostat*, a hydrogen gas balloon. Inside the airtight cabin hanging under his balloon, he had normal air pressure. However, he needed to introduce several ropes attached at the balloon into the cabin, in order to be able to pull and release them, as they controlled his balloon. How did he get the ropes into the cabin while at the same time preventing air from leaving?

Challenge 543 s

* *

A human cannot breathe at any depth under water, even if he has a tube going to the surface. At a few metres of depth, trying to do so is inevitably *fatal*! Even at a depth of 60 cm only, the human body can only breathe in this way for a few minutes, and can get hurt. Why?

Challenge 544 s

* *

A human in air falls with a limiting speed of about 50 m/s (the precise value depends on clothing). How long does it take to fall from a plane at 3000 m down to a height of 200 m?

Challenge 545 ny




FIGURE 210 Which funnel empties more rapidly?

To get an idea of the size of Avogadro's and Loschmidt's number, two questions are usually asked. First, on average, how many molecules or atoms that you breathe in with every breath have previously been exhaled by Caesar? Second, on average, how many atoms of Jesus do you eat every day? Even though the Earth is large, the resulting numbers are still telling.

A few drops of tea usually flow along the underside of the spout of a teapot (or fall onto the table). This phenomenon has even been simulated using supercomputer simulations of the motion of liquids, by Kistler and Scriven, using the Navier–Stokes equations. Teapots are still shedding drops, though.

* *

The best giant soap bubbles can be made by mixing 1.5 l of water, 200 ml of corn syrup and 450 ml of washing-up liquid. Mix everything together and then let it rest for four hours. You can then make the largest bubbles by dipping a metal ring of up to 100 mm diameter into the mixture. But why do soap bubbles burst?

* *

A drop of water that falls into a pan containing hot oil dances on the surface for a considerable time, if the oil is above 220°C. Cooks test the temperature of oil in this way. Why does this so-called *Leidenfrost effect** take place? The Leidenfrost effect allows one to plunge the bare hand into molten lead, to keep liquid nitrogen in one's mouth, to check whether a pressing iron is hot, or to walk over hot coal – if one follows several safety rules, as explained by Jearl Walker. (Do *not* try this yourself! Many things can go wrong.) The main condition is that the hand, the mouth or the feet must be wet. Walker lost two teeth in a demonstration and badly burned his feet in a walk when the condition was not met.

* *

Challenge 549 s Why don't air molecules fall towards the bottom of the container and stay there?

* *

Challenge 546 s

Ref. 255

Challenge 548 s

Challenge 547 s

Ref. 256

^{*} It is named after Johann Gottlieb Leidenfrost (1715–1794), German physician.

Challenge 550 s Which of the two water funnels in Figure 210 is emptied more rapidly? Apply energy conservation to the fluid's motion (the Bernoulli equation) to find the answer.

* *

As we have seen, fast flow generates an underpressure. How do fish prevent their eyes Challenge 551 s from popping when they swim rapidly?

* *

Golf balls have dimples for the same reasons that tennis balls are hairy and that shark and dolphin skin is not flat: deviations from flatness reduce the flow resistance because many small eddies produce less friction than a few large ones. Why?

Challenge 552 ny

* *

The recognized record height reached by a helicopter is 12 442 m above sea level, though 12 954 m has also been claimed. (The first height was reached in 1972, the second in 2002, both by French pilots in French helicopters.) Why, then, do people still continue to use their legs in order to reach the top of Mount Sagarmatha, the highest mountain in the world?

Challenge 553 s

* *

A loosely knotted sewing thread lies on the surface of a bowl filled with water. Putting a bit of washing-up liquid into the area surrounded by the thread makes it immediately become circular. Why?

Challenge 554 e

Ref. 258

* *

Challenge 555 s How can you put a handkerchief under water using a glass, while keeping it dry?

* *

Are you able to blow a ping-pong ball out of a funnel? What happens if you blow through a funnel towards a burning candle?

* *

The fall of a leaf, with its complex path, is still a topic of investigation. We are far from being able to predict the time a leaf will take to reach the ground; the motion of the air around a leaf is not easy to describe. One of the simplest phenomena of hydrodynamics remains one of its most difficult problems.

* *

Fluids exhibit many interesting effects. Soap bubbles in air are made of a thin spherical film of liquid with air on both sides. In 1932, anti-bubbles, thin spherical films of air with liquid on both sides, were first observed. In 2004, the Belgian physicist Stéphane Dorbolo and his team showed that it is possible to produce them in simple experiments, and in particular, in Belgian beer.

* *

Have you ever dropped a Mentos candy into a Diet Coca Cola bottle? You will get an



FIGURE 211 A smoke ring, around 100 m in size, ejected from Mt. Etna's Bocca Nova in 2000 (© Daniela Szczepanski at www.vulkanarchiv.de and www.vulkane.net).




FIGURE 212 Two leapfrogging vortex rings (QuickTime film © Lim Tee Tai)

Challenge 556 e interesting effect. (Do it at your own risk...) Is it possible to build a rocket in this way?

* *

Challenge 557 e A needle can swim on water, if you put it there carefully. Just try, using a fork. Why does it float?

* *

The Rhine emits about 2 300 m 3 /s of water into the North Sea, the Amazon River about 120 000 m 3 /s into the Atlantic. How much is this less than c^3 /4G?

* *

Fluids exhibit many complex motions. To see an overview, have a look at the beautiful col-



FIGURE 213 How can you move the coin into the glass without touching anything?

Challenge 559 e

lection on the web site serve.me.nus.edu.sg/limtt. Among fluid motion, vortex rings, as emitted by smokers or volcanoes, have often triggered the imagination. (See Figure 211.) One of the most famous examples of fluid motion is the leapfrogging of vortex rings, shown in Figure 212. Lim Tee Tai explains that more than two leapfrogs are extremely hard to achieve, because the slightest vortex ring misalignment leads to the collapse of the system.

Ref. 259

* *

A surprising effect can be observed when pouring *shampoo* on a plate: sometimes a thin stream is ejected from the region where the shampoo hits the plate. This so-called Kaye effect is best enjoyed in the beautiful movie produced by the University of Twente found on the youtube.com/watch?v=GX4 3cV 3Mw website.

* *

Aeroplanes toilets are dangerous places. In the 1990s, a fat person sat on the toilet seat and pushed the 'flush' button while sitting. (Never try this yourself.) The underpressure exerted by the toilet was so strong that it pulled out the intestine and the person had to be brought into hospital. (Everything ended well, by the way.)

* *

If one surrounds water droplets with the correct type of dust, the droplets can roll along inclined planes. They can roll with a speed of up to 1 m/s, whereas on the same surface, water would flow hundred times more slowly. When the droplets get too fast, they become flat disks; at even higher speed, they get a doughnut shape. Such droplets can even jump and swim.

Ref. 260

WHAT CAN MOVE IN NATURE? - FLOWS

Before we continue with the next way to describe motion globally, we have a look at the possibilities of motion in everyday life. One overview is given in Table 43. The domains that belong to everyday life – motion of fluids, of matter, of matter types, of heat, of light and of charge – are the domains of continuum physics.

Within continuum physics, there are three domains we have not yet studied: the motion of charge and light, called *electrodynamics*, the motion of heat, called *thermodynamics*, and the motion of the vacuum. Once we have explored these domains, we will have

Motion Mountain – The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christoph Schiller November 1997–M

TABLE 43 Extensive quantities in nature, i.e., quantities that flow and accumulate.

Domain	EXTENSIVE	Current	INTEN- SIVE QUANTITY	ENERGY FLOW	RESISTANCE TO TRANSPORT
	(ENERGY CARRIER)	(FLOW INTENSITY)	(DRIVING STRENGTE	•	(INTENSITY OF ENTROPY GENERA- TION)
Rivers	mass m	mass flow m/t	height difference <i>gh</i>	P = gh m/t	$R_{\rm m} = ght/m$ $[{\rm m}^2/{\rm skg}]$
Gases	volume V	volume flow V/t	pressure p	P = pV/t	$R_{\rm V} = pt/V$ $[kg/s {\rm m}^5]$
Mechanics	momentum p	force $F = \mathrm{d} \boldsymbol{p} / \mathrm{d} t$	velocity v	$P = \boldsymbol{v} F$	$R_{\rm p} = \Delta V/F = t/m [\rm s/kg]$
	angular momentum <i>L</i>	torque $M = dL/dt$	angular velocity ω	$P = \boldsymbol{\omega} \boldsymbol{M}$	$R_{\rm L} = t/mr^2$ [s/kg m ²]
Chemistry	amount of substance <i>n</i>	substance flow $I_n = dn/dt$	chemical potential μ	$P = \mu I_n$	$R_n = \mu t/n$ [Js/mol ²]
Thermo- dynamics	entropy S	entropy flow $I_S = dS/dt$	temperature T	$P = T I_S$	$R_S = Tt/S$ $[K^2/W]$
Light	like all massless radiation, it can flow but cannot accumulate				
Electricity	charge q	electrical current $I = dq/dt$	electrical potential U	P = U I	$R = U/I$ $[\Omega]$
Magnetism	no accumulable magnetic sources are found in nature				
Nuclear physics	extensive quantities exist, but do not appear in everyday life				
			_		

Gravitation empty space can move and flow, but the motion is not observed in everyday life

completed the first step of our description of motion: continuum physics. In continuum physics, motion and moving entities are described with continuous quantities that can take any value, including arbitrarily small or arbitrarily large values.

But nature is *not* continuous. We have already seen that matter cannot be indefinitely divided into ever-smaller entities. In fact, we will discover that there are precise experiments that provide limits to the observed values for *every* domain of continuum physics. There is a limit to mass, to speed, to angular momentum, to force, to entropy and to change of charge. The consequences of these discoveries lead to the next legs of our description of motion: relativity and quantum theory. Relativity is based on upper limits, quantum theory on lower limits. The last leg of our description of motion will be formed by the unification of quantum theory and general relativity.

Every domain of physics, regardless of which one of the above legs it belongs to, de-

Ref. 272

scribes change in terms of two quantities: energy, and an extensive quantity characteristic of the domain. An observable quantity is called *extensive* if it increases with system size. Table 43 provides an overview. The extensive and intensive quantities fro fluids – what flows and why it flows – are volume and pressure. The extensive and intensive quantities corresponding to what in everyday language is called 'heat' – what flows and why it flows – are *entropy* and *temperature*.

The analogies of the table can be carried even further. In all domains, the *capacity* of a system is defined by the extensive quantity divided by the intensive quantity. The capacity measures, how much easily stuff flows into the system. For electric charge the capacity is the usual electric capacity. For momentum, the capacity is called mass. Mass measures, how easily one can put momentum into a system. Can you determine the quantities that measure capacity in the other cases?

Challenge 560 e

Similarly, in all fields it is possible to store energy by using the intensive quantity – such as $E = CU^2/2$ in a capacitor or $E = mv^2/2$ in a moving body – or by using the extensive quantity – such as $E = LI^2/2$ in a coil or $E = F^2/2k$ in a spring. Combining the two, we get oscillations. Can you extend the analogy to the other cases?

Challenge 561 e

SUMMARY ON FLUIDS

The motion of fluids is the motion of its constituent particles. Fluid motion can be laminar or turbulent. Laminar motion is described by Bernoulli's equation, i.e., by energy conservation. The motion of fluids allows for swimming, flying, vortices and turbulence. The exact description of fluid motion is the most complicated problem of physics.



CHAPTER 13

FROM HEAT TO TIME-INVARIANCE

Spilled milk never returns into its container by itself. Any hot object, left alone, tarts to cool down with time; it never heats up. These and many other observations how that numerous processes in nature are *irreversible*. Further observations show that irreversibility is only found in systems composed of a many particles, and that all irreversible systems involve heat.

We are thus led to explore the next global approach for the description of motion: *statistical physics*. Statistical physics, which includes *themodynamics*, the study of temperature and heat, explains the origin of irreversibility and of many material properties.

Does irreversibility mean that motion, at a fundamental level, is not invariant under reversal, as Nobel Prize winner Ilya Prigogine, one of the fathers of self-organization, thought? In this chapter we show that he was wrong.* To deduce this result, we first need to know the basic facts about temperature and heat; then we discuss irreversibility and motion reversal.

Ref. 261

TEMPERATURE

Macroscopic bodies, i.e., bodies made of many atoms, have *temperature*. Bodies made of few atoms do not have a temperature. Ovens have high temperature, refrigerators low temperature. Temperature changes have important effects: matter changes from solid to liquid to gaseous to plasma state. With a change in temperature, matter also changes size, colour, magnetic properties, stiffness and many more.

Temperature is an aspect of the *state* of a body. In other words, two identical bodies can be characterized and distinguished by their temperature. This is well-known to criminal organizations around the world that rig lotteries. When a blind-folded child is asked to draw a numbered ball from a set of such balls, such as in Figure 215, it is often told beforehand to draw only *hot* or *cold* balls. The blindfolding also helps to hide the tears due to the pain.

Ref. 263

The temperature of a macroscopic body is an aspect of its state. In particular, temperature is an *intensive* quantity or variable. In short, temperature describes the intensity of heat. An overview of temperatures is given in Table 44.

^{*} Many even less serious thinkers often ask the question in the following term: is motion *time-invariant*? The cheap press goes even further, and asks whether motion has an 'arrow' or whether time has a preferred 'direction of flow'. We have already shown above that this is nonsense and steer clear of such phrases in the following.






FIGURE 214 Braking generates heat on the floor and in the tire (© Klaus-Peter Möllmann and Michael Vollmer).



FIGURE 215 A rigged lottery shows that temperature is an aspect of the state of a body (© ISTA).

We observe that any two bodies in contact tend towards the *same* temperature: temperature is contagious. In other words, temperature describes an equilibrium situation. The existence and contagiousness of temperature is often called the *zeroth principle of thermodynamics*. We call *heating* the increase of temperature, and *cooling* its decrease.

How is temperature measured? The eighteenth century produced the clearest answer: temperature is best defined and measured by the *expansion of gases*. For the simplest, so-called *ideal* gases, the product of pressure *p* and volume *V* is proportional to temperature:

$$pV \sim T$$
 . (106)

The proportionality constant is fixed by the *amount* of gas used. (More about it shortly.) The ideal gas relation allows us to determine temperature by measuring pressure and volume. This is the way (absolute) temperature has been defined and measured for about a century. To define the *unit* of temperature, we only have to fix the amount of gas used. It is customary to fix the amount of gas at 1 mol; for oxygen this is 32 g. The proportionality constant for 1 mol, called the *ideal gas constant R*, is defined to be $R = 8.3145 \, \text{J/mol K}$.

Ref. 262 Page 383

This number has been chosen in order to yield the best approximation to the independently defined Celsius temperature scale. Fixing the ideal gas constant in this way defines 1 K, or one Kelvin, as the unit of temperature. In simple terms, a temperature increase of one Kelvin is defined as the temperature increase that makes the volume of an ideal gas increase – keeping the pressure fixed – by a fraction of 1/273.15 or 0.3661 %.

In general, if we need to determine the temperature of an object, we take a mole of gas, put it in contact with the object, wait a while, and then measure the pressure and the volume of the gas. The ideal gas relation (106) then gives the temperature.

Most importantly, the ideal gas relation shows that there is a lowest temperature in nature, namely that temperature at which an ideal gas would have a vanishing volume. That would happen at T=0 K, i.e., at -273.15° C. In reality, other effects, like the volume of the atoms themselves, prevent the volume of the gas from ever reaching zero exactly. In fact, the unattainability of absolute zero is called the *third principle of thermodynamics*.

In fact, the temperatures achieved by a civilization can be used as a measure of its technological achievements. We can define the *Bronze Age* (1.1 kK, 3500 bCE), the *Iron Age* (1.8 kK, 1000 bCE), the *Electric Age* (3 kK from *c.* 1880) and the *Atomic Age* (several MK, from 1944) in this way. Taking into account also the quest for lower temperatures, we can define the *Quantum Age* (4 K, starting 1908). But what exactly is heating or cooling? What happens in these processes?

THERMAL ENERGY

Heating and cooling is the flow of disordered energy. For example, friction slows down moving bodies, and, while doing so, heats them up. The 'creation' of heat by friction can be tested experimentally. An example is shown in Figure 214. Heat can be generated from friction, just by continuous rubbing, without any limit. This endless 'creation' of heat implies that heat is not a material fluid or substance extracted from the body – which in this case would be consumed after a certain time – but something else. Indeed, today we know that heat, even though it behaves in some ways like a fluid, is due to disordered motion of particles. The conclusion of these studies is simple. Friction is the transformation of mechanical (i.e., ordered) energy into (disordered) thermal energy, i.e., into disordered motion of the particles making up a material.

In order to increase the temperature of 1 kg of water by 1 K using friction, 4.2 kJ of mechanical energy must be supplied. The first to measure this quantity with precision was, in 1842, the German physician Julius Robert Mayer (1814–1878). He described his experiments as proofs of the conservation of energy; indeed, he was the first person to state energy conservation! It is something of an embarrassment to modern physics that a medical doctor was the first to show the conservation of energy, and furthermore, that he was ridiculed by most physicists of his time. Worse, conservation of energy was accepted by scientists only when it was publicized many years later by two authorities: Hermann von Helmholtz – himself also a physician turned physicist – and William Thomson, who also cited similar, but later experiments by James Joule.* All of them acknowledged Mayer's

Ref. 264


Challenge 562 e

Ref. 265

^{*} Hermann von Helmholtz (b. 1821 Potsdam, d. 1894 Berlin), important Prussian scientist. William Thomson-Kelvin (b. 1824 Belfast, d. 1907 Netherhall), important Irish physicist. James Prescott Joule (b. 1818 Salford, d. 1889 Sale), English physicist. Joule is pronounced so that it rhymes with 'cool', as his descendants like to stress. (The pronunciation of the name 'Joule' varies from family to family.)

TABLE 44 Some temperature values.

TABLE 44 Some temperature values.	
O B S E R V A T I O N	Темрекатике
Lowest, but unattainable, temperature	0 K = −273.15°C
In the context of lasers, it sometimes makes (almost) sense to talk about negative temperature.	
Temperature a perfect vacuum would have at Earth's surface Vol. V, Page 102	40 zK
Sodium gas in certain laboratory experiments – coldest matter system achieved by man and possibly in the universe	0.45 nK
Temperature of neutrino background in the universe	c. 2 K
Temperature of photon gas background (or background radiation) in the universe	2.7 K
Liquid helium	4.2 K
Oxygen triple point	54.3584 K
Liquid nitrogen	77 K
Coldest weather ever measured (Antarctic)	$185 \text{K} = -88 ^{\circ} \text{C}$
Freezing point of water at standard pressure	$273.15 \mathrm{K} = 0.00 \mathrm{^{\circ}C}$
Triple point of water	$273.16 \mathrm{K} = 0.01 \mathrm{^{\circ}C}$
Average temperature of the Earth's surface	287.2 K
Smallest uncomfortable skin temperature	316 K (10 K above normal)
Interior of human body	$310.0 \pm 0.5 \mathrm{K} = 36.8 \pm 0.5 ^{\circ}\mathrm{C}$
Temperature of most land mammals	$310 \pm 3 \text{ K} = 36.8 \pm 2^{\circ}\text{C}$
Hottest weather ever measured	$343.8 \text{K} = 70.7^{\circ} \text{C}$
Boiling point of water at standard pressure	373.13 K or 99.975°C
Temperature of hottest living things: thermophile bacteria	395 K = 122°C
Large wood fire	c. 1100 K
Liquid bronze	c. 1100 K
Freezing point of gold	1337.33 K
Liquid, pure iron	1810 K
Bunsen burner flame	up to 1870 K
Light bulb filament	2.9 kK
Earth's centre	4 kK
Melting point of hafnium carbide	4.16 kK
Sun's surface	5.8 kK
Air in lightning bolt	30 kK
Hottest star's surface (centre of NGC 2240)	250 kK
Space between Earth and Moon (no typo)	up to 1 MK
Centre of white dwarf	5 to 20 MK
Sun's centre	20 MK
Centre of the accretion disk in X-ray binary stars	10 to 100 MK
Inside the JET fusion tokamak	100 MK
Centre of hottest stars	1 GK
Maximum temperature of systems without electron–positron pair generation	ca. 6 GK
Universe when it was 1 s old	100 GK
Hagedorn temperature	1.9 TK
Heavy ion collisions - highest man-made value	up to 3.6 TK
Planck temperature – nature's upper temperature limit	10^{32}K



priority. Marketing by William Thomson eventually led to the naming of the unit of energy after Joule.

In summary, two medical doctors proved:

▶ *In a closed system, the sum of mechanical energy and thermal energy is constant.* This is usually called the *first principle of thermodynamics*.

Equivalently, it is impossible to produce mechanical energy without paying for it with some other form of energy. This is an important statement, because among others it means that humanity will stop living one day. Indeed, we live mostly on energy from the Sun; since the Sun is of finite size, its energy content will eventually be consumed.




FIGURE 217 Which balloon wins when the tap is opened?

Challenge 563 s Page 101 Can you estimate when this will happen?

The first principle of thermodynamics, the conservation of energy, implies that there is no perpetuum mobile of the first kind: no machine can run without energy input. For the same reason, we need food to eat: the energy in the food keeps us alive. If we stop eating, we die. The conservation of energy also makes most so-called 'wonders' impossible: in nature, energy cannot be created, but is conserved.

Thermal energy is a form of energy. Thermal energy can be stored, accumulated, transferred, transformed into mechanical energy, electrical energy or light. In short, thermal energy can be transformed into motion, into work, and thus into money.

The first principle of thermodynamics also allows us to formulate what a car engine achieves. Car engines are devices that transform hot matter – the hot exploding fuel inside the cylinders – into motion of the car wheels. Car engines, like steam engines, are thus examples of *heat engines*.

The study of heat and temperature is called *thermostatics* if the systems concerned are at equilibrium, and *thermodynamics* if they are not. In the latter case, we distinguish situations *near* equilibrium, when equilibrium concepts such as temperature can still be used, from situations *far* from equilibrium, such as self-organization, where such concepts often cannot be applied.

Page 348

Does it make sense to distinguish between thermal energy and heat? It does. Many older texts use the term 'heat' to mean the *same* as thermal energy. However, this is confusing; in this text, 'heat' is used, in accordance with modern approaches, as the everyday term for *entropy*. Both thermal energy and heat flow from one body to another, and both accumulate. Both have no measurable mass.* Both the amount of thermal energy and the amount of heat inside a body increase with increasing temperature. The precise relation will be given shortly. But heat has many other interesting properties and stories to tell. Of these, two are particularly important: first, heat is due to particles; and secondly, heat is at the heart of the difference between past and future. These two stories are intertwined.

Why do balloons take up space? - The end of continuity

Heat properties are material-dependent. Studying thermal properties therefore should enable us to understand something about the constituents of matter. Now, the simplest

Vol. II, page 65

^{*} This might change in future, when mass measurements improve in precision, thus allowing the detection of relativistic effects. In this case, temperature increase may be detected through its related mass increase. However, such changes are noticeable only with twelve or more digits of precision in mass measurements.



FIGURE 218 What happened here? (© Johan de Jong)

materials of all are gases.* Gases need space: any amount of gas has pressure and volume. It did not take a long time to show that gases *could not* be continuous. One of the first scientists to think about gases as made up of atoms or molecules was Daniel Bernoulli. Bernoulli reasoned that if gases are made up of small particles, with mass and momentum, he should be able to make quantitative predictions about the behaviour of gases, and check them with experiment. If the particles fly around in a gas, then the *pressure* of a gas in a container is produced by the steady flow of particles hitting the wall. It was then easy to conclude that if the particles are assumed to behave as tiny, hard and perfectly elastic balls, the pressure p, the volume V and the temperature T must be related by

Challenge 564 ny

$$pV = kNT (107)$$

where N is the number of particles contained in the gas. (The Boltzmann constant k, one of the fundamental constants of nature, is defined below.) A gas made of particles with such textbook behaviour is called an *ideal gas*. Relation (107), often called the *ideal gas 'law'*, has been confirmed by experiments at room and higher temperatures, for all known gases.

Bernoulli thus derived the ideal gas relation, with a specific prediction for the proportionality constant, from the single assumption that gases are made of small massive particles. This derivation provides a clear argument for the existence of atoms and for their behaviour as normal, though small objects. We have already seen above how N can be determined experimentally.

Page 289

The ideal gas model helps us to answer questions such as the one illustrated in Figure 217. Two *identical* rubber balloons, one filled up to a larger size than the other, are connected via a pipe and a valve. The valve is opened. Which one deflates?

Challenge 565 s

^{*} By the way, the word gas is a modern construct. It was coined by the Brussels alchemist and physician Johan Baptista van Helmont (1579–1644), to sound similar to 'chaos'. It is one of the few words which have been invented by one person and then adopted all over the world.

Challenge 566 e

The ideal gas relation states that hotter gases, at given pressure, need more volume. The relation thus explains why winds and storms exist, why hot air balloons rise – even those of Figure 218 – why car engines work, why the ozone layer is destroyed by certain gases, or why during the extremely hot summer of 2001 in the south of Turkey, oxygen masks were necessary to walk outside during the day.

The ideal gas relation also explains why on the 21st of August 1986, over a thousand people and three-thousand livestock where found dead in their homes in Cameroon. They were living below a volcano whose crater contains a lake, *Lake Nyos*. It turns out that the volcano continuously emits carbon dioxide, or CO_2 , into the lake. The carbon dioxide is usually dissolved in the water. But in August 1986, an unknown event triggered the release of a bubble of around one million tons of CO_2 , about a cubic kilometre, into the atmosphere. Because carbon dioxide $(2.0 \, \text{kg/m}^3)$ is denser than air $(1.2 \, \text{kg/m}^3)$, the gas flowed down into the valleys and villages below the volcano. The gas has no colour and smell, and it leads to asphyxiation. It is unclear whether the outgassing system installed in the lake after the event is sufficiently powerful to avoid a recurrence of the event.

Using the ideal gas relation you are now able to explain why balloons increase in size as they rise high up in the atmosphere, even though the air is colder there. The largest balloon built so far had a diameter, at high altitude, of 170 m, but only a fraction of that value at take-off. How much?

Now you can also take up the following challenge: how can you measure the weight of a car or a bicycle with a ruler only?

The picture of gases as being made of hard constituents without any long-distance interactions breaks down at very low temperatures. However, the ideal gas relation (107) can be improved to overcome these limitations, by taking into account the deviations due to interactions between atoms or molecules. This approach is now standard practice and allows us to measure temperatures even at extremely low values. The effects observed below 80 K, such as the solidification of air, frictionless transport of electrical current, or frictionless flow of liquids, form a fascinating world of their own, the beautiful domain of low-temperature physics. The field will be explored later on.

Challenge 567 ny

Challenge 568 s

Ref. 266


Ref. 267

Vol. V, page 62 Vol. V, page 70

BROWNIAN MOTION

If fluids are made of particles moving randomly, this random motion should have observable effects. Indeed, under a microscope it is easy to observe that small particles, such as coal dust, in or on a liquid never come to rest. An example of the observed motion is shown in Figure 220. The particles seem to follow a random zig zag movement. This was first described by Lucretius, in the year 60 BCE, in his poem *De natura rerum*. He describes what everybody has seen: the dance of dust particles in air that is illuminated by the Sun.

In 1785, Jan Ingenhousz saw that coal dust particles never come to rest. He discovered what is called *Brownian motion* today. 40 years after him, the botanist Robert Brown was the first Englishman to repeat the observation, this time for small particles floating in vacuoles *inside* pollen. Further experiments showed that the observation of a random motion is independent of the type of particle and of the type of liquid. In other words, Ingenhousz had discovered a fundamental form of *noise* in nature. Around 1860, the random motion of particles in liquids was attributed to the molecules of the liquid colliding



Vol. III, page 151 FIGURE 219 An image of pollen grains – field size about 0.3 mm – made with an electron microscope (Wikimedia).

with the particles by various people. In 1905 and 1906, Marian von Smoluchowski and, independently, Albert Einstein argued that this attribution could be tested experimentally, even though at that time nobody was able to observe molecules directly. The test makes use of the specific properties of thermal noise.

It had already been clear for a long time that if molecules, i.e., indivisible matter particles, really existed, then thermal energy had to be disordered motion of these constituents and temperature had to be the average energy per degree of freedom of the constituents. Bernoulli's model of Figure 223 implies that for monatomic gases the kinetic energy $T_{\rm kin}$ per particle is given by

Challenge 569 ny

Ref. 268


$$T_{\rm kin} = \frac{3}{2}kT\tag{108}$$

where T is temperature. The so-called *Boltzmann constant* $k = 1.4 \cdot 10^{-23} \text{ J/K}$ is the standard conversion factor between temperature and energy.* At a room temperature of 293 K, the kinetic energy of a particle is thus 6 zJ.

Vol. III, page 125

Planck named the Boltzmann constant after the important Austrian physicist Ludwig Boltzmann (b.

^{*} The Boltzmann constant k was discovered and named by Max Planck, in the same work in which he also discovered what is now called Planck's constant \hbar , the quantum of action. For more details on Max Planck, see later on.






FIGURE 220 Example paths for particles in Brownian motion and their displacement distribution.

Challenge 570 e

Using relation (108) to calculate the speed of air molecules at room temperature yields values of several hundred metres per second, about the speed of sound! Given this large speed, why does smoke from a candle take so long to diffuse through a room? Rudolph Clausius (1822–1888) answered this question in the mid-nineteenth century: smoke diffusion is slowed by the collisions with air molecules, in the same way as pollen particles collide with molecules in liquids.

The description of Brownian motion can be tested by following the displacement of pollen particles under the microscope. At first sight, we might guess that the average distance the pollen particle has moved after n collisions should be zero, because the molecule velocities are random. However, this is wrong, as experiment shows.

An increasing average *square* displacement, written $\langle d^2 \rangle$, is observed for the pollen particle. It cannot be predicted in which direction the particle will move, but it does move. If the distance the particle moves after one collision is l, the average square displacement after n collisions is given, as you should be able to show yourself, by

Challenge 571 ny

$$\langle d^2 \rangle = nl^2 \ . \tag{109}$$

For molecules with an average velocity v over time t this gives

$$\langle d^2 \rangle = nl^2 = vlt \ . \tag{110}$$

In other words, the average square displacement increases proportionally with time. Of course, this is only valid because the liquid is made of separate molecules. Repeatedly measuring the position of a particle should give the distribution shown in Figure 220 for the probability that the particle is found at a given distance from the starting point. This is called the (Gaussian) normal distribution. In 1908, Jean Perrin* performed ex-

Ref. 269

¹⁸⁴⁴ Vienna, d. 1906 Duino), who is most famous for his work on thermodynamics. Boltzmann he explained all thermodynamic phenomena and observables, above all entropy itself, as results of the behaviour of molecules. It is said that Boltzmann committed suicide partly because of the animosities of his fellow physicists towards his ideas and himself. Nowadays, his work is standard textbook material.

^{*} Jean Perrin (1870-1942), important French physicist, devoted most of his career to the experimental

Motion Mountain – The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christoph Schiller November 1997–Ma

tensive experiments in order to test this prediction. He found that equation (110) corresponded completely with observations, thus convincing everybody that Brownian motion is indeed due to collisions with the molecules of the surrounding liquid, as had been expected.* Perrin received the 1926 Nobel Prize for these experiments.

Einstein also showed that the same experiment could be used to determine the number of molecules in a litre of water (or equivalently, the Boltzmann constant k). Can you work out how he did this?

Challenge 572 d

WHY STONES CAN BE NEITHER SMOOTH NOR FRACTAL, NOR MADE OF LITTLE HARD BALLS

The exploration of temperature yields another interesting result. Researchers first studied gases, and measured how much energy was needed to heat them by 1 K. The result is simple: all gases share only a few values, when the number of molecules N is taken into account. Monatomic gases (in a container with constant volume) require 3Nk/2, diatomic gases (and those with a linear molecule) 5Nk/2, and almost all other gases 3Nk, where $k = 1.4 \cdot 10^{-23}$ J/K is the Boltzmann constant.

Page 327

The explanation of this result was soon forthcoming: each thermodynamic degree of freedom** contributes the energy kT/2 to the total energy, where T is the temperature. So the number of degrees of freedom in physical bodies is finite. Bodies are not continuous, nor are they fractals: if they were, their specific thermal energy would be infinite. Matter is indeed made of small basic entities.

All degrees of freedom contribute to the specific thermal energy. At least, this is what classical physics predicts. Solids, like stones, have 6 thermodynamic degrees of freedom and should show a specific thermal energy of 3Nk. At high temperatures, this is indeed observed. But measurements of solids at room temperature yield lower values, and the lower the temperature, the lower the values become. Even gases show values lower than those just mentioned, when the temperature is sufficiently low. In other words, molecules and atoms behave differently at low energies: atoms are not immutable little hard balls. The deviation of these values is one of the first hints of quantum theory.

ENTROPY

- It's irreversible.- Like my raincoat!

Mel Brooks, Spaceballs, 1987

proof of the atomic hypothesis and the determination of Avogadro's number; in pursuit of this aim he perfected the use of emulsions, Brownian motion and oil films. His Nobel Prize speech (nobelprize.org/physics/laureates/1926/perrin-lecture.html) tells the interesting story of his research. He wrote the influential book *Les atomes* and founded the Centre National de la Recherche Scientifique. He was also the first to speculate, in 1901, that an atom is similar to a small solar system.

Ref. 270 Page 357

^{*} In a delightful piece of research, Pierre Gaspard and his team showed in 1998 that Brownian motion is also chaotic, in the strict physical sense given later on.

^{**} A thermodynamic degree of freedom is, for each particle in a system, the number of dimensions in which it can move plus the number of dimensions in which it is kept in a potential. Atoms in a solid have six, particles in monatomic gases have only three; particles in diatomic gases or rigid linear molecules have five. The number of degrees of freedom of larger molecules depends on their shape.




FIGURE 221 The fire pump.

Every domain of physics describes change in terms of three quantities: energy, as well as an intensive and an extensive quantity characteristic of the domain. In the domain of thermal physics, the intensive quantity is temperature. What is the corresponding extensive quantity?

The obvious guess would be 'heat'. Unfortunately, the quantity that physicists usually call 'heat' is not the same as what we call 'heat' in our everyday speech. For this historical reason, we need to introduce a new term. The extensive quantity corresponding to what we call 'heat' in everyday speech is called *entropy* in physics.*

Entropy describes the *amount* of everyday heat. Entropy is measured in joule per kelvin or J/K; some example values (per amount of matter) are listed in Table 45 and Table 46. Entropy describes everyday heat in the same way as momentum describes everyday motion. Entropy describes the amount of heat in the same way that momentum describes the amount of motion. Correspondingly, temperature describes the *intensity* of heat, in the same way that speed describes the intensity of motion.

When two objects of different speeds collide, a flow of momentum takes place between them. Similarly, when two objects differing in temperature are brought into contact, an entropy flow takes place between them. We now define the concept of entropy – 'everyday heat' – more precisely and explore its properties in some more detail.

Entropy measures the degree to which energy is mixed up inside a system, that is, the degree to which energy is spread or shared among the components of a system. When all components of a system – usually the molecules or atoms – move in the same way, in concert, the entropy of the system is low. When the components of the system move completely independently, randomly, the entropy is large. In short, entropy measures the amount of disordered energy content per temperature in a system. That is the reason that it is measured in J/K.

^{*} The term 'entropy' was invented by the German physicist Rudolph Clausius (b. 1822 Köslin, d. 1888 Bonn) in 1865. He formed it from the Greek èv 'in' and $\tau\rho \acute{o}\pi o\varsigma$ 'direction', to make it sound similar to 'energy'. The term entropy has always had the meaning given here.

In contrast, what physicists call 'heat' is a form of energy and not an extensive quantity in general.

TABLE 45 Some measured specific entropy values.

PROCESS/SYSTEM	ENTROPY VALUE
Melting of ice	1.21 kJ/K kg = 21.99 J/K mol
Iron under standard conditions	27.2 J/K mol
Water under standard conditions	70.1 J/K mol
Boiling of 1 kg of liquid water at 101.3 kPa	6.03 kJ/K = 110 J/K mol
Oxygen under standard conditions	161.1 J/K mol

TABLE 46 Some typical entropy values per particle at *standard* temperature and pressure as multiples of the Boltzmann constant.

MATERIAL	ENTROPY PER PAR-		
	TICLE		
Monatomic solids	0.3 k to 10 k		
Diamond	0.29 k		
Graphite	$0.68 \ k$		
Lead	7.79 k		
Monatomic gases	15-25 <i>k</i>		
Helium	15.2 <i>k</i>		
Radon	21.2 <i>k</i>		
Diatomic gases	15 k to 30 k		
Polyatomic solids	10 k to 60 k		
Polyatomic liquids	10 k to 80 k		
Polyatomic gases	20 k to 60 k		
Icosane	112 <i>k</i>		

Since entropy measures an amount, an extensive quantity, and not an intensity, entropy *adds up* when identical systems are composed into one. When two one-litre bottles of water at the same temperature are poured together, the entropy of the water adds up. Again, this corresponds to the behaviour of momentum: it also adds up when systems are composed. In contrast to

Like any other extensive quantity, entropy can be accumulated in a body, and entropy can flow into or out of bodies. When we transform water into steam by heating it, we say that we add entropy to the water. After the water has become steam, the added entropy is contained in the steam. In short, entropy is the exact term for what we call 'heat' in everyday speech.

When we dissolve a block of salt in water, the entropy of the total system must increase, because the disorder increases. We now explore this effect.

ENTROPY FROM PARTICLES

Once it had become clear that heat and temperature are due to the motion of microscopic particles, people asked what entropy *was* microscopically. The answer can be formulated in various ways. The two most extreme answers are:

- ▶ Entropy measures the (logarithm of the) number *W* of possible microscopic states. A given macroscopic state can have many microscopic realizations. The logarithm of this number, multiplied by the Boltzmann constant *k*, gives the entropy.*
- \triangleright Entropy is the expected number of yes-or-no questions, multiplied by $k \ln 2$, the answers of which would tell us everything about the system, i.e., about its microscopic state.

In short, the higher the entropy, the more microstates are possible. Through either of these definitions, entropy measures the quantity of randomness in a system. In other words, entropy measures the transformability of energy: higher entropy means lower transformability. Alternatively, entropy measures the *freedom* in the choice of microstate that a system has. High entropy means high freedom of choice for the microstate. For example, when a molecule of glucose (a type of sugar) is produced by photosynthesis, about 40 bits of entropy are released. This means that after the glucose is formed, 40 additional yes-or-no questions must be answered in order to determine the full microscopic state of the system. Physicists often use a macroscopic unit; most systems of interest are large, and thus an entropy of 10^{23} bits is written as 1 J/K. (This is only approximate. Can you find the precise value?)

Challenge 573 ny

Ref. 273

To sum up, entropy is thus a specific measure for the characterization of disorder of thermal systems. Three points are worth making here. First of all, entropy is not *the* measure of disorder, but *one* measure of disorder. It is therefore *not* correct to use entropy as a *synonym* for the concept of disorder, as is often done in the popular literature. Entropy is only defined for systems that have a temperature, in other words, only for systems that are in or near equilibrium. (For systems far from equilibrium, no measure of disorder has been found yet; probably none is possible.) In fact, the use of the term entropy has degenerated so much that sometimes one has to call it *thermodynamic* entropy for clarity.

Secondly, entropy is related to information *only if* information is defined also as $-k \ln W$. To make this point clear, take a book with a mass of one kilogram. At room temperature, its entropy content is about $4 \, \text{kJ/K}$. The printed information inside a book, say 500 pages of 40 lines with each containing 80 characters out of 64 possibilities, corresponds to an entropy of $4 \cdot 10^{-17} \, \text{J/K}$. In short, what is usually called 'information' in everyday life is a negligible fraction of what a physicist calls information. Entropy is defined using the *physical* concept of information.

Finally, entropy is *not* a measure for what in normal life is called the *complexity* of a situation. In fact, nobody has yet found a quantity describing this everyday notion. The task is surprisingly difficult. Have a try!

Ref. 274 Challenge 574 ny

^{*} When Max Planck went to Austria to search for the anonymous tomb of Boltzmann in order to get him buried in a proper grave, he inscribed the formula $S = k \ln W$ on the tombstone. (Which physicist would finance the tomb of another, nowadays?)

Ref. 275

Ref. 276

Ref. 278

Ref. 279

Ref 276

Vol. VI, page 27

In summary, if you hear the term entropy used with a different meaning than the expression $S = k \ln W$, beware. Somebody is trying to get you, probably with some ideology.

THE MINIMUM ENTROPY OF NATURE - THE QUANTUM OF INFORMATION

Before we complete our discussion of thermal physics we must point out in another way the importance of the Boltzmann constant k. We have seen that this constant appears whenever the granularity of matter plays a role; it expresses the fact that matter is made of small basic entities. The most striking way to put this statement is the following:

▶ *There is a smallest entropy in nature.* Indeed, for all systems, the entropy obeys

$$S \geqslant k$$
 . (111)

This result is almost 100 years old; it was stated most clearly (with a different numerical factor) by the Hungarian–German physicist Leo Szilard. The same point was made by the French physicist Léon Brillouin (again with a different numerical factor). The statement can also be taken as the *definition* of the Boltzmann constant.

The existence of a smallest entropy in nature is a strong idea. It eliminates the possibility of the continuity of matter and also that of its fractality. A smallest entropy implies that matter is made of a finite number of small components. The lower limit to entropy expresses the fact that matter is made of particles.* The limit to entropy also shows that Galilean physics cannot be correct: Galilean physics assumes that arbitrarily small quantities do exist. The entropy limit is the first of several limits to motion that we will encounter in our adventure. After we have found all limits, we can start the final leg that leads to the unified description of motion.

The existence of a smallest quantity implies a limit on the precision of measurements. Measurements cannot have infinite precision. This limitation is usually stated in the form of an indeterminacy relation. Indeed, the existence of a smallest entropy can be rephrased as an indeterminacy relation between the temperature T and the inner energy U of a system:

$$\Delta \frac{1}{T} \Delta U \geqslant \frac{k}{2} . \tag{112}$$

This relation** was given by Niels Bohr; it was discussed by Werner Heisenberg, who called it one of the basic indeterminacy relations of nature. The Boltzmann constant (divided by 2) thus fixes the smallest possible entropy value in nature. For this reason, Gilles Cohen-Tannoudji calls it the *quantum of information* and Herbert Zimmermann calls it the *quantum of entropy*.

^{*} The minimum entropy implies that matter is made of tiny spheres; the minimum *action*, which we will encounter in quantum theory, implies that these spheres are actually small clouds.

Ref. 277 ** It seems that the historical value for the right hand side, k, has to be corrected to k/2, for the same reason that the quantum of action \hbar appears with a factor 1/2 in Heisenberg's indeterminacy relations.

334

antum

t sheds
osolute
ntropy
for ev-

Vol. IV, page 13

The relation (112) points towards a more general pattern. For every minimum value for an observable, there is a corresponding indeterminacy relation. We will come across this several times in the rest of our adventure, most importantly in the case of the quantum of action and Heisenberg's indeterminacy relation.

The existence of a smallest entropy has numerous consequences. First of all, it sheds light on the third principle of thermodynamics. A smallest entropy implies that absolute zero temperature is not achievable. Secondly, a smallest entropy explains why entropy values are finite instead of infinite. Thirdly, it fixes the absolute value of entropy for every system; in continuum physics, entropy, like energy, is only defined up to an additive constant. The quantum of entropy settles all these issues.

The existence of a minimum value for an observable implies that an indeterminacy relation appears for any two quantities whose product yields that observable. For example, entropy production rate and time are such a pair. Indeed, an indeterminacy relation connects the entropy production rate P = dS/dt and the time t:

$$\Delta P \ \Delta t \geqslant \frac{k}{2} \ . \tag{113}$$

Ref. 279, Ref. 277

Challenge 575 ny

From this and the previous relation (112) it is possible to deduce all of statistical physics, i.e., the precise theory of thermostatics and thermodynamics. We will not explore this further here. (Can you show that the zeroth and third principle follows from the existence of a smallest entropy?) We will limit ourselves to one of the cornerstones of thermodynamics: the second principle.

IS EVERYTHING MADE OF PARTICLES?

A physicist is the atom's way of knowing about atoms.

Ref. 280

George Wald

Historically, the study of statistical mechanics has been of fundamental importance for physics. It provided the first demonstration that physical objects are made of interacting particles. The story of this topic is in fact a long chain of arguments showing that all the properties we ascribe to objects, such as size, stiffness, colour, mass density, magnetism, thermal or electrical conductivity, result from the interaction of the many particles they consist of. The discovery that *all objects are made of interacting particles* has often been called the main result of modern science.

Page 317

How was this discovery made? Table 43 listed the main extensive quantities used in physics. Extensive quantities are able to flow. It turns out that all flows in nature are *composed* of elementary processes, as shown in Table 47. We have seen that flows of mass, volume, charge, entropy and substance are composed. Later, quantum theory will show the same for flows of angular momentum and of the nuclear quantum numbers. *All flows are made of particles*.

Ref. 281

This success of this idea has led many people to generalize it to the statement: 'Everything we observe is made of parts.' This approach has been applied with success to chemistry with molecules, materials science and geology with crystals, electricity with electrons, atoms with elementary particles, space with points, time with instants, light

Motion Mountain - The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright @ Christoph Schiller November 1997-March

TABLE 47 Some minimum flow values found in nature.

OBSERVATION	MINIMUM FLOW
Matter flow	one molecule or one atom or one particle
Volume flow	one molecule or one atom or one particle
Angular momentum flow	Planck's quantum of action
Chemical amount of substance	e one molecule, one atom or one particle
Entropy flow	the minimum entropy
Charge flow	one elementary charge
Light flow	one single photon, Planck's quantum of action




FIGURE 222 A 111 crystal surface of a gold single crystal, every bright dot being an atom, with a surface dislocation (© CNRS).

with photons, biology with cells, genetics with genes, neurology with neurons, mathematics with sets and relations, logic with elementary propositions, and even to linguistics with morphemes and phonemes. All these sciences have flourished on the idea that everything is made of related *parts*. The basic idea seems so self-evident that we find it difficult even to formulate an alternative. Just try!

Challenge 576 ny

Vol. VI, page 99

However, in the case of the *whole* of nature, the idea that nature is a sum of related parts is incorrect. It turns out to be a prejudice, and a prejudice so entrenched that it retarded further developments in physics in the latter decades of the twentieth century. In particular, it does *not* apply to elementary particles or to space-time. Finding the correct description for the whole of nature is the biggest challenge of our adventure, as it requires a complete change in thinking habits. There is a lot of fun ahead.

Jede Aussage über Komplexe läßt sich in eine Aussage über deren Bestandteile und in diejenigen Sätze zerlegen, welche die Komplexe vollständig beschreiben.*

Ludwig Wittgenstein, Tractatus, 2.0201

THE SECOND PRINCIPLE OF THERMODYNAMICS

In contrast to several other important extensive quantities, entropy is *not* conserved. On the one hand, in closed systems, entropy accumulates and never decreases; the sharing or mixing of energy among the components of a system cannot be undone. On the other hand, the sharing or mixing can increase spontaneously over time. Entropy is thus only 'half conserved'. What we call thermal equilibrium is simply the result of the highest possible mixing. Entropy allows us to define the concept of *equilibrium* more precisely as the state of maximum entropy, or maximum energy sharing among the components of a system. In short, the entropy of a closed system increases until it reaches the maximum possible value, the equilibrium value.

The non-conservation of entropy has far-reaching consequences. When a piece of rock is detached from a mountain, it falls, tumbles into the valley, heating up a bit, and eventually stops. The opposite process, whereby a rock cools and tumbles upwards, is never observed. Why? We could argue that the opposite motion does not contradict any rule or pattern about motion that we have deduced so far.

Rocks never fall upwards because mountains, valleys and rocks are made of *many* particles. Motions of many-particle systems, especially in the domain of thermodynamics, are called *processes*. Central to thermodynamics is the distinction between *reversible* processes, such as the flight of a thrown stone, and *irreversible* processes, such as the afore-mentioned tumbling rock. Irreversible processes are all those processes in which friction and its generalizations play a role. Irreversible processes are those processes that increase the sharing or mixing of energy. They are important: if there were no friction, shirt buttons and shoelaces would not stay fastened, we could not walk or run, coffee machines would not make coffee, and maybe most importantly of all, we would have no memory.

Irreversible processes, in the sense in which the term is used in thermodynamics, transform macroscopic motion into the disorganized motion of all the small microscopic components involved: they increase the sharing and mixing of energy. Irreversible processes are therefore not *strictly* irreversible – but their reversal is *extremely* improbable. We can say that entropy measures the 'amount of irreversibility': it measures the degree of mixing or decay that a collective motion has undergone.

Entropy is not conserved. Indeed, entropy – 'heat' – can appear out of nowhere, spontaneously, because energy sharing or mixing can happen by itself. For example, when two different liquids of the same temperature are mixed – such as water and sulphuric acid – the final temperature of the mix can differ. Similarly, when electrical current flows through material at room temperature, the system can heat up or cool down, depending on the material.

Challenge 577 s

Ref. 282

Vol. IV, page 137

^{* &#}x27;Every statement about complexes can be resolved into a statement about their constituents and into the propositions that describe the complexes completely.'

The second principle of thermodynamics states

The entropy in a closed system tends towards its maximum.

In sloppy terms, 'entropy ain't what it used to be.' Here, a closed system is a system that does not exchange energy or matter with its environment. Can you think of an example?

Challenge 578 ny

In a closed system, entropy never decreases. Even everyday life shows us that in a closed system, such as a room, the disorder increases with time, until it reaches some maximum. To reduce disorder, we need effort, i.e., work and energy. In other words, in order to reduce the disorder in a system, we need to connect the system to an energy source in some clever way. For this reason, refrigerators need electrical current or some other energy source.

Ref. 283

In 1866, Ludwig Boltzmann showed that the second principle of thermodynamics results from the principle of least action. Can you imagine and sketch the general ideas?

Challenge 579 ny

Because entropy never decreases in closed systems, white colour does not last. Whenever disorder increases, the colour white becomes 'dirty', usually grey or brown. Perhaps for this reason white objects, such as white clothes, white houses and white underwear, are valued in our society. White objects defy decay.

The second principle implies that heat cannot be transformed to work completely. In other words, every heat engine needs cooling: that is the reason for the holes in the front of cars. The first principle of thermodynamics then states that the mechanical power of a heat engine is the difference between the inflow of thermal energy at high temperature and the outflow of thermal energy at low temperature. If the cooling is insufficient – for example, because the weather is too hot or the car speed too low - the power of the engine is reduced. Every driver knows this from experience.

In summary, the concept of entropy, corresponding to what is called 'heat' in everyday life – but not to what is called 'heat' in physics! – describes the randomness of the internal motion in matter. Entropy is not conserved: in a closed system, entropy never decreases, but it can increase until it reaches a maximum value. The non-conservation of entropy is due to the many components inside everyday systems. The large number of components lead to the non-conservation of entropy and therefore explain, among many other things, that many processes in nature never occur backwards, even though they could do so in principle.

WHY CAN'T WE REMEMBER THE FUTURE?

((It's a poor sort of memory which only works backwards.

Lewis Carroll, Alice in Wonderland

When we first discussed time, we ignored the difference between past and future. But obviously, a difference exists, as we do not have the ability to remember the future. This is not a limitation of our brain alone. All the devices we have invented, such as tape recorders, photographic cameras, newspapers and books, only tell us about the past. Is there a way to build a video recorder with a 'future' button? Such a device would have to

Challenge 580 ny

Challenge 581 ny

solve a deep problem: how would it distinguish between the near and the far future? It does not take much thought to see that any way to do this would conflict with the second principle of thermodynamics. That is unfortunate, as we would need precisely the same device to show that there is faster-than-light motion. Can you find the connection?

In summary, the future cannot be remembered because entropy in closed systems tends towards a maximum. Put even more simply, memory exists because the brain is made of many particles, and so the brain is limited to the past. However, for the most simple types of motion, when only a few particles are involved, the difference between past and future disappears. For few-particle systems, there is no difference between times gone by and times approaching. We could say that the future differs from the past only in our brain, or equivalently, only because of friction. Therefore the difference between the past and the future is not mentioned frequently in this walk, even though it is an essential part of our human experience. But the fun of the present adventure is precisely to overcome our limitations.

FLOW OF ENTROPY

We know from daily experience that transport of an extensive quantity always involves friction. Friction implies generation of entropy. In particular, the flow of entropy itself produces additional entropy. For example, when a house is heated, entropy is produced in the wall. Heating means to keep a temperature difference ΔT between the interior and the exterior of the house. The heat flow J traversing a square metre of wall is given by

$$J = \kappa \Delta T = \kappa (T_{\rm i} - T_{\rm e}) \tag{114}$$

where κ is a constant characterizing the ability of the wall to conduct heat. While conducting heat, the wall also *produces* entropy. The entropy production σ is proportional to the difference between the interior and the exterior entropy flows. In other words, one has

$$\sigma = \frac{J}{T_{\rm e}} - \frac{J}{T_{\rm i}} = \kappa \frac{(T_{\rm i} - T_{\rm e})^2}{T_{\rm i} T_{\rm e}} \ . \tag{115}$$

Note that we have assumed in this calculation that everything is near equilibrium in each slice parallel to the wall, a reasonable assumption in everyday life. A typical case of a good wall has $\kappa = 1 \text{ W/m}^2\text{K}$ in the temperature range between 273 K and 293 K. With this value, one gets an entropy production of

$$\sigma = 5 \cdot 10^{-3} \,\text{W/m}^2\text{K} \,. \tag{116}$$

Challenge 582 ny

Can you compare the amount of entropy that is produced in the flow with the amount that is transported? In comparison, a good goose-feather duvet has $\kappa = 1.5 \, \text{W/m}^2 \text{K}$, which in shops is also called 15 tog.*

^{*} That unit is not as bad as the official (not a joke) BthU \cdot h/sqft/cm/°F used in some remote provinces of our galaxy.

The insulation power of materials is usually measured by the constant $\lambda = \kappa d$ which is independent of the thickness d of the insulating layer. Values in nature range from about 2000 W/K m for diamond, which




FIGURE 223 The basic idea of statistical mechanics about gases: gases are systems of moving particles, and pressure is due to their collisions with the container.

Entropy can be transported in three ways: through *heat conduction*, as just mentioned, via *convection*, used for heating houses, and through *radiation*, which is possible also through empty space. For example, the Earth radiates about 1.2 W/m²K into space, in total thus about 0.51 PW/K. The entropy is (almost) the same that the Earth receives from the Sun. If more entropy had to be radiated away than received, the temperature of the surface of the Earth would have to increase. This is called the *greenhouse effect* or *global warming*. Let's hope that it remains small in the near future.

DO ISOLATED SYSTEMS EXIST?

In all our discussions so far, we have assumed that we can distinguish the system under investigation from its environment. But do such *isolated* or *closed* systems, i.e., systems not interacting with their environment, actually exist? Probably our own human condition was the original model for the concept: we do experience having the possibility to act independently of our environment. An isolated system may be simply defined as a system not exchanging any energy or matter with its environment. For many centuries, scientists saw no reason to question this definition.

The concept of an isolated system had to be refined somewhat with the advent of quantum mechanics. Nevertheless, the concept provides useful and precise descriptions of nature also in that domain. Only in the final part of our walk will the situation change drastically. There, the investigation of whether the universe is an isolated system will lead to surprising results. (What do you think? A strange hint: your answer is almost surely wrong.) We'll take the first steps towards the answer shortly.

Challenge 583 s

CURIOSITIES AND FUN CHALLENGES ABOUT HEAT AND REVERSIBILITY

Compression of air increases its temperature. This is shown directly by the fire pump, a variation of a bicycle pump, shown in Figure 221. (For a working example, see the web page www.de-monstrare.nl). A match head at the bottom of an air pump made of transparent material is easily ignited by the compression of the air above it. The temperature of the air after compression is so high that the match head ignites spontaneously.

is the best conductor of all, down to between 0.1 W/K m and 0.2 W/K m for wood, between 0.015 W/K m and 0.05 W/K m for wools, cork and foams, and the small value of $5 \cdot 10^{-3}$ W/K m for krypton gas.

* *

Running backwards is an interesting sport. The 2006 world records for running backwards can be found on www.recordholders.org/en/list/backwards-running.html. You will be astonished how much these records are faster than your best personal *forward*-running time.

Challenge 584 e

* *

If heat really is disordered motion of atoms, a big problem appears. When two atoms collide head-on, in the instant of smallest distance, neither atom has velocity. Where does the kinetic energy go? Obviously, it is transformed into potential energy. But that implies that atoms can be deformed, that they have internal structure, that they have parts, and thus that they can in principle be split. In short, if heat is disordered atomic motion, atoms are not indivisible! In the nineteenth century this argument was put forward in order to show that heat cannot be atomic motion, but must be some sort of fluid. But since we know that heat really is kinetic energy, atoms must indeed be divisible, even though their name means 'indivisible'. We do not need an expensive experiment to show this.

* *

How long does it take to cook an egg? This issue has been researched in many details; of course, the time depends on what type of cooked egg you want, how large it is, and whether it comes from the fridge or not. There is even a formula for calculating the cooking time! Egg white starts hardening at 62°, the yolk starts hardening at 65°. The best-tasting hard eggs are formed at 69°, half-hard eggs at 65°, and soft eggs at 63°. If you cook eggs at 100° (for a long time) , the white gets the consistency of rubber and the yolk gets a green surface that smells badly, because the high temperature leads to the formation of the smelly H_2S , which then bonds to iron and forms the green FeS. Note that when temperature is controlled, the time plays no role; 'cooking' an egg at 65° for 10 minutes or 10 hours gives the *same* result.

* *

Challenge 585 s Challenge 586 e It is easy to cook an egg in such a way that the white is hard but the yolk remains liquid. Can you achieve the opposite? Research has even shown how you can cook an egg so that the yolk remains at the centre. Can you imagine the method?

* *

Ref. 285

Ref. 284

In 1912, Emile Borel noted that if a gram of matter on Sirius was displaced by one centimetre, it would change the gravitational field on Earth by a tiny amount only. But this tiny change would be sufficient to make it impossible to calculate the path of molecules in a gas after a fraction of a second.

* *

Not only gases, but also most other materials expand when the temperature rises. As a result, the electrical wires supported by pylons hang much lower in summer than in winter. True?

Challenge 587 s




FIGURE 224 Can you boil water in this paper cup?

The following is a famous problem asked by Fermi. Given that a human corpse cools down in four hours after death, what is the minimum number of calories needed per day in our food? Challenge 588 nv

The energy contained in thermal motion is not negligible. A 1 g bullet travelling at the speed of sound has a kinetic energy of only 0.01 kcal.

How does a typical, 1500 m³ hot-air balloon work? Challenge 589 s

If you do not like this text, here is a proposal. You can use the paper to make a cup, as shown in Figure 224, and boil water in it over an open flame. However, to succeed, you have to be a little careful. Can you find out in what way?

Challenge 590 s

Challenge 591 ny

Mixing 1 kg of water at 0°C and 1 kg of water at 100°C gives 2 kg of water at 50°C. What is the result of mixing 1 kg of ice at 0°C and 1 kg of water at 100°C?

The highest recorded air temperature in which a man has survived is 127°C. This was tested in 1775 in London, by the secretary of the Royal Society, Charles Blagden, together with a few friends, who remained in a room at that temperature for 45 minutes. Interestingly, the raw steak which he had taken in with him was cooked ('well done') when he and his friends left the room. What condition had to be strictly met in order to avoid

cooking the people in the same way as the steak? Challenge 592 s

The Swedish astronomer Anders Celsius (1701–1744) originally set the freezing point of water at 100 degrees and the boiling point at 0 degrees. Shortly afterwards, the scale was reversed to the one in use now. However, this is not the whole story. With the official definition of the kelvin and the degree Celsius, at the standard pressure of 101 325 Pa, water boils at 99.974°C. Can you explain why it is not 100°C any more?

Ref. 288

Challenge 593 s




FIGURE 225 The invisible loudspeaker.

* *

Challenge 594 s Can you fill a bottle precisely with 1 ± 10^{-30} kg of water?

* *

One gram of fat, either butter or human fat, contains 38 kJ of chemical energy (or, in ancient units more familiar to nutritionists, 9 kcal). That is the same value as that of petrol. Why are people and butter less dangerous than petrol?

Challenge 595 s

Challenge 596 s

* *

In 1992, the Dutch physicist Martin van der Mark invented a loudspeaker which works by heating air with a laser beam. He demonstrated that with the right wavelength and with a suitable modulation of the intensity, a laser beam in air can generate sound. The effect at the basis of this device, called the *photoacoustic effect*, appears in many materials. The best laser wavelength for air is in the infrared domain, on one of the few absorption lines of water vapour. In other words, a properly modulated infrared laser beam that shines through the air generates sound. Such light can be emitted from a small matchbox-sized semiconductor laser hidden in the ceiling and shining downwards. The sound is emitted in all directions perpendicular to the beam. Since infrared laser light is not visible, Martin van der Mark thus invented an invisible loudspeaker! Unfortunately, the efficiency of present versions is still low, so that the power of the speaker is not yet sufficient for practical applications. Progress in laser technology should change this, so that in the future we should be able to hear sound that is emitted from the centre of an otherwise empty room.

* *

A famous exam question: How can you measure the height of a building with a barometer, a rope and a ruler? Find at least six different ways.

* *

What is the approximate probability that out of one million throws of a coin you get

Challenge 597 ny

exactly 500 000 heads and as many tails? You may want to use Stirling's formula $n! \approx \sqrt{2\pi n} (n/e)^n$ to calculate the result.*

* *

Challenge 598 s Does it make sense to talk about the entropy of the universe?

* *

Challenge 599 ny Can a helium balloon lift the tank which filled it?

* *

All friction processes, such as osmosis, diffusion, evaporation, or decay, are *slow*. They take a characteristic time. It turns out that any (macroscopic) process with a time-scale is irreversible. This is no real surprise: we know intuitively that undoing things always takes more time than doing them. That is again the second principle of thermodynamics.

* *

Ref. 289

It turns out that *storing* information is possible with negligible entropy generation. However, *erasing* information requires entropy. This is the main reason why computers, as well as brains, require energy sources and cooling systems, even if their mechanisms would otherwise need no energy at all.

* *

Challenge 600 ny

When mixing hot rum and cold water, how does the increase in entropy due to the mixing compare with the entropy increase due to the temperature difference?

* *

Challenge 601 s

Why aren't there any small humans, say 10 mm in size, as in many fairy tales? In fact, there are no warm-blooded animals of that size at all. Why not?

* *

Shining a light onto a body and repeatedly switching it on and off produces sound. This is called the *photoacoustic effect*, and is due to the thermal expansion of the material. By changing the frequency of the light, and measuring the intensity of the noise, one reveals a characteristic photoacoustic spectrum for the material. This method allows us to detect gas concentrations in air of one part in 10^9 . It is used, among other methods, to study the gases emitted by plants. Plants emit methane, alcohol and acetaldehyde in small quantities; the photoacoustic effect can detect these gases and help us to understand the processes behind their emission.

* *

Challenge 602 ny

What is the rough probability that all oxygen molecules in the air would move away from a given city for a few minutes, killing all inhabitants?

^{*} There are many improvements to Stirling's formula. A simple one is Gosper's formula $n! \approx \sqrt{(2n+1/3)\pi}(n/e)^n$. Another is $\sqrt{2\pi n}(n/e)^n e^{1/(12n+1)} < n! < \sqrt{2\pi n}(n/e)^n e^{1/(12n)}$.

* *

If you pour a litre of water into the sea, stir thoroughly through all the oceans and then Challenge 603 ny take out a litre of the mixture, how many of the original atoms will you find?

* *

Challenge 604 ny How long would you go on breathing in the room you are in if it were airtight?

* *

What happens if you put some ash onto a piece of sugar and set fire to the whole? Challenge 605 ny (Warning: this is dangerous and not for kids.)

* *

Entropy calculations are often surprising. For a system of N particles with two states each, there are $W_{\rm all}=2^N$ states. For its most probable configuration, with exactly half the particles in one state, and the other half in the other state, we have $W_{\rm max}=N!/((N/2)!)^2$. Now, for a macroscopic system of particles, we might typically have $N=10^{24}$. That gives $W_{\rm all}\gg W_{\rm max}$; indeed, the former is 10^{12} times larger than the latter. On the other hand, we find that $\ln W_{\rm all}$ and $\ln W_{\rm max}$ agree for the first 20 digits! Even though the configuration with exactly half the particles in each state is much more rare than the general case, where the ratio is allowed to vary, the entropy turns out to be the same. Why?

* *

If heat is due to motion of atoms, our built-in senses of heat and cold are simply detectors of motion. How could they work?

By the way, the senses of smell and taste can also be seen as motion detectors, as they signal the presence of molecules flying around in air or in liquids. Do you agree?

* *

The Moon has an atmosphere, although an extremely thin one, consisting of sodium (Na) and potassium (K). This atmosphere has been detected up to nine Moon radii from its surface. The atmosphere of the Moon is generated at the surface by the ultraviolet radiation from the Sun. Can you estimate the Moon's atmospheric density?

* *

Does it make sense to add a line in Table 43 for the quantity of physical action? A column? Why?

* *

Diffusion provides a length scale. For example, insects take in oxygen through their skin. As a result, the interiors of their bodies cannot be much more distant from the surface than about a centimetre. Can you list some other length scales in nature implied by diffusion processes?

* *

Rising warm air is the reason why many insects are found in tall clouds in the evening. Many insects, especially that seek out blood in animals, are attracted to warm and humid

Challenge 606 ny

Challenge 607 ny


Challenge 608 ny

Challenge 609 ny

Challenge 610 s

Challenge 611 ny

Challenge 612 s






FIGURE 226 The design of the Wirbelrohr or Ranque–Hilsch vortex tube, and a commercial version, about 40 cm in size, used to cool manufacturing processes (© Coolquip).

air.

* *

Thermometers based on mercury can reach 750°C. How is this possible, given that mercury boils at 357°C?

Challenge 613 s

* *

Challenge 614 s What does a burning candle look like in weightless conditions?

* *

It is possible to build a power station by building a large chimney, so that air heated by the Sun flows upwards in it, driving a turbine as it does so. It is also possible to make a power station by building a long vertical tube, and letting a gas such as ammonia rise into it which is then liquefied at the top by the low temperatures in the upper atmosphere; as it falls back down a second tube as a liquid – just like rain – it drives a turbine. Why are such schemes, which are almost completely non-polluting, not used yet?

Challenge 615 s

* *

One of the most surprising devices ever invented is the *Wirbelrohr* or Ranque–Hilsch vortex tube. By blowing compressed air at room temperature into it at its midpoint, two flows of air are formed at its ends. One is extremely cold, easily as low as –50°C, and one extremely hot, up to 200°C. No moving parts and no heating devices are found inside. How does it work?

Challenge 616 s

* *

Thermoacoustic engines, pumps and refrigerators provide many strange and fascinating applications of heat. For example, it is possible to use loud sound in closed metal chambers to move heat from a cold place to a hot one. Such devices have few moving parts and are being studied in the hope of finding practical applications in the future.

Ref. 290

* *

Does a closed few-particle system contradict the second principle of thermodynamics?

Challenge 617 ny






FIGURE 227 What happens to the ink stripe if the inner cylinder is turned a few times in one direction, and then turned back by the same amount?

* *

What happens to entropy when gravitation is taken into account? We carefully left gravitation out of our discussion. In fact, gravitation leads to many new problems – just try to think about the issue. For example, Jacob Bekenstein has discovered that matter reaches its highest possible entropy when it forms a black hole. Can you confirm this?

Challenge 618 ny

* *

The numerical values (but not the units!) of the Boltzmann constant $k = 1.38 \cdot 10^{-23}$ J/K and the combination h/ce agree in their exponent and in their first three digits, where h is Planck's constant and e is the electron charge. Can you dismiss this as mere coincidence?

Challenge 619 ny

* *

Mixing is not always easy to perform. The experiment of Figure 227 gives completely different results with water and glycerine. Can you guess them?

Challenge 620 s

* *

Challenge 621 s How do you get rid of chewing gum in clothes?

* *

Challenge 622 ny

There are less-well known arguments about atoms. In fact, two everyday prove the existence of atoms: reproduction and memory. Why?

Challenge 623 s Why is this not really sensible?

In the context of lasers and of spin systems, it is fun to talk about negative temperature.

Summary on heat and time-invariance

Microscopic motion due to gravity and electric interactions, thus all microscopic motion in everyday life, is *reversible*: such motion can occur backwards in time. In other words, motion due to gravity and electromagnetism is time-reversal-invariant or motion-reversal-invariant.

Nevertheless, everyday motion is *irreversible*, because there are no completely closed systems in everyday life. Lack of closure leads to fluctuations; fluctuations lead to friction. Equivalently, irreversibility results from the extremely low probabilities required to perform a motion inversion. Macroscopic irreversibility does not contradict microscopic reversibility.

For these reasons, in everyday life, entropy in closed systems always increases. This leads to a famous issue: how can biological evolution be reconciled with entropy increase? Let us have a look.



SELF-ORGANIZATION AND CHAOS THE SIMPLICITY OF COMPLEXITY

To speak of non-linear physics is like calling zoology the study of non-elephant animals. Stanislaw Ulam

Ref. 291

Page 208

Ref. 292

Challenge 624 s

N our list of global descriptions of motion, the the study of self-organization s the high point. Self-organization is the appearance of order. In physics, order s a term that includes shapes, such as the complex symmetry of snowflakes; patterns, such as the stripes of zebras and the ripples on sand; and cycles, such as the creation of sound when singing. When we look around us, we note that every example of what we call beauty is a combination of shapes, patterns and cycles. (Do you agree?) Selforganization can thus be called the study of the origin of beauty. Table 48 shows how frequently the appearance of order shapes our environment.

TABLE 48 Some rhythms, patterns and shapes observed in nature.

OBSERVATION	DRIVING 'FORCE'	'RESTORING FORCE'	TYP. SCALE
Fingerprint	chemical reactions	diffusion	0.1 mm
Clock ticking	falling weight	friction	1 s
Chalk squeaking due to stick-slip instability	motion	friction	600 Hz
Musical note generation in violin	bow motion	friction	600 Hz
Musical note generation in flute	air flow	turbulence	400 Hz
Train oscillations transversally to the track	motion	friction	0.3 Hz
Flow structures in waterfalls and fountains	water flow	turbulence	10 cm
Jerky detachment of scotch tape	pulling speed	sticking friction	0.1 Hz
Radius oscillations in spaghetti and polymer fibre production	extrusion speed	friction	10 cm
Patterns on buckled metal plates and foils	deformation	stiffness	depend on thickness

Patterns in laser irradiated laser irradiation

Cd-Se alloys

50 µm

OBSERVATION	DRIVING 'FORCE'	'RESTORING FORCE'	Typ. Scale
Flapping of flags in steady wind	air flow	stiffness	20 cm
Dripping of water tap	water flow	surface tension	1 Hz
Bubble stream from a beer glass irregularity	dissolved gas pressure	surface tension	0.1 Hz, 1 mm
Raleigh–Bénard instability	temperature gradient	diffusion	0.1 Hz, 1 mm
Couette-Taylor flow	speed gradient	friction	0.1 Hz, 1 mm
Bénard–Marangoni flow, sea wave generation			0.1 Hz, 1 mm
Karman wakes, Emmon spots, Osborne-Reynolds flow			
Regular bangs in a car exhaustion pipe	flow	pressure resonances	0.3 Hz
Regular cloud arrangements	flow	diffusion	0.5 km
El Niño	flow	diffusion	5 to 7 years
Wine arcs on glass walls	surface tension	binary mixture	0.1 Hz, 1 mm
Ferrofluids surfaces in magnetic fields	magnetic energy	gravity	3 mm
Patterns in liquid crystals	electric energy	stress	1 mm, 3 s
Flickering of aging fluorescence lights	electron flow	diffusion	1 Hz
Surface instabilities of welding	electron flow	diffusion	1 cm
Tokamak plasma instabilities	electron flow	diffusion	10 s
Snowflake formation and other dendritic growth processes	concentration gradient	surface diffusion	10 μm
Solidification interface patterns, e.g. in CBr ₄	entropy flow	1 mm	
Periodic layers in metal corrosion	concentration gradients	diffusion	10 μm
Hardening of steel by cold working	strain	dislocation motion	5 μm
Labyrinth structures in proton irradiated metals	particle flow	dislocation motion	5 μm

diffusion

TABLE 48 (Continued) Some rhytms, patterns and shapes observed in nature.

TABLE 48 (Continued) Some rhytms, patterns and shapes observed in nature.

O B S E R V A T I O N	DRIVING 'FORCE'	'RESTORING FORCE'	TYP. SCALE
Dislocation patterns and density oscillations in fatigued Cu single crystals	strain	dislocation motion	10 μm 100 s
Laser light emission, its cycles and chaotic regimes	pumping energy	light losses	10 ps to 1 ms
Rotating patterns from shining laser light on the surface of certain electrolytes	light energy	diffusion	1 mm
Belousov-Zhabotinski reaction patterns and cycles	concentration gradients	diffusion	1 mm, 10 s
Flickering of a burning candle	heat and concentration gradients	thermal and substance diffusion	0.1 s
Regular sequence of hot and cold flames in carbohydrate combustion	heat and concentration gradients	thermal and substance diffusion	1 cm
Feedback whistle from microphone to loudspeaker	amplifiers	electric losses	1 kHz
Any electronic oscillator in radio sets, television sets, computers, mobile phones, etc.	power supply	resistive losses	1 kHz to 30 GHz
Periodic geysir eruptions	underground heating	evaporation	10 min
Periodic earthquakes at certain faults	tectonic motion	ruptures	1 Ms
Hexagonal patterns in basalt rocks	heating	heat diffusion	1 m
Hexagonal patterns on dry soil	regular temperature changes	water diffusion	0.5 m
Periodic intensity changes of the Cepheids and other stars	nuclear fusion	energy emission	3 Ms
Convection cells on the surface of the Sun	nuclear fusion	energy emission	1000 km
Formation and oscillations of the magnetic field of the Earth and other celestial bodies		resistive losses	100 ka
Wrinkling/crumpling transition	strain	stiffness	1 mm

TYP. SCALE OBSERVATION DRIVING 'RESTORING FORCE FORCE' Patterns of animal furs chemical diffusion 1 cm concentration Growth of fingers and chemical diffusion 1 cm limbs concentration probably molecular diffusion Symmetry breaking in 1 m embryogenesis, such as the chirality plus heart on the left chemical concentration Cell differentiation and chemical diffusion $10 \, \mu m$ to $30 \, m$ appearance of organs concentration during growth Prey-predator oscillations reproduction hunger 3 to 17 a

TABLE 48 (Continued) Some rhytms, patterns and shapes observed in nature.

neuron firing

APPEARANCE OF ORDER

Thinking

The appearance of order is a general observation across nature. *Fluids* in particular exhibit many phenomena where order appears and disappears. Examples include the more or less regular flickering of a burning candle, the flapping of a flag in the wind, the regular stream of bubbles emerging from small irregularities in the surface of a champagne glass, and the regular or irregular dripping of a water tap. Figure 202 shows some additional examples, and so do the figures in this chapter. Other examples include the appearance of clouds and of regular cloud arrangements in the sky. It can be fascinating to ponder, during an otherwise boring flight, the mechanisms behind the formation of the cloud shapes and patterns you see from the aeroplane.

heat dissipation

 $1 \text{ ms}, 100 \, \mu \text{m}$

Challenge 625 e

Page 303

Other cases of self-organization are mechanical, such as the formation of mountain ranges when continents move, the creation of earthquakes, or the formation of laughing fold at the corners of human eyes.

All *growth* processes are self-organization phenomena. The appearance of order is found from the cell differentiation in an embryo inside a woman's body; the formation of colour patterns on tigers, tropical fish and butterflies; the symmetrical arrangements of flower petals; the formation of biological rhythms; and so on.

Page 45

Have you ever pondered the incredible way in which teeth grow? A practically inorganic material forms shapes in the upper and the lower rows fitting exactly into each other. How this process is controlled is still a topic of research. Also the formation, before and after birth, of neural networks in the brain is another process of self-organization. Even the physical processes at the basis of thinking, involving changing electrical signals, is to be described in terms of self-organization.

Vol. III, page 187

Biological evolution is a special case of growth. Take the evolution of animal shapes. It turns out that snake tongues are forked because that is the most efficient shape for following chemical trails left by prey and other snakes of the same species. (Snakes smell with the help of their tongue.) The fixed numbers of fingers in human hands or of petals

Ref. 293

funnel



Vol. III, page 334 of flowers are also consequences of self-organization.

Studies into the conditions required for the appearance or disappearance of order have shown that their description requires only a few common concepts, independently of the details of the physical system. This is best seen looking at a few simple examples.

SELF-ORGANIZATION IN SAND

All the richness of self-organization reveals itself in the study of plain sand. Why do sand dunes have ripples, as does the sand floor at the bottom of the sea? How do avalanches occur on steep heaps of sand? How does sand behave in hourglasses, in mixers, or in vibrating containers? The results are often surprising.

totion Mountain – The Adventure of Physics 🏻 pdf file available free of charge at www.motionmountain.net 🖯 Copyright © Christoph Schiller November 1997–March

TABLE 49 Patterns and a cycle on horizontal sand and on sand-like surfaces in the sea and on land.

PATTERN/CYC	CLE PERIOD	AMPLITUDE	ORIGIN
Under water			
Ripples	5 cm	5 mm	water waves
Megaripples	1 m	0.1 m	tides
Sand waves	100 to 800 m	5 m	tides
Sand banks	2 to 10 km	2 to 20 m	tides
In air			
Ripples	0.1 m	0.05 m	wind
Singing sand	65 to 110 Hz	up to 105 dB	wind on sand dunes, avalanches making the dune vibrate
Road corrugations	0.3 to 0.9 m	0.05 m	wheels
Ski moguls	5 to 6 m	up to 1 m	skiers
Elsewhere			
On Mars	a few km	few tens of m	wind



Ref. 295

FIGURE 229 Road corrugations (courtesy David Mays).

An overview of self-organization in sand is given in Table 49. For example, as recently as 2006, the Cuban research group of Ernesto Altshuler and his colleagues discovered solitary waves on sand flows (shown in Figure 228). They had already discovered the revolving river effect on sand piles, shown in the same figure, in 2002. Even more surprisingly, these effects occur only for Cuban sand, and a few rare other types of sand. The reasons are still unclear.

Similarly, in 1996 Paul Umbanhowar and his colleagues found that when a flat container holding tiny bronze balls (around 0.165 mm in diameter) is shaken up and down in vacuum at certain frequencies, the surface of this bronze 'sand' forms stable heaps.



FIGURE 230 Oscillons formed by shaken bronze balls; horizontal size is about 2 cm (© Paul Umbanhowar)




FIGURE 231 Magic numbers: 21 spheres, when swirled in a dish, behave differently from non-magic numbers, like 23, of spheres (redrawn from photographs © Karsten Kötter).

They are shown in Figure 230. These heaps, so-called *oscillons*, also bob up and down. The oscillons can move and interact with one another.

Oscillons in bronze sand are a simple example for a general effect in nature: *discrete* systems with nonlinear interactions can exhibit *localized excitations*. This fascinating topic is just beginning to be researched. It might well be that one day it will yield results relevant to our understanding of the growth of organisms.

Sand shows many other pattern-forming processes.

- A mixture of sand and sugar, when poured onto a heap, forms regular layered structures that in cross section look like zebra stripes.
- Horizontally rotating cylinders with binary mixtures inside them separate the mixture out over time.
- Take a container with two compartments separated by a 1 cm wall. Fill both halves with sand and rapidly shake the whole container with a machine. Over time, all the sand will spontaneously accumulate in one half of the container.
- In sand, people have studied the various types of sand dunes that 'sing' when the wind blows over them.
- Also the corrugations formed by traffic on roads without tarmac, the washboard roads shown in Figure 229, are an example of self-organization. These corrugation patterns move, over time, *against* the traffic direction. Can you explain why? The moving ski moguls mentioned above also belong here.

In fact, the behaviour of sand and dust is proving to be such a beautiful and fascinating topic that the prospect of each human returning to dust does not look so grim after all.

Ref. 297

Challenglee 6 2298
Page 271



FIGURE 232 Self-organization: a growing snow flake (QuickTime film © Kenneth Libbrecht)

SELF-ORGANIZATION OF SPHERES

A stunningly simple and beautiful example of self-organization is the effect discovered in 1999 by Karsten Kötter and his group. They found that the behaviour of a set of spheres swirled in a dish depends on the number of spheres used. Usually, all the spheres get continuously mixed up. But for certain 'magic' numbers, such as 21, stable ring patterns emerge, for which the outside spheres remain outside and the inside ones remain inside. The rings, best seen by colouring the spheres, are shown in Figure 231.

APPEARANCE OF ORDER

The many studies of self-organizing systems have changed our understanding of nature in a number of ways. First of all, they have shown that patterns and shapes are similar to cycles: all are due to motion. Without motion, and thus without history, there is no order, neither patterns nor shapes nor rhythms. Every pattern has a history; every pattern is a result of *motion*. As an example, Figure 232 shows how a snowflake grows.

Secondly, patterns, shapes and rhythms are due to the organized motion of large numbers of small constituents. Systems which self-organize are always composite: they are *cooperative structures*.

Thirdly, all these systems obey evolution equations which are *nonlinear* in the macroscopic configuration variables. Linear systems do not self-organize.

Fourthly, the appearance and disappearance of order depends on the strength of a driving force, the so-called *order parameter*. Moreover, all order and all structure appears when two general types of motion compete with each other, namely a 'driving', energy-adding process, and a 'dissipating', braking mechanism. Thermodynamics thus plays a role in all self-organization. Self-organizing systems are always *dissipative systems*, and are always far from equilibrium. When the driving and the dissipation are of the same order of magnitude, and when the key behaviour of the system is not a linear function of the driving action, order may appear.*

^{*} To describe the 'mystery' of human life, terms like 'fire', 'river' or 'tree' are often used as analogies. These are all examples of self-organized systems: they have many degrees of freedom, have competing driving

THE MATHEMATICS OF ORDER APPEARANCE

Every pattern, every shape and every rhythm or cycle can be described by some observable *A* that describes the amplitude of the pattern, shape or rhythm. For example, the amplitude *A* can be a length for sand patterns, or a chemical concentration for biological systems, or a sound pressure for sound appearance.

Order appears when the amplitude *A* differs from zero. To understand the appearance of order, one has to understand the evolution of the amplitude *A*. The study of order has shown that this amplitude always follows similar evolution equations, *independently* of the physical mechanism of system. This surprising result unifies the whole field of self-organization.

All self-organizing systems at the onset of order appearance can be described by equations for the pattern amplitude *A* of the general form

$$\frac{\partial A(t,x)}{\partial t} = \lambda A - \mu |A|^2 A + \kappa \Delta A + \text{higher orders}.$$
 (117)

Here, the observable A – which can be a real or a complex number, in order to describe phase effects – is the observable that appears when order appears, such as the oscillation amplitude or the pattern amplitude. The first term λA is the *driving term*, in which λ is a parameter describing the strength of the driving. The next term is a typical *nonlinearity* in A, with μ a parameter that describes its strength, and the third term $\kappa \Delta A = \kappa (\partial^2 A/\partial x^2 + \partial^2 A/\partial y^2 + \partial^2 A/\partial z^2)$ is a typical diffusive and thus *dissipative* term.

We can distinguish two main situations. In cases where the dissipative term plays no role ($\kappa=0$), we find that when the driving parameter λ increases above zero, a *temporal* oscillation appears, i.e., a stable cycle with non-vanishing amplitude. In cases where the diffusive term does play a role, equation (117) describes how an amplitude for a *spatial* oscillation appears when the driving parameter λ becomes positive, as the solution A=0 then becomes spatially unstable.

In both cases, the onset of order is called a *bifurcation*, because at this critical value of the driving parameter λ the situation with amplitude zero, i.e., the homogeneous (or unordered) state, becomes unstable, and the ordered state becomes stable. *In nonlinear systems, order is stable*. This is the main conceptual result of the field. Equation (117) and its numerous variations allow us to describe many phenomena, ranging from spirals, waves, hexagonal patterns, and topological defects, to some forms of turbulence. For every physical system under study, the main task is to distil the observable A and the parameters λ , μ and κ from the underlying physical processes.

Self-organization is a vast field which is yielding new results almost by the week. To discover new topics of study, it is often sufficient to keep one's eye open; most effects are comprehensible without advanced mathematics. Enjoy the hunting!

Challenge 627 ny

Challenge 628 ny

Ref. 30

Challenge 629 nv

and braking forces, depend critically on their initial conditions, show chaos and irregular behaviour, and sometimes show cycles and regular behaviour. Humans and human life resemble them in all these respects; thus there is a solid basis to their use as metaphors. We could even go further and speculate that pure beauty *is* pure self-organization. The lack of beauty indeed often results from a disturbed equilibrium between external braking and external driving.




FIGURE 233 Examples of different types of motion in configuration space.

CHAOS

Most systems that show self-organization also show another type of motion. When the driving parameter of a self-organizing system is increased to higher and higher values, order becomes more and more irregular, and in the end one usually finds *chaos*. For physicists, $c^{\text{N}}a^{\text{O}}T$, c motion is the most irregular type of motion.* Chaos can be

For physicists, $c^{\infty}a^{\circ}1$. c motion is the most irregular type of motion.* Chaos can be defined independently of self-organization, namely as that motion of systems for which small changes in initial conditions evolve into large changes of the motion (exponentially with time). This is illustrated in Figure 234. More precisely, *chaos* is irregular motion characterized by a positive *Lyapounov exponent* in the presence of a strictly valid evolution.

A simple chaotic system is the damped pendulum above three magnets. Figure 235 shows how regions of predictability (around the three magnet positions) gradually change into a chaotic region, i.e., a region of effective unpredictability, for higher initial amplitudes. The weather is also a chaotic system, as are dripping water-taps, the fall of dice, and many other everyday systems. For example, research on the mechanisms by which the heart beat is generated has shown that the heart is not an oscillator, but a chaotic system with irregular cycles. This allows the heart to be continuously ready for demands for changes in beat rate which arise once the body needs to increase or decrease its efforts.

Ref. 302

Challenge 630 ny

There is chaotic motion also in machines: chaos appears in the motion of trains on the rails, in gear mechanisms, and in fire-fighter's hoses. The precise study of the motion in a zippo cigarette lighter will probably also yield an example of chaos. The mathematical description of chaos – simple for some textbook examples, but extremely involved for others – remains an important topic of research.

Incidentally, can you give a simple argument to show that the so-called *butterfly effect* does not exist? This 'effect' is often cited in newspapers. The claim is that nonlinearities

Challenge 631 s

^{*} On the topic of chaos, see the beautiful book by H.-O. Peitgen, H. Jürgens & D. Saupe, *Chaos and Fractals*, Springer Verlag, 1992. It includes stunning pictures, the necessary mathematical background, and some computer programs allowing personal exploration of the topic. 'Chaos' is an old word: according to Greek mythology, the first goddess, Gaia, i.e., the Earth, emerged from the chaos existing at the beginning. She then gave birth to the other gods, the animals and the first humans.



initial condition 1
initial condition 2
time

FIGURE 234 Chaos as sensitivity to initial conditions.




FIGURE 235 A simple chaotic system: a metal pendulum over three magnets (fractal © Paul Nylander).

imply that a small change in initial conditions can lead to large effects; thus a butterfly wing beat is alleged to be able to induce a tornado. Even though nonlinearities do indeed lead to growth of disturbances, the butterfly 'effect' has *never* been observed. Thus it does *not* exist. This 'effect' exists only to sell books and to get funding.

All the steps from disorder to order, quasiperiodicity and finally to chaos, are examples of self-organization. These types of motion, illustrated in Figure 233, are observed in many fluid systems. Their study should lead, one day, to a deeper understanding of the mysteries of turbulence. Despite the fascination of this topic, we will not explore it further, because it does not lead towards the top of Motion Mountain.

EMERGENCE

Self-organization is of interest also for a more general reason. It is sometimes said that our ability to formulate the patterns or rules of nature from observation does not imply the ability to predict *all* observations from these rules. According to this view, so-called 'emergent' properties exist, i.e., properties appearing in complex systems as something *new* that cannot be deduced from the properties of their parts and their interactions. (The ideological backdrop to this view is obvious; it is the latest attempt to fight the idea of determinism.) The study of self-organization has definitely settled this debate. The

properties of water molecules do allow us to predict Niagara Falls.* Similarly, the diffusion of signal molecules do determine the development of a single cell into a full human being: in particular, cooperative phenomena determine the places where arms and legs are formed; they ensure the (approximate) right–left symmetry of human bodies, prevent mix-ups of connections when the cells in the retina are wired to the brain, and explain the fur patterns on zebras and leopards, to cite only a few examples. Similarly, the mechanisms at the origin of the heart beat and many other cycles have been deciphered.

Self-organization provides general principles which allow us in principle to predict the behaviour of complex systems of any kind. They are presently being applied to the most complex system in the known universe: the human brain. The details of how it learns to coordinate the motion of the body, and how it extracts information from the images in the eye, are being studied intensely. The ongoing work in this domain is fascinating. (A neglected case of self-organization is *humour*.) If you plan to become a scientist, consider taking this path.

Ref. 305 Challenge 633 ny

Self-organization research provided the final arguments that confirmed what J. Offrey de la Mettrie stated and explored in his famous book *L'homme machine* in 1748: humans are complex machines. Indeed, the lack of understanding of complex systems in the past was due mainly to the restrictive teaching of the subject of motion, which usually concentrated – as we do in this walk – on examples of motion in *simple* systems. The concepts of self-organization allow us to understand and to describe what happens during the functioning and the growth of organisms.

Even though the subject of self-organization provides fascinating insights, and will do so for many years to come, we now leave it. We continue with our own adventure, namely to explore the *basics* of motion.

(Ich sage euch: man muss noch Chaos in sich haben, um einen tanzenden Stern gebären zu können. Ich sage euch: ihr habt noch Chaos in euch.**

Friedrich Nietzsche, Also sprach Zarathustra.

CURIOSITIES AND FUN CHALLENGES ABOUT SELF-ORGANIZATION

Ref. 306 Challenge 634 ny All icicles have a wavy surface, with a crest-to-crest distance of about 1 cm, as shown in Figure 236. The distance is determined by the interplay between water flow and surface cooling. How? (Indeed, stalagtites do not show the effect.)

* *

When a fine stream of water leaves a water tap, putting a finger in the stream leads to a Challenge 635 ny wavy shape, as shown in Figure 237. Why?

* *

The research on sand has shown that it is often useful to introduce the concept of granular

Ref. 304 Challenge 632 ny * Already small versions of Niagara Falls, namely dripping water taps, show a large range of cooperative phenomena, including the chaotic, i.e., non-periodic, fall of water drops. This happens when the water flow has the correct value, as you can verify in your own kitchen. Several cooperative fluid phenomena have been simulated even on the molecular level.

^{** &#}x27;I tell you: one must have chaos inside oneself, in order to give birth to a dancing star. I tell you: you still have chaos inside you.'






FIGURE 236 The wavy surface of icicles.




FIGURE 237 Water pearls.



FIGURE 238 A braiding water stream (© Vakhtang Putkaradze).

temperature, which quantifies how fast a region of sand moves. Research into this field is still in full swing.

* *

When water emerges from a oblong opening, the stream forms a braid pattern, as shown in Figure 238. This effect results from the interplay and competition between inertia and surface tension: inertia tends to widen the stream, while surface tension tends to narrow it. Predicting the distance from one narrow region to the next is still a topic of research.

If the experiment is done in free air, without a plate, one usually observes an additional effect: there is a *chiral* braiding at the narrow regions, induced by the asymmetries of the water flow. You can observe this effect in the toilet! Scientific curiosity knows no limits: are you a right-turner or a left-turner, or both? On every day?

Challenge 636 e

Ref. 307

* *

When wine is made to swirl in a wine glass, after the motion has calmed down, the wine flowing down the glass walls forms little arcs. Can you explain in a few words what forms them?

Challenge 637 ny

Challenge 638 ny

* *

How does the average distance between cars parked along a street change over time, assuming a constant rate of cars leaving and arriving?

* *



FIGURE 239 The Belousov-Zhabotinski reaction: the liquid periodically changes colour, both in space and time (© Yamaguchi University).

A famous case of order appearance is the *Belousov-Zhabotinski reaction*. This mixture of chemicals spontaneously produces spatial and temporal patterns. Thin layers produce slowly rotating spiral patterns, as shown in Figure 239; Large, stirred volumes oscillate back and forth between two colours. A beautiful movie of the oscillations can be found on www.uni-r.de/Fakultaeten/nat_Fak_IV/Organische_Chemie/Didaktik/Keusch/D-oscill-d.htm. The exploration of this reaction led to the Nobel Prize in Chemistry for Ilya Prigogine in 1997.

* *

Gerhard Müller has discovered a simple but beautiful way to observe self-organization in solids. His system also provides a model for a famous geological process, the formation of hexagonal columns in basalt, such as the Giant's Causeway in Northern Ireland. Similar formations are found in many other places of the Earth. Just take some rice flour or corn starch, mix it with about half the same amount of water, put the mixture into a pan and dry it with a lamp: hexagonal columns form. The analogy with basalt structures is possible because the drying of starch and the cooling of lava are diffusive processes governed by the same equations, because the boundary conditions are the same, and because both materials respond to cooling with a small reduction in volume.


* *

Water flow in pipes can be laminar (smooth) or turbulent (irregular and disordered). The transition depends on the diameter d of the pipe and the speed v of the water. The transition usually happens when the so-called *Reynolds number* – defined as $R = vd/\eta$ (η being the *kinematic viscosity* of the water, around $1 \text{ mm}^2/\text{s}$) – becomes greater than about 2000. (The *dynamic viscosity* μ is defined as $\mu = \eta \rho$, where ρ is the density of the fluid.) However, careful experiments show that with proper handling, laminar flows can be produced up to $R = 100\,000$. A linear analysis of the equations of motion of the fluid, the Navier–Stokes equations, even predicts stability of laminar flow for *all* Reynolds numbers. This riddle was solved only in the years 2003 and 2004. First, a complex mathematical analysis showed that the laminar flow is not always stable, and that the transition to turbulence in a long pipe occurs with travelling waves. Then, in 2004, careful experi-

Ref. 308 Challenge 639 e

Ref. 309

Ref. 310






FIGURE 240 A famous correspondence: on the left, hexagonal columns in starch, grown in a kitchen pan (the red lines are 1 cm in length), and on the right, hexagonal columns in basalt, grown from lava in Northern Ireland (top right, view of around 300 m, and middle right, view of around 40 m) and in Iceland (view of about 30 m, bottom right) (© Gerhard Müller, Raphael Kessler - www.raphaelk.co.uk, Bob Pohlad, and Cédric Hüsler).

ments showed that these travelling waves indeed appear when water is flowing through a pipe at large Reynolds numbers.

* *

For more beautiful pictures on self-organization in fluids, see the mentioned serve.me. nus.edu.sg/limtt website.

* *

Also *dance* is an example of self-organization. This type of self-organization takes place in the brain. Like for all complex movements, learning them is often a challenge. Nowadays

suntain − The Adventure of Physics | pdf file available free of charge at www.motionmountain.net | Copyright © Christoph Schiller November 1997–March 2012




FIGURE 241 How the shape of snow crystals depend on temperature and saturation (© Kenneth Libbrecht).

Ref. 311 there are beautiful books that tell how physics can help you improve your dancing skills and the grace of your movements.

* *

Do you want to enjoy working on your PhD? Go into a scientific toy shop, and look for any toy that moves in a complex way. There are high chances that the motion is chaotic; explore the motion and present a thesis about it. For example, go to the extreme: explore the motion of a hanging rope whose upper end is externally driven. This simple system is fascinating in its range of complex motion behaviours.

* *

Self-organization is also observed in liquid corn starch-water mixtures. Enjoy the film at www.youtube.com/watch?v=f2XQ97XHjVw and watch even more bizarre effects, for humans walking over a pool filled with the liquid, on www.youtube.com/watch? v=nq3ZjY0Uf-g.

* *

Snow flakes and snow crystals have already been mentioned as examples of selforganization. Figure 241 shows the general connection. To learn more about this fasciRef. 313

Ref. 314

Ref. 315

nating topic, explore the wonderful website snowcrystals.com by Kenneth Libbrecht. A complete classification of snow crystals has also been developed.

* *

A famous example of self-organization whose mechanisms are not well-known so far, is the *hiccup*. It is known that the vagus nerve plays a role in it. Like for many other examples of self-organization, it takes quite some energy to get rid of a hiccup. Modern experimental research has shown that orgasms, which strongly stimulate the vagus nerve, are excellent ways to overcome hiccups. One of these researchers has won the 2006 IgNobel Prize for medicine for his work.

* *

Another important example of self-organization is the *weather*. If you want to know more about the known connections between the weather and the quality of human life on Earth, free of any ideology, read the wonderful book by Reichholf. It explains how the weather between the continents is connected and describes how and why the weather changed in the last one thousand years.

* *

Does self-organization or biological evolution contradict the second principle of thermodynamics? Of course not. Self-organization can even be shown to follow from the second principle, as any textbook on the topic will explain. Also for evolution there is no contradiction, as the Earth is not a closed thermodynamic system. Statements of the opposite are only made by crooks.

* *

Are systems that show self-organization the most complex ones that can be studied with evolution equations? No. The most complex systems are those that consist of many interacting self-organizing systems. The obvious example are *swarms*. Swarms of birds, as shown in Figure 242, of fish, of insects and of people – for example in a stadium or in cars on a highway – have been studied extensively and are still a subject of research. Their beauty is fascinating.

The other example of many interconnected self-organized systems is the brain; the exploration of how the interconnected neurons work will occupy researchers for many years. We will explore some aspects in the next volumes.

SUMMARY ON SELF-ORGANIZATION AND CHAOS

Appearance of *order*, in form of patterns, shapes and cycles, is not due to a decrease in entropy, but to a competition between driving causes and dissipative effects in open systems. Such appearance of order is predictable with (quite) simple equations. Also biological evolution is the appearance of order. It occurs automatically and obeys simple equations.

Chaos, the sensitivity to initial conditions, is common in strongly driven open systems, is at the basis of everyday chance, and often is described by simple equations. In nature, complexity is apparent. Motion is simple.




FIGURE 242 A typical swarm of starlings that visitors in Rome can observe every fall (© Andrea Cavagna/Physics Today).



FROM THE LIMITATIONS OF PHYSICS TO THE LIMITS OF MOTION

I only know that I know nothing.
Socrates, as cited by Plato

E have explored, so far, the everyday concept of motion. e called this exploration of moving objects and fluids *Galilean physics*. e found that in all these cases, motion is *predictable*: nature shows no surprises and no miracles. In particular, we have found six important aspects of this predictability:

- 1. Everyday motion is *continuous*. Motion allows us to define space and time.
- 2. Everyday motion *conserves* mass, momentum, energy and angular momentum. Nothing appears out of nothing.
- 3. Everyday motion is *relative*: motion depends on the observer.
- 4. Everyday motion is *reversible*: everyday motion can occur backwards.
- 5. Everyday motion is *mirror-invariant*: everyday motion can occur in a mirror-reversed way.

The last property is the most important, and in addition contains the previous ones:

6. Everyday motion is *lazy*: motion happens in a way that minimizes change, i.e., physical action.

This Galilean description of nature made engineering possible: textile machines, steam engines, combustion motors, kitchen appliances, many children toys, fitness machines and all the progress in the quality of life that came with these devices are due to the results of Galilean physics. But despite these successes, Socrates' saying, cited above, still applies to Galilean physics. Let us see why.

RESEARCH TOPICS IN CLASSICAL DYNAMICS

Even though mechanics and thermodynamics are now several hundred years old, research into its details is still ongoing. For example, we have already mentioned above that it is unclear whether the solar system is stable. The long-term future of the planets is unknown! In general, the behaviour of few-body systems interacting through gravitation is still a research topic of mathematical physics. Answering the simple question of how long a given set of bodies gravitating around each other will stay together is a formidable challenge. The history of this so-called *many-body problem* is long and involved. Interest-

Ref. 316

TABLE 50 Examples of errors in state-of-the art measurements (numbers in brackets give one standard deviation in the last digits), partly taken from physics.nist.gov/constants.

OBSERVATION	MEASUREMENT	PRECISION / ACCURACY
Highest precision achieved: ratio between the electron magnetic moment and the Bohr magneton $\mu_e/\mu_{\rm B}$	-1.001 159 652 180 76(24)	$2.6 \cdot 10^{-13}$
High precision: Rydberg constant	$10973731.568539(55)\mathrm{m}^{-1}$	$5.0 \cdot 10^{-12}$
High precision: astronomical unit	149 597 870.691(30) km	$2.0 \cdot 10^{-10}$
Industrial precision: part dimension tolerance in an automobile engine	1 μm of 20 cm	$5\cdot 10^{-6}$
Low precision: gravitational constant <i>G</i>	$6.67428(67)\cdot 10^{-11}\mathrm{Nm}^2/\mathrm{kg}^2$	$1.0\cdot 10^{-4}$
Everyday precision: human circadian clock governing sleep	15 h to 75 h	2

ing progress has been achieved, but the final answer still eludes us.

Many challenges remain in the fields of self-organization, of nonlinear evolution equations, and of chaotic motion. In these fields, *turbulence* is the most famous example: a precise description of turbulence has not yet been achieved. This and the other challenges motivate numerous researchers in mathematics, physics, chemistry, biology, medicine and the other natural sciences.

But apart from these research topics, classical physics leaves unanswered several basic questions.

WHAT IS CONTACT?

Democritus declared that there is a unique sort of motion: that ensuing from collision.

Simplicius, Commentary on the Physics of Aristotle, 42, 10

Ref. 317

Page 92

Of the questions unanswered by classical physics, the details of *contact* and *collisions* are among the most pressing. Indeed, we defined mass in terms of velocity changes during collisions. But why do objects change their motion in such instances? Why are collisions between two balls made of chewing gum different from those between two stainless-steel balls? What happens during those moments of contact?

Contact is related to material properties, which in turn influence motion in a complex way. The complexity is such that the sciences of material properties developed independently from the rest of physics for a long time; for example, the techniques of metallurgy (often called the oldest science of all), of chemistry and of cooking were related to the properties of motion only in the twentieth century, after having been independently pursued for thousands of years. Since material properties determine the essence of contact, we *need* knowledge about matter and about materials to understand the notion of mass, and thus of motion. The parts of our mountain ascent that deal with quantum theory will reveal these connections.

PRECISION AND ACCURACY

Ref. 318 Challenge 640 e Precision has its own fascination. How many digits of π , the ratio between circumference and diameter of a circle, do you know by heart? What is the largest number of digits of π you have calculated yourself?

Challenge 641 s

Is it possible to draw or cut a rectangle for which the ratio of lengths is a real number, e.g. of the form 0.131520091514001315211420010914..., whose digits encode a book? (A simple method would code a space as 00, the letter 'a' as 01, 'b' as 02, 'c' as 03, etc. Even more interestingly, could the number be printed inside its own book?)

Why are so many measurement results, such as those of Table 50, of limited precision, even if the budget is almost unlimited? These are questions about precision.

When we started climbing Motion Mountain, we explained that gaining height means increasing the *precision* of our description of nature. To make even this statement itself more precise, we distinguish between two terms: *precision* is the degree of reproducibility; *accuracy* is the degree of correspondence to the actual situation. Both concepts apply to measurements,* to statements and to physical concepts.

Challenge 642 s

Statements with false accuracy and false precision abound. What should we think of a car company – Ford – who claim that the drag coefficient $c_{\rm w}$ of a certain model is 0.375? Or of the official claim that the world record in fuel consumption for cars is 2315.473 km/l? Or of the statement that 70.3% of all citizens share a certain opinion? One lesson we learn from investigations into measurement errors is that we should never provide more digits for a result than we can put our hand into fire for.

Page 383

In short, precision and accuracy are *limited*. At present, the record number of reliable digits ever measured for a physical quantity is 13. Why so few? Classical physics doesn't provide an answer at all. What is the maximum number of digits we can expect in measurements; what determines it; and how can we achieve it? These questions are still open at this point in our ascent; they will be covered in the parts on quantum theory.

In our walk we aim for highest possible precision and accuracy, while avoiding false accuracy. Therefore, concepts have mainly to be *precise*, and descriptions have to be *accurate*. Any inaccuracy is a proof of lack of understanding. To put it bluntly, in our adventure, 'inaccurate' means *wrong*. Increasing the accuracy and precision of our description of nature implies leaving behind us all the mistakes we have made so far. This quest raises several issues.

CAN ALL OF NATURE BE DESCRIBED IN A BOOK?

Could the perfect physics publication, one that describes *all* of nature, exist? If it does, it must also describe itself, its own production – including its readers and its author – and most important of all, its own contents. Is such a book possible? Using the concept of information, we can state that such a book should contain all information contained in the universe. Is this possible? Let us check the options.

If nature requires an *infinitely* long book to be fully described, such a publication obviously cannot exist. In this case, only approximate descriptions of nature are possible and a perfect physics book is impossible.

^{*} For measurements, both precision and accuracy are best described by their *standard deviation*, as explained on page 388.

If nature requires a *finite* amount of information for its description, there are two options. One is that the information of the universe is so large that it cannot be summarized in a book; then a perfect physics book is again impossible. The other option is that the universe does contain a finite amount of information and that it can be summarized in a few short statements. This would imply that the rest of the universe would not add to the information already contained in the perfect physics book. But in this case, it seems that the entropy of the book and the entropy of the universe must be similar. This is also impossible, or at least unlikely.

We note that the answer to this puzzle also implies the answer to another puzzle: whether a brain can contain a full description of nature. In other words, the real question is: can we understand nature? Is our hike to the top of motion mountain possible? We usually believe this.

But the arguments just given imply that we believe something which seems unlikely: we believe that the universe does not contain more information than what our brain could contain or even contains already. Do we have an error in our arguments? Yes, we do. The terms 'universe' and 'information' are not used correctly in the above reasoning, as you might want to verify. We will solve this puzzle later in our adventure. Until then, do make up your own mind.

Vol. III, page 102 Challenge 643 e

SOMETHING IS WRONG ABOUT OUR DESCRIPTION OF MOTION

Darum kann es in der Logik auch *nie* Überraschungen geben.* Ludwig Wittgenstein, *Tractatus*, 6.1251

We described nature in a rather simple way. *Objects* are permanent and massive entities localized in space-time. *States* are changing properties of objects, described by position in space and instant in time, by energy and momentum, and by their rotational equivalents. *Time* is the relation between events measured by a clock. *Clocks* are devices in undisturbed motion whose position can be observed. *Space* and position is the relation between objects measured by a metre stick. *Metre sticks* are devices whose shape is subdivided by some marks, fixed in an invariant and observable manner. *Motion* is change of position with time (times mass); it is determined, does not show surprises, is conserved (even in death), and is due to gravitation and other interactions.

Even though this description works rather well in practice, it contains a circular definition. Can you spot it? Each of the two central concepts of motion is defined with the help of the other. Physicists worked for about 200 years on classical mechanics without noticing or wanting to notice the situation. Even thinkers with an interest in discrediting science did not point it out. Can an exact science be based on a circular definition? Obviously yes, and physics has done quite well so far. Is the situation unavoidable in principle? Undoing this logical loop is one of the aims of the rest of our walk. We will achieve the solution in the last leg of our adventure. To achieve the solution, we need to increase

Whenever precision is increased, imagination is restricted. We will discover that many types of motion that seem possible are not. Motion is limited. Nature limits speed, size,

substantially the level of precision in our description of motion.

Challenge 644 s

Challenge 645 s

^{* &#}x27;Hence there can *never* be surprises in logic.'

acceleration, mass, force, power and many other quantities. Continue reading the other parts of this adventure only if you are prepared to exchange fantasy for precision. It will be no loss, because exploring the precise working of nature will turn out to be more fascinating than any fantasy.

WHY IS MEASUREMENT POSSIBLE?

In the description of gravity given so far, the one that everybody learns – or should learn – at school, acceleration is connected to mass and distance via $a = GM/r^2$. That's all. But this simplicity is deceiving. In order to check whether this description is correct, we have to measure lengths and times. However, it is *impossible* to measure lengths and time intervals with any clock or any ruler based on the gravitational interaction alone! Try to conceive such an apparatus and you will be inevitably be disappointed. You always need a non-gravitational method to start and stop the stopwatch. Similarly, when you measure length, e.g. of a table, you have to hold a ruler or some other device near it. The interaction necessary to line up the ruler and the table cannot be gravitational.

A similar limitation applies even to mass measurements. Try to measure mass using gravitation alone. Any scale or balance needs other – usually mechanical, electromagnetic or optical – interactions to achieve its function. Can you confirm that the same applies to speed and to angle measurements? In summary, whatever method we use, *in order to measure velocity, length, time, and mass, interactions other than gravity are needed.* Our ability to measure shows that gravity is not all there is.

In short, Galilean physics does not explain our ability to measure. In fact, it does not even explain the existence of measurement standards. Why do objects have fixed lengths? Why do clocks work with regularity? Galilean physics cannot explain these observations; we will need relativity and quantum physics to find out.

IS MOTION UNLIMITED?

Galilean physics suggests that motion could go on forever. In fact, Galilean physics makes no clear statements on the universe as a whole. It tacitly suggests that it is infinite. Indeed, finitude does not fit with the Galilean description of motion. On the other hand, we know that the universe is *not* infinite: if it were infinite, the night would not be dark. Galilean physics is thus limited in its explanations because it disregards this and other limits to motion.

In particular, Galilean physics suggests that speeds can have any value. But the existence of infinite speeds in nature would not allow us to define time sequences. Clocks would be impossible. In other words, a description of nature that allows unlimited speeds is not precise. Precision requires limits. To achieve the highest possible precision, we need to discover all limits to motion. So far, we have discovered only one: there is a smallest entropy. We now turn to another, more striking one: the limit for speed of energy, objects and signals. To understand this limit, in the next volume we will explore the most rapid motion of energy, objects and signals that we know: the motion of light.

Challenge 646 s

Challenge 647 s

Challenge 648 s

Challenge 649 e

APPENDIX A

NOTATION AND CONVENTIONS

Page 383

Page 383 Vol. V, page 264 EWLY introduced concepts are indicated, throughout this text, by *italic typeface*. ew definitions are also referred to in the index. In this text, aturally we use the international SI units; they are defined in Appendix B. Experimental results are cited with limited precision, usually only two digits, as this is almost always sufficient for our purposes. High-precision reference values for important quantities can also be found in Appendix B. Additional precision values on composite physical systems are given in volume V.

But the information that is provided in this volume uses some additional conventions that are worth a second look.

THE LATIN ALPHABET

Ref. 319

What is written without effort is in general read without pleasure.

Samuel Johnson

Books are collections of symbols. *Writing* was probably invented between 3400 and 3300 BCE by the Sumerians in Mesopotamia (though other possibilities are also discussed). It then took over a thousand years before people started using symbols to represent sounds instead of concepts: this is the way in which the first alphabet was created. This happened between 2000 and 1600 BCE (possibly in Egypt) and led to the Semitic alphabet. The use of an alphabet had so many advantages that it was quickly adopted in all neighbouring cultures, though in different forms. As a result, the Semitic alphabet is the forefather of all alphabets used in the world.

This text is written using the Latin alphabet. At first sight, this seems to imply that its pronunciation *cannot* be explained in print, in contrast to the pronunciation of other alphabets or of the International Phonetic Alphabet (IPA). (They can be explained using the alphabet of the main text.) However, it *is* in principle possible to write a text that describes exactly how to move lips, mouth and tongue for each letter, using physical concepts where necessary. The descriptions of pronunciations found in dictionaries make indirect use of this method: they refer to the memory of pronounced words or sounds found in nature.

Historically, the Latin alphabet was derived from the Etruscan, which itself was a derivation of the Greek alphabet. There are two main forms.

Motion Mountain – The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christoph Schiller November 1997–Ma

The *ancient* Latin alphabet, used from the sixth century BCE onwards:

A B C D E F Z H I K L M N O P Q R S T V X

The classical Latin alphabet,

used from the second century BCE until the eleventh century:

A B C D E F G H I K L M N O P Q R S T V X Y Z

The letter G was added in the third century BCE by the first Roman to run a fee-paying school, Spurius Carvilius Ruga. He added a horizontal bar to the letter C and substituted the letter Z, which was not used in Latin any more, for this new letter. In the second century BCE, after the conquest of Greece, the Romans included the letters Y and Z from the Greek alphabet at the end of their own (therefore effectively reintroducing the Z) in order to be able to write Greek words. This classical Latin alphabet was stable for the next thousand years.*

The classical Latin alphabet was spread around Europe, Africa and Asia by the Romans during their conquests; due to its simplicity it began to be used for writing in numerous other languages. Most modern 'Latin' alphabets include a few other letters. The letter W was introduced in the eleventh century in French and was then adopted in most European languages.** The letter U was introduced in the mid fifteenth century in Italy, the letter J at the end of that century in Spain, to distinguish certain sounds which had previously been represented by V and I. The distinction proved a success and was already common in most European languages in the sixteenth century. The contractions æ and æ date from the Middle Ages. The German alphabet includes the *sharp s*, written ß, a contraction of 'ss' or 'sz', and the Nordic alphabets added *thorn*, written Þ or þ, and *eth*, written Đ or ð, both taken from the futhorc,*** and other signs.

Lower-case letters were not used in classical Latin; they date only from the Middle Ages, from the time of Charlemagne. Like most accents, such as \hat{e} , ς or \ddot{a} , which were also first used in the Middle Ages, lower-case letters were introduced to save the then expensive paper surface by shortening written words.

Outside a dog, a book is a man's best friend. Inside a dog, it's too dark to read.

Groucho Marx

Ref. 320

Ref. 321

Ref. 322

^{*} To meet Latin speakers and writers, go to www.alcuinus.net.

^{**} In Turkey, still in 2008, you can be convoked in front of a judge if you use the letters w, q or x in an official letter; these letters only exist in the Kurdish language, not in Turkish. Using them is 'unturkish' behaviour and punishable by law. It is not generally known how physics teachers cope with this situation.

^{***} The Runic script, also called *Futhark* or *Futhorc*, a type of alphabet used in the Middle Ages in Germanic, Anglo–Saxon and Nordic countries, probably also derives from the Etruscan alphabet. The name derives from the first six letters: f, u, th, a (or o), r, k (or c). The third letter is the letter thorn mentioned above; it is often written 'Y' in Old English, as in 'Ye Olde Shoppe.' From the runic alphabet Old English also took the letter *wyn* to represent the 'w' sound, and the already mentioned eth. (The other letters used in Old English – not from futhorc – were the *yogh*, an ancient variant of g, and the ligatures æ or Æ, called *ash*, and œ or Œ, called *ethel*.)

TABLE 51 The ancient and classical Greek alphabets, and the correspondence with Latin and Indian digits.

								_			
Anc.	Сг	ASS.	Name	Corr	ESP.	Anc.	Сг	A	ss.	ss. Name	ss. Name Cori
A	A	α	alpha	a	1	N	N	ν		nu	nu n
В	В	β	beta	b	2	Ξ	Ξ	ξ		xi	xi x
Γ	Γ	γ	gamma	g, n ¹	3	O	Ο	o		omicron	omicron o
Δ	Δ	δ	delta	d	4	П	П	π		pi	pi p
E	E	ε	epsilon	e	5	9 p, 4				qoppa ³	qoppa ³ q
F , ς			digamma,	w	6	P	P	ρ		rho	
			stigma ²			Σ	Σ	σ, ς		$sigma^4$	sigma ⁴ s
Z	Z	ζ	zeta	z	7	T	T	τ		tau	tau t
Н	Н	η	eta	e	8		Υ	υ		upsilon	upsilon y, u ⁵
Θ	Θ	θ	theta	th	9		Φ	φ		phi	phi ph, f
I	I	ι	iota	i, j	10		X	χ		chi	chi ch
K	K	κ	kappa	k	20		Ψ	Ψ		psi	psi ps
Λ	Λ	λ	lambda	1	30		Ω	ω		omega	omega o
M	M	μ	mu	m	40	$\Lambda \geqslant$				sampi ⁶	sampi ⁶ s

The regional archaic letters yot, sha and san are not included in the table. The letter san was the ancestor of sampi.

- 1. Only if before velars, i.e., before kappa, gamma, xi and chi.
- 2. 'Digamma' is the name used for the F-shaped form. It was mainly used as a letter (but also sometimes, in its lower-case form, as a number), whereas the shape and name 'stigma' is used only for the number. Both names were derived from the respective shapes; in fact, the stigma is a medieval, uncial version of the digamma. The name 'stigma' is derived from the fact that the letter looks like a sigma with a tau attached under it though unfortunately not in all modern fonts. The original letter name, also giving its pronunciation, was 'waw'.
- 3. The version of qoppa that looks like a reversed and rotated z is still in occasional use in modern Greek. Unicode calls this version 'koppa'.
- 4. The second variant of sigma is used only at the end of words.
- 5. Uspilon corresponds to 'u' only as the second letter in diphthongs.
- 6. In older times, the letter sampi was positioned between pi and qoppa.

THE GREEK ALPHABET

The Latin alphabet was derived from the Etruscan; the Etruscan from the Greek. The Greek alphabet was itself derived from the Phoenician or a similar northern Semitic alphabet in the tenth century BCE. The Greek alphabet, for the first time, included letters also for vowels, which the Semitic alphabets lacked (and often still lack). In the Phoenician alphabet and in many of its derivatives, such as the Greek alphabet, each letter has a proper name. This is in contrast to the Etruscan and Latin alphabets. The first two Greek letter names are, of course, the origin of the term *alphabet* itself.

In the tenth century BCE, the Ionian or *ancient* (eastern) Greek alphabet consisted of the upper-case letters only. In the sixth century BCE several letters were dropped, while a few new ones and the lower-case versions were added, giving the *classical* Greek alpha-

Ref. 323

bet. Still later, accents, subscripts and breathings were introduced. Table 51 also gives the values signified by the letters took when they were used as numbers. For this special use, the obsolete ancient letters were kept during the classical period; thus they also acquired lower-case forms.

The Latin correspondence in the table is the standard classical one, used for writing Greek words. The question of the correct *pronunciation* of Greek has been hotly debated in specialist circles; the traditional *Erasmian* pronunciation does not correspond either to the results of linguistic research, or to modern Greek. In classical Greek, the sound that sheep make was $\beta\eta-\beta\eta$. (Erasmian pronunciation wrongly insists on a narrow η ; modern Greek pronunciation is different for β , which is now pronounced 'v', and for η , which is now pronounced as 'i:' – a long 'i'.) Obviously, the pronunciation of Greek varied from region to region and over time. For Attic Greek, the main dialect spoken in the classical period, the question is now settled. Linguistic research has shown that chi, phi and theta were less aspirated than usually pronounced in English and sounded more like the initial sounds of 'cat', 'perfect' and 'tin'; moreover, the zeta seems to have been pronounced more like 'zd' as in 'buzzed'. As for the vowels, contrary to tradition, epsilon is closed and short whereas eta is open and long; omicron is closed and short whereas omega is wide and long, and upsilon is really a sound like a French 'u' or German 'ü.'

The Greek vowels can have rough or smooth *breathings*, *subscripts*, and acute, grave, circumflex or diaeresis *accents*. Breathings – used also on ρ – determine whether the letter is aspirated. Accents, which were interpreted as stresses in the Erasmian pronunciation, actually represented pitches. Classical Greek could have up to three of these added signs per letter; modern Greek never has more than one.

Another descendant of the Greek alphabet* is the *Cyrillic alphabet*, which is used with slight variations, in many Slavic languages, such as Russian and Bulgarian. However, there is no standard transcription from Cyrillic to Latin, so that often the same Russian name is spelled differently in different countries or even in the same country on different occasions.

TABLE 52 The beginning of the Hebrew abjad.

LETTER	Name	Corr	ESPONDENCE
×	aleph	a	1
コ	beth	b	2
ጏ	gimel	g	3
٦	daleth	d	4
etc.			

^{*} The Greek alphabet is also the origin of the *Gothic alphabet*, which was defined in the fourth century by Wulfila for the Gothic language, using also a few signs from the Latin and futhorc scripts.

The Gothic alphabet is not to be confused with the so-called *Gothic letters*, a style of the *Latin* alphabet used all over Europe from the eleventh century onwards. In Latin countries, Gothic letters were replaced in the sixteenth century by the *Antiqua*, the ancestor of the type in which this text is set. In other countries, Gothic letters remained in use for much longer. They were used in type and handwriting in Germany until 1941, when the National Socialist government suddenly abolished them, in order to comply with popular demand. They remain in sporadic use across Europe. In many physics and mathematics books, Gothic letters are used to denote vector quantities.

THE HEBREW ALPHABET AND OTHER SCRIPTS

Ref. 324 Vol. III, page 226 The Phoenician alphabet is also the origin of the Hebrew consonant alphabet or *abjad*. Its first letters are given in Table 52. Only the letter aleph is commonly used in mathematics, though others have been proposed.

Around one hundred writing systems are in use throughout the world. Experts classify them into five groups. *Phonemic alphabets*, such as Latin or Greek, have a sign for each consonant and vowel. *Abjads* or consonant alphabets, such as Hebrew or Arabic, have a sign for each consonant (sometimes including some vowels, such as aleph), and do not write (most) vowels; most abjads are written from right to left. *Abugidas*, also called *syllabic alphabets* or *alphasyllabaries*, such as Balinese, Burmese, Devanagari, Tagalog, Thai, Tibetan or Lao, write consonants and vowels; each consonant has an inherent vowel which can be changed into the others by diacritics. *Syllabaries*, such as Hiragana or Ethiopic, have a sign for each syllable of the language. Finally, *complex scripts*, such as Chinese, Mayan or the Egyptian hieroglyphs, use signs which have both sound and meaning. Writing systems can have text flowing from right to left, from bottom to top, and can count book pages in the opposite sense to this book.

Ref. 325

Even though there are about 7000 languages on Earth, there are only about one hundred writing systems used today. About fifty other writing systems have fallen out of use.* For physical and mathematical formulae, though, the sign system used in this text, based on Latin and Greek letters, written from left to right and from top to bottom, is a standard the world over. It is used independently of the writing system of the text containing it.

NUMBERS AND THE INDIAN DIGITS

Ref. 326

Both the digits and the method used in this text to write numbers originated in India. They were brought to the Mediterranean by Arabic mathematicians in the Middle Ages. The number system used in this text is thus much younger than the alphabet.** The Indian numbers were made popular in Europe by Leonardo of Pisa, called Fibonacci,*** in his book *Liber Abaci* or 'Book of Calculation', which he published in 1202. That book revolutionized mathematics. Anybody with paper and a pen (the pencil had not yet been invented) was now able to calculate and write down numbers as large as reason allowed, or even larger, and to perform calculations with them. Fibonacci's book started:

Novem figure indorum he sunt 9 8 7 6 5 4 3 2 1. Cum his itaque novem figuris, et cum hoc signo 0, quod arabice zephirum appellatur, scribitur quilibet numerus, ut inferius demonstratur.****

^{*} A well-designed website on the topic is www.omniglot.com. The main present and past writing systems are encoded in the Unicode standard, which at present contains 52 writing systems. See www.unicode.org. ** It is not correct to call the digits 0 to 9 *Arabic*. Both the digits used in Arabic texts and the digits used in Latin texts such as this one derive from the Indian digits. Only the digits 0, 2, 3 and 7 resemble those used in Arabic writing, and then only if they are turned clockwise by 90°.

^{***} Leonardo di Pisa, called Fibonacci (b. c. 1175 Pisa, d. 1250 Pisa), Italian mathematician, and the most important mathematician of his time.

^{**** &#}x27;The nine figures of the Indians are: 9 8 7 6 5 4 3 2 1. With these nine figures, and with this sign 0 which in Arabic is called zephirum, any number can be written, as will be demonstrated below.'

The Indian method of writing numbers, the *Indian number system*, introduced two innovations: a large one, the *positional system*, and a small one, the digit zero. The positional system, as described by Fibonacci, was so much more efficient that it completely replaced the previous *Roman number system*, which writes 1996 as IVMM or MCMIVC or MCMXCVI, as well as the *Greek number system*, in which the Greek letters were used for numbers in the way shown in Table 51, thus writing 1996 as $,\alpha \geqslant 0$, Compared to these systems, the Indian numbers are a much better technology.

The Indian number system proved so practical that calculations done on paper completely eliminated the need for the *abacus*, which therefore fell into disuse. The abacus is still in use only in those countries which do not use a positional system to write numbers.

The Indian number system also eliminated the need for systems to represent numbers with fingers. Such ancient systems, which could show numbers up to 10 000 and more, have left only one trace: the term 'digit' itself, which derives from the Latin word for finger.

The power of the positional number system is often forgotten. But only a positional number system allows mental calculations and makes calculating prodigies possible.*

THE SYMBOLS USED IN THE TEXT

To avoide the tediouse repetition of these woordes: is equalle to: I will sette as I doe often in woorke use, a paire of paralleles, or Gemowe lines of one lengthe, thus: = , bicause noe .2. thynges, can be moare equalle.

Robert Recorde*

Besides text and numbers, physics books contain other symbols. Most symbols have been developed over hundreds of years, so that only the clearest and simplest are now in use. In this mountain ascent, the symbols used as abbreviations for *physical* quantities are all taken from the Latin or Greek alphabets and are always defined in the context where they are used. The symbols designating units, constants and particles are defined in Appendix B and in Appendix B of volume V. There is an international standard for them (ISO 80000, formerly ISO 31), but it is shamefully expensive and thus virtually inaccessible; the symbols used in this text therefore are those in common use.

The *mathematical* symbols used in this text, in particular those for operations and relations, are given in the following list, together with their origins. The details of their history have been extensively studied in the literature.

Ref. 327

Page 383

Ref. 324

Ref. 328

Ref. 328

^{*} Currently, the shortest time for finding the thirteenth (integer) root of a hundred-digit (integer) number, a result with 8 digits, is 11.8 seconds. For more about the stories and the methods of calculating prodigies, see the bibliography.

^{**} Robert Recorde (*c.* 1510–1558), English mathematician and physician; he died in prison because of debts. The quotation is from his *The Whetstone of Witte*, 1557. An image showing the quote can be found at en. wikipedia.org/wiki/Equals_sign. It is usually suggested that the quote is the first introduction of the equal sign; claims that Italian mathematicians used the equal sign before Recorde are not backed up by convincing examples.

Motion Mountain
- The Adventure of Phy
sics pdf file ava
ilable free of char
ge at www.motionmountain.n
et Copyright
© Christoph Schiller
November 1997-
997–March 2012

SYMBOL	MEANING	ORIGIN
+,-	plus, minus	Johannes Widmann 1489; the plus sign is derived from Latin 'et'.
$\sqrt{}$	read as 'square root'	used by Christoff Rudolff in 1525; the sign evolved from a point.
=	equal to	Robert Recorde 1557
{},[],()	grouping symbols	use starts in the sixteenth century
>,<	larger than, smaller than	Thomas Harriot 1631
×	multiplied with, times	England <i>c.</i> 1600, made popular by William Oughtred 1631
a^n	<i>a</i> to the power n , $a \cdot \cdot a$ (n factors)	René Descartes 1637
x, y, z	coordinates, unknowns	René Descartes 1637
ax + by + c = 0	constants and equations for unknowns	René Descartes 1637
∞	infinity	John Wallis 1655
d/dx , dx , $\int y dx$	derivative, differential, integral	Gottfried Wilhelm Leibniz 1675
:	divided by	Gottfried Wilhelm Leibniz 1684
	multiplied with, times	Gottfried Wilhelm Leibniz c. 1690
a_1, a_n	indices	Gottfried Wilhelm Leibniz c. 1690
	similar to	Gottfried Wilhelm Leibniz c. 1690
π	circle number, 4 arctan 1	William Jones 1706
φx	function of <i>x</i>	Johann Bernoulli 1718
fx, f(x)	function of <i>x</i>	Leonhard Euler 1734
e	$\sum_{n=0}^{\infty} \frac{1}{n!} = \lim_{n \to \infty} (1 + 1/n)^n$	Leonhard Euler 1736
f'(x)	derivative of function at x	Giseppe Lagrangia 1770
$\Delta x, \sum$	difference, sum	Leonhard Euler 1755
i	imaginary unit, $+\sqrt{-1}$	Leonhard Euler 1777
#	is different from	Leonhard Euler eighteenth century
$\partial/\partial x$	partial derivative, read like ' d/dx '	it was derived from a cursive form of 'd'
		or of the letter 'dey' of the Cyrillic alphabet by Adrien-Marie Legendre in 1786 and made popular by Carl Gustav Jacobi in 1841
n!	faculty, 1 · 2 ··	Christian Kramp 1808
Δ	Laplace operator	Robert Murphy 1833
x	absolute value	Karl Weierstrass 1841
∇	read as 'nabla' (or 'del')	introduced by William Hamilton in 1853 and Peter Tait in 1867, named after the shape of an old Egyptian musical in- strument
C , ⊃	set inclusion	Ernst Schröder in 1890
∪, ∩	set union and intersection	Giuseppe Peano 1888
€	element of	Giuseppe Peano 1888
\otimes	dyadic product or tensor product or outer product	unknown
$\langle \psi , \psi \rangle$	bra and ket state vectors	Paul Dirac 1930
Ø	empty set	André Weil as member of the Nicolas Bourbaki group in the early twentieth century
[<i>x</i>]	the measurement unit of a quantity x	twentieth century
		·

Other signs used here have more complicated origins. The & sign is a contraction of Latin et meaning 'and', as is often more clearly visible in its variations, such as &, the common italic form.

Each of the punctuation signs used in sentences with modern Latin alphabets, such as , . ; :!? ' ' » « – () ... has its own history. Many are from ancient Greece, but the question mark is from the court of Charlemagne, and exclamation marks appear first in the sixteenth century.* The @ or at-sign probably stems from a medieval abbreviation of Latin ad, meaning 'at', similarly to how the & sign evolved from Latin et. In recent years, the smiley:-) and its variations have become popular. The smiley is in fact a new version of the 'point of irony' which had been formerly proposed, without success, by A. de Brahm (1868–1942).

The section sign § dates from the thirteenth century in northern Italy, as was shown by the German palaeographer Paul Lehmann. It was derived from ornamental versions of the capital letter C for *capitulum*, i.e., 'little head' or 'chapter.' The sign appeared first in legal texts, where it is still used today, and then spread into other domains.

The paragraph sign \P was derived from a simpler ancient form looking like the Greek letter Γ , a sign which was used in manuscripts from ancient Greece until well into the Middle Ages to mark the start of a new text paragraph. In the Middle Ages it took the modern form, probably because a letter c for *caput* was added in front of it.

One of the most important signs of all, the *white space* separating words, was due to Celtic and Germanic influences when these people started using the Latin alphabet. It became commonplace between the ninth and the thirteenth century, depending on the language in question.

CALENDARS

Ref 330

Ref. 331

Ref. 333

The many ways to keep track of time differ greatly from civilization to civilization. The most common calendar, and the one used in this text, is also one of the most absurd, as it is a compromise between various political forces who tried to shape it.

In ancient times, independent localized entities, such as tribes or cities, preferred *lu-nar* calendars, because lunar timekeeping is easily organized locally. This led to the use of the month as a calendar unit. Centralized states imposed *solar* calendars, based on the year. Solar calendars require astronomers, and thus a central authority to finance them. For various reasons, farmers, politicians, tax collectors, astronomers, and some, but not all, religious groups wanted the calendar to follow the solar year as precisely as possible. The compromises necessary between days and years are the origin of leap days. The compromises necessary between months and year led to the varying lengths of the months; they are different in different calendars. The most commonly used year–month structure was organized over 2000 years ago by Gaius Julius Caesar, and is thus called the *Julian calendar*.

The system was destroyed only a few years later: August was lengthened to 31 days when it was named after Augustus. Originally, the month was only 30 days long; but in order to show that Augustus was as important as Caesar, after whom July is named, all

^{*} On the parenthesis see the beautiful book by J. Lennard, *But I Digress*, Oxford University Press, 1991.

month lengths in the second half of the year were changed, and February was shortened by one additional day.

Ref. 334

Page 191

The week is an invention of Babylonia. One day in the Babylonian week was 'evil' or 'unlucky', so it was better to do nothing on that day. The modern week cycle with its resting day descends from that superstition. (The way astrological superstition and astronomy cooperated to determine the order of the weekdays is explained in the section on gravitation.) Although about three thousand years old, the week was fully included into the Julian calendar only around the year 300, towards the end of the Western Roman Empire. The final change in the Julian calendar took place between 1582 and 1917 (depending on the country), when more precise measurements of the solar year were used to set a new method to determine leap days, a method still in use today. Together with a reset of the date and the fixation of the week rhythm, this standard is called the *Gregorian calendar* or simply the *modern calendar*. It is used by a majority of the world's population.

Despite its complexity, the modern calendar does allow you to determine the day of the week of a given date in your head. Just execute the following six steps:

- 1. take the last two digits of the year, and divide by 4, discarding any fraction;
- 2. add the last two digits of the year;
- 3. subtract 1 for January or February of a leap year;
- 4. add 6 for 2000s or 1600s, 4 for 1700s or 2100s,
 - 2 for 1800s and 2200s, and 0 for 1900s or 1500s;
- 5. add the day of the month;
- 6. add the month key value, namely 144 025 036 146 for JFM AMJ JAS OND.

The remainder after division by 7 gives the day of the week, with the correspondence 1-2-3-4-5-6-0 meaning Sunday-Monday-Tuesday-Wednesday-Thursday-Friday-Saturday.*

When to start counting the years is a matter of choice. The oldest method not attached to political power structures was that used in ancient Greece, when years were counted from the first Olympic games. People used to say, for example, that they were born in the first year of the twenty-third Olympiad. Later, political powers always imposed the counting of years from some important event onwards.** Maybe reintroducing the Olympic counting is worth considering?

Some other standards set by the Roman Empire explain several abbreviations used in the text:

- c. is a Latin abbreviation for circa and means 'roughly';
- i.e. is a Latin abbreviation for id est and means 'that is';
- e.g. is a Latin abbreviation for exempli gratia and means 'for the sake of example';
- ibid. is a Latin abbreviation for *ibidem* and means 'at that same place';
- inf. is a Latin abbreviation for infra and means '(see) below';
- op. cit. is a Latin abbreviation for opus citatum and means 'the cited work';
- et al. is a Latin abbreviation for et alii and means 'and others'.

By the way, *idem* means 'the same' and *passim* means 'here and there' or 'throughout'. Many terms used in physics, like frequency, acceleration, velocity, mass, force, momentum, inertia, gravitation and temperature,

^{*} Remembering the intermediate result for the current year can simplify things even more, especially since the dates 4.4, 6.6, 8.8, 10.10, 12.12, 9.5, 5.9, 7.11, 11.7 and the last day of February all fall on the same day of the week, namely on the year's intermediate result plus 4.

^{**} The present counting of years was defined in the Middle Ages by setting the date for the foundation of Rome to the year 753 BCE, or 753 *before the Common Era*, and then counting backwards, so that the BCE years behave almost like negative numbers. However, the year 1 follows directly after the year 1 BCE: there was no year 0.

PEOPLE NAMES

In the Far East, such as Corea*, Japan or China, family names are put in front of the given name. For example, the first Japanese winner of the Nobel Prize in Physics was Yukawa Hideki. In *India*, often, but not always, there is no family name; in those cases, the father's first name is used. In Russia, the family name is rarely used in conversation; instead, the first name of the father is. For example, Lev Landau was addressed as Lev Davidovich ('son of David'). In addition, Russian transliteration is not standardized; it varies from country to country and from tradition to tradition. For example, one finds the spellings Dostojewski, Dostoevskij, Dostoëvski and Dostoyevsky for the same person. In the Netherlands, the official given names are never used; every person has a semiofficial first name by which he is called. For example, Gerard 't Hooft's official given name is Gerardus. In Germany, some family names have special pronunciations. For example, Voigt is pronounced 'Fohgt'. In Italy, during the Middle Age and the Renaissance, people were called by their first name only, such as Michelangelo or Galileo, or often by first name plus a personal surname that was not their family name, but was used like one, such as Niccolò Tartaglia or Leonardo Fibonacci. In ancient Rome, the name by which people are known is usually their surname. The family name was the middle name. For example, Cicero's family name was Tullius. The law introduced by Cicero was therefore known as 'lex Tullia'. In ancient Greece, there were no family names. People had only one name. In the English language, the Latin version of the Greek name is used, such as Democritus.

ABBREVIATIONS AND EPONYMS OR CONCEPTS?

Sentences like the following are the scourge of modern physics:

The EPR paradox in the Bohm formulation can perhaps be resolved using the GRW approach, using the WKB approximation of the Schrödinger equation.

Using such vocabulary is the best way to make language unintelligible to outsiders. (In fact, the sentence is nonsense anyway, because the 'GRW approach' is false.) First of all, the sentence uses abbreviations, which is a shame. On top of this, the sentence uses people's names to characterize concepts, i.e., it uses *eponyms*. Originally, eponyms were intended as tributes to outstanding achievements. Today, when formulating radical new laws or variables has become nearly impossible, the spread of eponyms intelligible to a steadily decreasing number of people simply reflects an increasingly ineffective drive to fame.

Ref. 335

are derived from Latin. In fact, it is arguable that the language of science has been Latin for over two thousand years. In Roman times it was Latin vocabulary with Latin grammar, in modern times it switched to Latin vocabulary with French grammar, then for a short time to Latin vocabulary with German grammar, after which it changed to Latin vocabulary with British/American grammar.

Many units of measurement also date from Roman times, as explained in the next appendix. Even the infatuation with Greek technical terms, as shown in coinages such as 'gyroscope', 'entropy' or 'proton', dates from Roman times.

^{*} Corea was temporarily forced to change its spelling to 'Korea' by the Japanese Army because the generals could not bear the fact that Corea preceded Japan in the alphabet. This is not a joke.

Eponyms are a proof of scientist's lack of imagination. We avoid them as much as possible in our walk and give *common* names to mathematical equations or entities wherever possible. People's names are then used as appositions to these names. For example, 'Newton's equation of motion' is never called 'Newton's equation'; 'Einstein's field equations' is used instead of 'Einstein's equations'; and 'Heisenberg's equation of motion' is used instead of 'Heisenberg's equation'.

However, some exceptions are inevitable: certain terms used in modern physics have no real alternatives. The Boltzmann constant, the Planck scale, the Compton wavelength, the Casimir effect and Lie groups are examples. In compensation, the text makes sure that you can look up the definitions of these concepts using the index. In addition, the text tries to provide pleasurable reading.

Aountain – The Adventure of Physics - pdf file available free of charge at www.motionmountain.net - Copyright © Christoph Sc

UNITS, MEASUREMENTS AND CONSTANTS

any different systems of units have been used throughout the world. ost standards confer power to the organization in charge of them. Such power can be misused; this is the case today, for example in the computer industry, and was so in the distant past. The solution is the same in both cases: organize an independent and global standard. For measurement units, this happened in the eighteenth century: in order to avoid misuse by authoritarian institutions, to eliminate problems with differing, changing and irreproducible standards, and – this is not a joke – to simplify tax collection and to make it more just, a group of scientists, politicians and economists agreed on a set of units. It is called the *Système International d'Unités*, abbreviated *SI*, and is defined by an international treaty, the 'Convention du Mètre'. The units are maintained by an international organization, the 'Conférence Générale des Poids et Mesures', and its daughter organizations, the 'Commission Internationale des Poids et Mesures' and the 'Bureau International des Poids et Mesures' (BIPM). All originated in the times just before the French revolution.

Ref. 337

SI UNITS

All SI units are built from seven *base units*, whose official definitions, translated from French into English, are given below, together with the dates of their formulation:

- 'The *second* is the duration of 9 192 631 770 periods of the radiation corresponding to the transition between the two hyperfine levels of the ground state of the caesium 133 atom.' (1967)*
- 'The *metre* is the length of the path travelled by light in vacuum during a time interval of 1/299 792 458 of a second.' (1983)*
- 'The *kilogram* is the unit of mass; it is equal to the mass of the international prototype of the kilogram.' (1901)*
- 'The *ampere* is that constant current which, if maintained in two straight parallel conductors of infinite length, of negligible circular cross-section, and placed 1 metre apart in vacuum, would produce between these conductors a force equal to $2 \cdot 10^{-7}$ newton per metre of length.' (1948)*
- 'The *kelvin*, unit of thermodynamic temperature, is the fraction 1/273.16 of the thermodynamic temperature of the triple point of water.' (1967)*
- 'The *mole* is the amount of substance of a system which contains as many elementary entities as there are atoms in 0.012 kilogram of carbon 12.' (1971)*

• 'The *candela* is the luminous intensity, in a given direction, of a source that emits monochromatic radiation of frequency $540 \cdot 10^{12}$ hertz and has a radiant intensity in that direction of (1/683) watt per steradian.' (1979)*

Note that both time and length units are defined as certain properties of a standard example of motion, namely light. In other words, also the Conférence Générale des Poids et Mesures makes the point that the observation of motion is a *prerequisite* for the definition and construction of time and space. *Motion is the fundament each observation and measurements*. By the way, the use of light in the definitions had been proposed already in 1827 by Jacques Babinet.**

From these basic units, all other units are defined by multiplication and division. Thus, all SI units have the following properties:

- SI units form a system with *state-of-the-art precision*: all units are defined with a precision that is higher than the precision of commonly used measurements. Moreover, the precision of the definitions is regularly being improved. The present relative uncertainty of the definition of the second is around 10^{-14} , for the metre about 10^{-10} , for the kilogram about 10^{-9} , for the ampere 10^{-7} , for the mole less than 10^{-6} , for the kelvin 10^{-6} and for the candela 10^{-3} .
- SI units form an *absolute* system: all units are defined in such a way that they can be reproduced in every suitably equipped laboratory, independently, and with high precision. This avoids as much as possible any misuse by the standard-setting organization. (The kilogram, still defined with the help of an artefact, is the last exception to this requirement; extensive research is under way to eliminate this artefact from the definition an international race that will take a few more years. There are two approaches: counting particles, or fixing \hbar . The former can be achieved in crystals, the latter using any formula where \hbar appears, such as the formula for the de Broglie wavelength or that of the Josephson effect.)
- SI units form a *practical* system: the base units are quantities of everyday magnitude. Frequently used units have standard names and abbreviations. The complete list includes the seven base units just given, the supplementary units, the derived units and the admitted units.

The *supplementary* SI units are two: the unit for (plane) angle, defined as the ratio of arc length to radius, is the *radian* (rad). For solid angle, defined as the ratio of the subtended area to the square of the radius, the unit is the *steradian* (sr).

The *derived* units with special names, in their official English spelling, i.e., without capital letters and accents, are:

Page 92 Ref. 338

^{*} The respective symbols are s, m, kg, A, K, mol and cd. The international prototype of the kilogram is a platinum–iridium cylinder kept at the BIPM in Sèvres, in France. For more details on the levels of the caesium atom, consult a book on atomic physics. The Celsius scale of temperature θ is defined as: θ /°C = T/K - 273.15; note the small difference with the number appearing in the definition of the kelvin. SI also states: 'When the mole is used, the elementary entities must be specified and may be atoms, molecules, ions, electrons, other particles, or specified groups of such particles.' In the definition of the mole, it is understood that the carbon 12 atoms are unbound, at rest and in their ground state. In the definition of the candela, the frequency of the light corresponds to 555.5 nm, i.e., green colour, around the wavelength to which the eye is most sensitive.

^{**} Jacques Babinet (1794–1874), French physicist who published important work in optics.

NAME	ABBREVIATION
hertz	Hz = 1/s
pascal	$Pa = N/m^2 = kg/m s^2$
watt	$W = kg m^2/s^3$
volt	$V = kg m^2 / As^3$
ohm	$\Omega = V/A = kg m^2/A^2 s^3$
weber	$Wb = Vs = kg m^2 / As^2$
henry	$H = Vs/A = kg m^2/A^2 s^2$
lumen	lm = cd sr
becquerel	Bq = 1/s
sievert	$Sv = J/kg = m^2/s^2$

N а м е	ABBREVIATION
newton	$N = kg m/s^2$
joule	$J = Nm = kg m^2/s^2$
coulomb	C = As
farad	$F = As/V = A^2s^4/kg m^2$
siemens	$S = 1/\Omega$
tesla	$T = Wb/m^2 = kg/As^2 = kg/Cs$
degree Celsius	°C (see definition of kelvin)
lux	$lx = lm/m^2 = cd sr/m^2$
gray	$Gy = J/kg = m^2/s^2$
katal	kat = mol/s

We note that in all definitions of units, the kilogram only appears to the powers of 1, 0 and -1. Can you try to formulate the reason?

The admitted non-SI units are minute, hour, day (for time), degree $1^{\circ} = \pi/180 \, \text{rad}$, minute $1' = \pi/10 \, 800 \, \text{rad}$, second $1'' = \pi/648 \, 000 \, \text{rad}$ (for angles), litre and tonne. All other units are to be avoided.

All SI units are made more practical by the introduction of standard names and abbreviations for the powers of ten, the so-called *prefixes*:*

Pov	VER NAME	Powi	er Nai	м Е	Pow	er Nami	Ξ	Power	R NAME	
10 ¹	deca da	10^{-1}	deci	d	10 ¹⁸	Exa	E	10^{-18}	atto	a
10^2	hecto h	10^{-2}	centi	c	10^{21}	Zetta	Z	10^{-21}	zepto	z
10^3	kilo k	10^{-3}	milli	m	10^{24}	Yotta	Y	10^{-24}	yocto	y
10^{6}	Mega M	10^{-6}	micro	μ	unoffi	cial:		Ref. 339		
10^{9}	Giga G	10^{-9}	nano	n	10^{27}	Xenta	X	10^{-27}	xenno	X
10^{12}	Tera T	10^{-12}	pico	p	10^{30}	Wekta	W	10^{-30}	weko	w
10^{15}	Peta P	10^{-15}	femto	f	10^{33}	Vendekta	V	10^{-33}	vendeko	v
					10^{36}	Udekta	U	10^{-36}	udeko	u

• SI units form a *complete* system: they cover in a systematic way the full set of observables of physics. Moreover, they fix the units of measurement for all other sciences as well.

Translate: I was caught in such a traffic jam that I needed a microcentury for a picoparsec and that my car's fuel consumption was two tenths of a square millimetre.

Challenge 650 s

^{*} Some of these names are invented (yocto to sound similar to Latin *octo* 'eight', zepto to sound similar to Latin *septem*, yotta and zetta to resemble them, exa and peta to sound like the Greek words ἑξάκις and πεντάκις for 'six times' and 'five times', the unofficial ones to sound similar to the Greek words for nine, ten, eleven and twelve); some are from Danish/Norwegian (atto from *atten* 'eighteen', femto from *femten* 'fifteen'); some are from Latin (from *mille* 'thousand', from *centum* 'hundred', from *decem* 'ten', from *nanus* 'dwarf'); some are from Italian (from *piccolo* 'small'); some are Greek (micro is from μικρός 'small', deca/deka from δέκα 'ten', hecto from έκατόν 'hundred', kilo from χίλιοι 'thousand', mega from μέγας 'large', giga from γίγας 'giant', tera from τέρας 'monster').

- SI units form a *universal* system: they can be used in trade, in industry, in commerce, at home, in education and in research. They could even be used by extraterrestrial civilizations, if they existed.
- SI units form a *coherent* system: the product or quotient of two SI units is also an SI unit. This means that in principle, the same abbreviation, e.g. 'SI', could be used for every unit.

The SI units are not the only possible set that could fulfil all these requirements, but they are the only existing system that does so.*

THE MEANING OF MEASUREMENT

Challenge 652 e

Every measurement is a comparison with a standard. Therefore, any measurement requires *matter* to realize the standard (even for a speed standard), and *radiation* to achieve the comparison. The concept of measurement thus assumes that matter and radiation exist and can be clearly separated from each other.

Every measurement is a comparison. Measuring thus implies that space and time exist, and that they differ from each other.

Every measurement produces a measurement result. Therefore, every measurement implies the *storage* of the result. The process of measurement thus implies that the situation before and after the measurement can be distinguished. In other terms, every measurement is an *irreversible* process.

Every measurement is a process. Thus every measurement takes a certain amount of time and a certain amount of space.

All these properties of measurements are simple but important. Beware of anybody who denies them.

CURIOSITIES AND FUN CHALLENGES ABOUT UNITS

Not using SI units can be expensive. In 1999, NASA lost a satellite on Mars because some software programmers had used provincial units instead of SI units in part of the code. As a result of using feet instead of meters, the Mars Climate Orbiter crashed into the planet, instead of orbiting it; the loss was around 100 million euro.**

* *

The second does not correspond to 1/86 400th of the day any more, though it did in the year 1900; the Earth now takes about 86 400.002 s for a rotation, so that the *International Earth Rotation Service* must regularly introduce a leap second to ensure that the Sun is

^{*} Apart from international units, there are also *provincial* units. Most provincial units still in use are of Roman origin. The mile comes from *milia passum*, which used to be one thousand (double) strides of about 1480 mm each; today a nautical mile, once defined as minute of arc on the Earth's surface, is exactly 1852 m). The inch comes from *uncia/onzia* (a twelfth – now of a foot). The pound (from *pondere* 'to weigh') is used as a translation of *libra* – balance – which is the origin of its abbreviation lb. Even the habit of counting in dozens instead of tens is Roman in origin. These and all other similarly funny units – like the system in which all units start with 'f', and which uses furlong/fortnight as its unit of velocity – are now officially defined as multiples of SI units.

^{**} This story revived an old but false urban legend that states that only three countries in the world do not use SI units: Liberia, the USA and Myanmar.

Ref. 340

Ref. 341

Page 395

Challenge 653 s

at the highest point in the sky at 12 o'clock sharp.* The time so defined is called *Universal Time Coordinate*. The speed of rotation of the Earth also changes irregularly from day to day due to the weather; the average rotation speed even changes from winter to summer because of the changes in the polar ice caps; and in addition that average decreases over time, because of the friction produced by the tides. The rate of insertion of leap seconds is therefore higher than once every 500 days, and not constant in time.

* *

The most precise clock ever built, using microwaves, had a stability of 10^{-16} during a running time of 500 s. For longer time periods, the record in 1997 was about 10^{-15} ; but values around 10^{-17} seem within technological reach. The precision of clocks is limited for short measuring times by *noise*, and for long measuring times by *drifts*, i.e., by systematic effects. The region of highest stability depends on the clock type; it usually lies between 1 ms for optical clocks and 5000 s for masers. Pulsars are the only type of clock for which this region is not known yet; it certainly lies at more than 20 years, the time elapsed at the time of writing since their discovery.

* *

The least precisely measured of the fundamental constants of physics are the gravitational constant G and the strong coupling constant α_s . Even less precisely known are the age of the universe and its density (see Table 57).

* *

The precision of mass measurements of solids is limited by such simple effects as the adsorption of water. Can you estimate the mass of a monolayer of water – a layer with thickness of one molecule – on a metal weight of 1 kg?

* *

In the previous millennium, thermal energy used to be measured using the unit *calorie*, written as cal. 1 cal is the energy needed to heat 1 g of water by 1 K. To confuse matters, 1 kcal was often written 1 Cal. (One also spoke of a large and a small calorie.) The value of 1 kcal is 4.1868 kJ.

* *

SI units are adapted to humans: the values of heartbeat, human size, human weight, human temperature and human substance are no more than a couple of orders of magnitude near the unit value. SI units thus (roughly) confirm what Protagoras said 25 centuries ago: 'Man is the measure of all things.'

* *

Some units systems are particularly badly adapted to humans. The most infamous is shoe

^{*} Their website at hpiers.obspm.fr gives more information on the details of these insertions, as does maia. usno.navy.mil, one of the few useful military websites. See also www.bipm.fr, the site of the BIPM.

size *S*. It is a pure number calculated as

$$S_{\text{France}} = 1.5 \text{ cm}^{-1} (l + (1 \pm 1) \text{ cm})$$

$$S_{\text{central Europe}} = 1.5748 \text{ cm}^{-1} (l + (1 \pm 1) \text{ cm})$$

$$S_{\text{Anglosaxon men}} = 1.181 \text{ cm}^{-1} (l + (1 \pm 1) \text{ cm}) - 22$$
(118)

where l is the length of a foot and the correction length depends on the manufacturing company. In addition, the Anglosaxon formula is not valid for women and children, where the first factor depends, for marketing reasons, both on manufacturer and size itself. The ISO standard for shoe size requires, unsurprisingly, to use foot length in millimetres.

* *

The table of SI prefixes covers 72 orders of magnitude. How many additional prefixes will be needed? Even an extended list will include only a small part of the infinite range of possibilities. Will the Conférence Générale des Poids et Mesures have to go on forever, defining an infinite number of SI prefixes? Why?

Challenge 654 s

* *

The French philosopher Voltaire, after meeting Newton, publicized the now famous story that the connection between the fall of objects and the motion of the Moon was discovered by Newton when he saw an apple falling from a tree. More than a century later, just before the French Revolution, a committee of scientists decided to take as the unit of force precisely the force exerted by gravity on a *standard apple*, and to name it after the English scientist. After extensive study, it was found that the mass of the standard apple was 101.9716 g; its weight was called 1 newton. Since then, visitors to the museum in Sèvres near Paris have been able to admire the standard metre, the standard kilogram and the standard apple.*

PRECISION AND ACCURACY OF MEASUREMENTS

Measurements are the basis of physics. Every measurement has an *error*. Errors are due to lack of precision or to lack of accuracy. *Precision* means how well a result is reproduced when the measurement is repeated; *accuracy* is the degree to which a measurement corresponds to the actual value.

Lack of precision is due to accidental or *random errors*; they are best measured by the *standard deviation*, usually abbreviated σ ; it is defined through

$$\sigma^2 = \frac{1}{n-1} \sum_{i=1}^{n} (x_i - \bar{x})^2 , \qquad (119)$$

^{*} To be clear, this is a joke; no standard apple exists. It is *not* a joke however, that owners of several apple trees in Britain and in the US claim descent, by rerooting, from the original tree under which Newton had his insight. DNA tests have even been performed to decide if all these derive from the same tree. The result was, unsurprisingly, that the tree at MIT, in contrast to the British ones, is a fake.




FIGURE 243 A precision experiment and its measurement distribution. The precision is high if the width of the distribution is narrow; the accuracy is high if the centre of the distribution agrees with the actual value.

Challenge 655 s

where \bar{x} is the average of the measurements x_i . (Can you imagine why n-1 is used in the formula instead of n?)

For most experiments, the distribution of measurement values tends towards a normal distribution, also called *Gaussian distribution*, whenever the number of measurements is increased. The distribution, shown in Figure 243, is described by the expression

$$N(x) \approx e^{-\frac{(x-\bar{x})^2}{2\sigma^2}} \ . \tag{120}$$

Challenge 656 e

The square σ^2 of the standard deviation is also called the *variance*. For a Gaussian distribution of measurement values, 2.35σ is the full width at half maximum.

Lack of accuracy is due to *systematic errors*; usually these can only be estimated. This estimate is often added to the random errors to produce a *total experimental error*, sometimes also called *total uncertainty*. The *relative* error or uncertainty is the ratio between the error and the measured value.

For example, a professional measurement will give a result such as 0.312(6) m. The number between the parentheses is the standard deviation σ , in units of the last digits. As above, a Gaussian distribution for the measurement results is assumed. Therefore, a value of 0.312(6) m implies that the actual value is expected to lie

Challenge 657 e

- within 1σ with 68.3% probability, thus in this example within 0.312 \pm 0.006 m;
- within 2σ with 95.4% probability, thus in this example within 0.312 ± 0.012 m;
- within 3σ with 99.73% probability, thus in this example within 0.312 ± 0.018 m;
- within 4σ with 99.9937% probability, thus in this example within 0.312 ± 0.024 m;
- within 5σ with 99.999 943% probability, thus in this example within 0.312 ± 0.030 m;
- within 6σ with 99.999 999 80% probability, thus in this example within

- $0.312 \pm 0.036 \,\mathrm{m}$;
- within 7σ with 99.999 999 74% probability, thus in this example within 0.312 ± 0.041 m.

Challenge 658 s (Do the latter numbers make sense?)

Note that standard deviations have one digit; you must be a world expert to use two, and a fool to use more. If no standard deviation is given, a (1) is assumed. As a result, among professionals, 1 km and 1000 m are *not* the same length!

What happens to the errors, if two measured values A and B are added or subtracted? If the all measurements are independent – or uncorrelated – the standard deviation of the sum and that of difference is given by $\sigma = \sqrt{\sigma_A^2 + \sigma_B^2}$. For both the product or ratio of two measured and uncorrelated values C and D, the result is $\rho = \sqrt{\rho_C^2 + \rho_D^2}$, where the ρ terms are the P terms are t

Assume you measure that an object moves 1.0 m in 3.0 s: what is the measured speed value?

Challenge 659 s

LIMITS TO PRECISION

What are the limits to accuracy and precision? There is no way, even in principle, to measure a length x to a *precision* higher than about 61 digits, because in nature, the ratio between the largest and the smallest measurable length is $\Delta x/x > l_{\rm Pl}/d_{\rm horizon} = 10^{-61}$. (Is this ratio valid also for force or for volume?) In the final volume of our text, studies of clocks and metre bars strengthen this theoretical limit.

Challenge 660 e Vol. VI, page 87

But it is not difficult to deduce more stringent practical limits. No imaginable machine can measure quantities with a higher precision than measuring the diameter of the Earth within the smallest length ever measured, about 10^{-19} m; that is about 26 digits of precision. Using a more realistic limit of a 1000 m sized machine implies a limit of 22 digits. If, as predicted above, time measurements really achieve 17 digits of precision, then they are nearing the practical limit, because apart from size, there is an additional practical restriction: cost. Indeed, an additional digit in measurement precision often means an additional digit in equipment cost.

PHYSICAL CONSTANTS

In physics, general observations are deduced from more fundamental ones. As a consequence, many measurements can be deduced from more fundamental ones. The most fundamental measurements are those of the physical constants.

The following tables give the world's best values of the most important physical constants and particle properties – in SI units and in a few other common units – as published in the standard references. The values are the world averages of the best measurements made up to the present. As usual, experimental errors, including both random and estimated systematic errors, are expressed by giving the standard deviation in the last digits. In fact, behind each of the numbers in the following tables there is a long story which is worth telling, but for which there is not enough room here.

Ref. 344

Ref. 344

In principle, all quantitative properties of matter can be calculated with quantum theory and the values of certain physical constants. For example, colour, density and elastic properties can be predicted using the equations of the standard model of high-energy

physics and the values of the following basic constants. Vol. V, page 193

TABLE 54 Basic physical constants.

Quantity	SYMBOL	VALUE IN SI UNITS	Uncert.
Number of space-time dimen	sions	3 + 1	0 ^b
Vacuum speed of light ^c	С	299 792 458 m/s	0
Vacuum permeability ^c	μ_0	$4\pi\cdot 10^{-7}~H/m$	0
		= 1.256 637 061 435 μH/1	m0
Vacuum permittivity ^c	$\varepsilon_0 = 1/\mu_0 c^2$	8.854 187 817 620 pF/m	0
Original Planck constant	h	$6.62606876(52)\cdot10^{-34}\mathrm{Js}$	$7.8 \cdot 10^{-8}$
Reduced Planck constant	\hbar	$1.054571596(82)\cdot 10^{-34}\mathrm{Js}$	$7.8\cdot10^{-8}$
Positron charge	e	0.160 217 646 2(63) aC	$3.9 \cdot 10^{-8}$
Boltzmann constant	k	$1.3806503(24)\cdot 10^{-23}\mathrm{J/K}$	$1.7\cdot 10^{-6}$
Gravitational constant	G	$6.673(10) \cdot 10^{-11} \mathrm{Nm^2/kg^2}$	$1.5\cdot 10^{-3}$
Gravitational coupling consta	$nt\kappa = 8\pi G/c^4$	$2.076(3) \cdot 10^{-43} \text{s}^2/\text{kg m}$	$1.5\cdot 10^{-3}$
Fine structure constant, ^d	$\alpha = \frac{e^2}{4\pi\varepsilon_0\hbar c}$	1/137.035 999 76(50)	$3.7\cdot10^{-9}$
e.m. coupling constant	$=g_{\rm em}(m_{\rm e}^2c^2)$	= 0.007 297 352 533(27)	$3.7 \cdot 10^{-9}$
Fermi coupling constant, ^d	$G_{\rm F}/(\hbar c)^3$	$1.16639(1) \cdot 10^{-5} \mathrm{GeV}^{-2}$	$8.6 \cdot 10^{-6}$
weak coupling constant	$\alpha_{\rm w}(M_{\rm Z}) = g_{\rm w}^2/4\pi$	1/30.1(3)	$1 \cdot 10^{-2}$
Weak mixing angle	$\sin^2 \theta_{\rm W}(\overline{MS})$	0.231 24(24)	$1.0\cdot10^{-3}$
	$\sin^2 \theta_{\rm W}$ (on shell)	0.2224(19)	$8.7\cdot10^{-3}$
	$=1-(m_{\rm W}/m_{\rm Z})^2$		
Strong coupling constant ^d	$\alpha_{\rm s}(M_{\rm Z}) = g_{\rm s}^2/4\pi$	0.118(3)	$25\cdot 10^{-3}$
Electron mass	$m_{\rm e}$	$9.10938188(72)\cdot 10^{-31}\mathrm{kg}$	$7.9 \cdot 10^{-8}$
		$5.485799110(12) \cdot 10^{-4} \mathrm{u}$	$2.1\cdot 10^{-9}$
		0.510 998 902(21) MeV	$4.0\cdot 10^{-8}$
Muon mass	m_{u}	$1.88353109(16)\cdot 10^{-28}\mathrm{kg}$	$8.4\cdot 10^{-8}$
	r.	0.113 428 9168(34) u	$3.0 \cdot 10^{-8}$
		105, 658 3568(52) MeV	$4.9\cdot 10^{-8}$
Proton mass	$m_{ m p}$	$1,67262158(13)\cdot 10^{-27} \mathrm{kg}$	$7.9 \cdot 10^{-8}$
	r	1.007 276 466 88(13) u	$1.3\cdot 10^{-10}$
		938, 271 998(38) MeV	$4.0\cdot10^{-8}$
Neutron mass	$m_{ m n}$	$1,67492716(13)\cdot 10^{-27} \mathrm{kg}$	$7.9 \cdot 10^{-8}$
		1.008 664 915 78(55) u	$5.4\cdot10^{-10}$
		939, 565 330(38) MeV	$4.0\cdot 10^{-8}$
Atomic mass unit	$m_{yy} = m(^{12}C)/12 =$	1 u1.660 538 73(13) yg	$7.9 \cdot 10^{-8}$

a. Uncertainty: standard deviation of measurement errors. b. Only down to 10^{-19} m and up to 10^{26} m.

c. Defining constant.

d. All coupling constants depend on the 4-momentum transfer, as explained in the section on renormalization. Fine structure constant is the traditional name for the electromagnetic coupling con-

stant $g_{\rm em}$ in the case of a 4-momentum transfer of $Q^2 = m_{\rm e}^2 c^2$, which is the smallest one possible. At higher momentum transfers it has larger values, e.g. $g_{\rm em}(Q^2 = M_{\rm W}^2 c^2) \approx 1/128$. In contrast, the strong coupling constant has lover values at higher momentum transfers; e.g., $\alpha_s(34~{\rm GeV}) = 0.14(2)$.

Why do all these basic constants have the values they have? For any basic constant with a dimension, such as the quantum of action \hbar , the numerical value has only historical meaning. It is $1.054 \cdot 10^{-34}$ Js because of the SI definition of the joule and the second. The question why the value of a dimensional constant is not larger or smaller therefore always requires one to understand the origin of some dimensionless number giving the ratio between the constant and the corresponding natural unit that is defined with c, G, \hbar and α . More details and the values of the natural units are given later. Understanding the sizes of atoms, people, trees and stars, the duration of molecular and atomic processes, or the mass of nuclei and mountains, implies understanding the ratios between these values and the corresponding natural units. The key to understanding nature is thus the understanding of all ratios, and thus of all dimensionless constants. The quest of understanding all ratios, including the fine structure constant α itself, is completed only in the final volume of our adventure.

The basic constants yield the following useful high-precision observations.

TABLE 55 Derived physical constants.

Symbol	VALUE IN SI UNITS	Uncert.
$Z_0 = \sqrt{\mu_0/\varepsilon_0}$	376.730 313 461 77 Ω	0
$N_{ m A}$	$6.022\ 141\ 99(47)\cdot 10^{23}$	$7.9\cdot10^{-8}$
$N_{ m L}$	$2.6867775(25) \cdot 10^{23}$	$1.7\cdot 10^{-6}$
$F = N_{\rm A} e$	96 485.3415(39) C/mol	$4.0\cdot 10^{-8}$
$R = N_{\rm A}k$	8.314 472(15) J/mol K	$1.7\cdot 10^{-6}$
V = RT/p	22.413 996(39) l/mol	$1.7 \cdot 10^{-6}$
$R_{\infty} = m_{\rm e} c \alpha^2 / 2h$	$10973731.568549(83)\mathrm{m}^{-1}$	$7.6\cdot10^{-12}$
$G_0 = 2e^2/h$	$77.48091696(28)\mu\text{S}$	$3.7\cdot10^{-9}$
$\varphi_0 = h/2e$	2.067 833 636(81) pWb	$3.9 \cdot 10^{-8}$
2e/h	483.597 898(19) THz/V	$3.9 \cdot 10^{-8}$
$h/e^2 = \mu_0 c/2\alpha$	$25812.807572(95)\Omega$	$3.7 \cdot 10^{-9}$
e^2/h	$3.874045848(14)\cdot 10^{-5}\mathrm{S}$	$3.7 \cdot 10^{-9}$
$\mu_{\rm B} = e\hbar/2m_{\rm e}$	9.274 008 99(37) yJ/T	$4.0\cdot 10^{-8}$
$f_{\rm c}/B = e/2\pi m_{\rm e}$	27.992 4925(11) GHz/T	$4.0 \cdot 10^{-8}$
$r_{\rm e} = e^2/4\pi\varepsilon_0 m_{\rm e}c^2$	2.817 940 285(31) fm	$1.1\cdot 10^{-8}$
$\lambda_{\rm C} = h/m_{\rm e}c$	2.426 310 215(18) pm	$7.3\cdot10^{-9}$
$\lambda_{\rm c} = \hbar/m_{\rm e}c = r_{\rm e}/\alpha$	0.386 159 264 2(28) pm	$7.3\cdot10^{-9}$
$a_{\infty} = r_{\rm e}/\alpha^2$	52.917 720 83(19) pm	$3.7 \cdot 10^{-9}$
$h/2m_{\rm e}$	$3.636947516(27)\cdot 10^{-4}\mathrm{m}^2/\mathrm{s}$	$7.3\cdot10^{-9}$
$e/m_{\rm e}$	$1.758820174(71)\cdot 10^{11}\mathrm{C/kg}$	$4.0\cdot10^{-8}$
$\mu_{ m e}$	$-9.28476362(37) \cdot 10^{-24} \text{ J/T}$	$4.0\cdot10^{-8}$
	$Z_0 = \sqrt{\mu_0/\varepsilon_0}$ N_A N_L $F = N_A e$ $R = N_A k$ $V = RT/p$ $R_{\infty} = m_e c \alpha^2 / 2h$ $G_0 = 2e^2 / h$ $\varphi_0 = h / 2e$ $2e / h$ $h / e^2 = \mu_0 c / 2\alpha$ e^2 / h $\mu_B = e h / 2m_e$ $f_c / B = e / 2\pi m_e$ $r_e = e^2 / 4\pi \varepsilon_0 m_e c^2$ $\lambda_C = h / m_e c$	$Z_0 = \sqrt{\mu_0/\epsilon_0} \qquad 376.73031346177\Omega$ $N_A \qquad 6.02214199(47)\cdot 10^{23}$ $N_L \qquad 2.6867775(25)\cdot 10^{23}$ $F = N_A e \qquad 96485.3415(39)\text{C/mol}$ $R = N_A k \qquad 8.314472(15)\text{J/mol}\text{K}$ $V = RT/p \qquad 22.413996(39)\text{l/mol}$ $R_\infty = m_e c\alpha^2/2h \qquad 10973731.568549(83)\text{m}^{-1}$ $G_0 = 2e^2/h \qquad 77.48091696(28)\mu\text{S}$ $\varphi_0 = h/2e \qquad 2.067833636(81)\text{pWb}$ $2e/h \qquad 483.597898(19)\text{THz/V}$ $h/e^2 = \mu_0 c/2\alpha \qquad 25812.807572(95)\Omega$ $e^2/h \qquad 3.874045848(14)\cdot 10^{-5}\text{S}$ $\mu_B = e\hbar/2m_e \qquad 9.27400899(37)\text{yJ/T}$ $f_c/B = e/2\pi m_e \qquad 27.9924925(11)\text{GHz/T}$ $r_e = e^2/4\pi\epsilon_0 m_e c^2 \qquad 2.817940285(31)\text{fm}$ $\lambda_C = h/m_e c \qquad 2.426310215(18)\text{pm}$ $\lambda_C = h/m_e c \qquad 3.636947516(27)\cdot 10^{-4}\text{m}^2/\text{s}$ $e/m_e \qquad 3.636947516(27)\cdot 10^{-4}\text{m}^2/\text{s}$ $e/m_e \qquad 1.758820174(71)\cdot 10^{11}\text{C/kg}$

Vol. IV, page 181

Motion Mountain – The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christopl

TABLE 55 (Continued) Derived physical constants.

Quantity	Symbol	VALUE IN SI UNITS	Uncert.
	$\mu_{\rm e}/\mu_{ m B}$	-1.001 159 652 1869(41)	$4.1\cdot 10^{-12}$
	$\mu_{\rm e}/\mu_{ m N}$	$-1.8382819660(39)\cdot 10^3$	$2.1\cdot10^{-9}$
Electron g-factor	g_{e}	-2.0023193043737(82)	$4.1\cdot10^{-12}$
Muon magnetic moment	$\mu_{ m \mu}$	$-9.28476362(37) \cdot 10^{-24} \text{J/T}$	$4.0\cdot10^{-8}$
muon g-factor	g_{μ}	-2.002 331 8320(13)	$6.4\cdot10^{-10}$
Proton-electron mass ratio	$m_{\rm p}/m_{\rm e}$	1 836.152 6675(39)	$2.1\cdot 10^{-9}$
Proton-muon mass ratio	$m_{\rm p}/m_{\rm \mu}$	8.880 244 08(27)	$3.0\cdot10^{-8}$
Specific proton charge	$e/m_{\rm p}$	9, 578 834 $08(38) \cdot 10^7$ C/kg	$4.0\cdot10^{-8}$
Proton Compton wavelength	$\lambda_{C,p} = h/m_p c$	1.321 409 847(10) fm	$7.6\cdot10^{-9}$
Nuclear magneton	$\mu_{\rm N} = e\hbar/2m_{\rm p}$	$5.05078317(20) \cdot 10^{-27} \text{ J/T}$	$4.0\cdot10^{-8}$
Proton magnetic moment	$\mu_{ m p}$	$1.410606633(58)\cdot 10^{-26}\mathrm{J/T}$	$4.1\cdot10^{-8}$
	$\mu_{\rm p}/\mu_{\rm B}$	$1.521032203(15)\cdot 10^{-3}$	$1.0\cdot10^{-8}$
	$\mu_{ m p}/\mu_{ m N}$	2.792 847 337(29)	$1.0\cdot10^{-8}$
Proton gyromagnetic ratio	$\gamma_p = 2\mu_p/\hbar$	$2.67522212(11)\cdot 10^8\mathrm{Hz/T}$	$4.1\cdot10^{-8}$
Neutron-electron mass ratio	$m_{\rm n}/m_{\rm e}$	1 838.683 6550(40)	$2.2 \cdot 10^{-9}$
Neutron-proton mass ratio	$m_{\rm n}/m_{\rm p}$	1.001 378 418 87(58)	$5.8 \cdot 10^{-10}$
Neutron Compton wavelength	$\lambda_{\rm C,n} = h/m_{\rm n}c$	1.319 590 898(10) fm	$7.6 \cdot 10^{-9}$
Neutron magnetic moment	$\mu_{ m n}$	$-0.96623640(23)\cdot 10^{-26}\mathrm{J/T}$	$2.4\cdot10^{-7}$
	$\mu_{\rm n}/\mu_{\rm B}$	$-1.04187563(25)\cdot 10^{-3}$	$2.4 \cdot 10^{-7}$
	$\mu_{ m n}/\mu_{ m N}$	-1.913 042 72(45)	$2.4 \cdot 10^{-7}$
Stefan-Boltzmann constant	$\sigma = \pi^2 k^4 / 60 \hbar^3 c^2$	$56.70400(40)\mathrm{nW/m^2K^4}$	$7.0 \cdot 10^{-6}$
Wien's displacement constant	$b = \lambda_{\max} T$	2.897 768 6(51) mmK	$1.7 \cdot 10^{-6}$
Electron volt	eV	$1.602176462(63)\cdot 10^{-19}\mathrm{J}$	$3.9 \cdot 10^{-8}$
Bits to entropy conversion const.		10^{23} bit = 0.956 994 5(17) J/K	
TNT energy content		3.7 to 4.0 MJ/kg	$4 \cdot 10^{-2}$

a. For infinite mass of the nucleus.

Some useful properties of our local environment are given in the following table.

TABLE 56 Astronomical constants.

QUANTITY SYMBOL VALUE Tropical year 1900 a a $31556925.9747s$ Tropical year 1994 a $31556925.2s$ Mean sidereal day d $23^h56'4.09053''$ Astronomical unit b AU $149597870.691(30)km$ Light year al $9.460528173\ldots Pm$ Parsec pc $30.856775806Pm = 3.261634al$ Earth's mass $M_{\frac{1}{5}}$ $5.973(1) \cdot 10^{24}kg$				
Tropical year 1994 a $31556925.2\mathrm{s}$ Mean sidereal day d $23^h56'4.09053''$ Astronomical unit b AU $149597870.691(30)\mathrm{km}$ 149 597 870.691(30) km Light year al $9.460528173\ldots\mathrm{Pm}$ Parsec pc $30.856775806\mathrm{Pm} = 3.261634\mathrm{al}$	QUANTITY	Sумвог	Value	Christo
24	Tropical year 1900 ^a	а	31 556 925.974 7 s	ph Sch
24	Tropical year 1994	а	31 556 925.2 s	er Z
24	Mean sidereal day	d	23 ^h 56'4.090 53"	ovem
24	Astronomical unit b	AU	149 597 870.691(30) km	ber 1
24	Light year	al	9.460 528 173 Pm	997-1
Earth's mass $M_{\stackrel{\leftarrow}{\Box}}$ 5.973(1) · 10 ²⁴ kg	Parsec	pc	$30.856775806 \mathrm{Pm} = 3.261634 \mathrm{al}$	March
	Earth's mass	M_{\coloredge}	$5.973(1) \cdot 10^{24} \mathrm{kg}$	2012

TABLE 56 (Continued) Astronomical constants.

TABLE 30 (Continued) Astronomical Con-	starits.	
QUANTITY	S y м в о l	Value
Geocentric gravitational constant	GM	$3.986004418(8)\cdot 10^{14}\mathrm{m}^3/\mathrm{s}^2$
Earth's gravitational length	$l_{\uparrow} = 2GM/c^2$	8.870 056 078(16) mm
Earth's equatorial radius ^c	$R_{\rm bullet_{eq}}$	6378.1366(1) km
Earth's polar radius ^c	$R_{\stackrel{\circ}{O}p}$	6356.752(1) km
Equator-pole distance ^c	o _F	10 001.966 km (average)
Earth's flattening ^c	$e_{ m t}$	1/298.25642(1)
Earth's av. density	$ ho_{\circ}$	$5.5 \mathrm{Mg/m^3}$
Earth's age	$T_{\dot{\Box}}$	4.50(4) Ga = 142(2) Ps
Earth's normal gravity	g	$9.80665 \mathrm{m/s^2}$
Earth's standard atmospheric pressure	p_0	101 325 Pa
Moon's radius	$R_{\text{((v)}}$	1738 km in direction of Earth
Moon's radius	$R_{\text{(1)}h}$	1737.4 km in other two directions
Moon's mass	$M_{\mathbb{Q}}$	$7.35 \cdot 10^{22} \mathrm{kg}$
Moon's mean distance d	$d_{\mathbb{Q}}$	384 401 km
Moon's distance at perigee ^d		typically 363 Mm, historical minimum 359 861 km
Moon's distance at apogee ^d		typically 404 Mm, historical maximum 406 720 km
Moon's angular size ^e		average 0.5181° = 31.08′, minimum 0.49′ maximum - shortens line 0.55°
Moon's average density	$ ho_{\mathbb{Q}}$	3.3 Mg/m ³
Moon's surface gravity	90	$1.62 \mathrm{m/s^2}$
Moons's atmospheric pressure	PC	from 10^{-10} Pa (night) to 10^{-7} Pa (day)
Jupiter's mass	M_{2}	$1.90 \cdot 10^{27} \mathrm{kg}$
Jupiter's radius, equatorial	R_{2}	71.398 Mm
Jupiter's radius, polar	R_{2}	67.1(1) Mm
Jupiter's average distance from Sun	$D_{2_{+}}$	778 412 020 km
Jupiter's surface gravity	$g_{7\!+}$	$24.9 \mathrm{m/s^2}$
Jupiter's atmospheric pressure	P 2 ₊	from 20 kPa to 200 kPa
Sun's mass	M_{\odot}	$1.98843(3) \cdot 10^{30} \mathrm{kg}$
Sun's gravitational length	$l_{\odot} = 2GM_{\odot}/c^2$	2.953 250 08 km
Sun's luminosity	L_{\odot}	384.6 YW
Solar equatorial radius	R_{\odot}	695.98(7) Mm
Sun's angular size		0.53° average; minimum on fourth of July (aphelion) 1888", maximum on fourth of January (perihelion) 1952"
Sun's average density	$ ho_{\odot}$	$1.4\mathrm{Mg/m^3}$
Sun's average distance	AU	149 597 870.691(30) km
Sun's age	T_{\odot}	4.6 Ga
Solar velocity	$v_{\odot g}$	January (perihelion) 1952" 1.4 Mg/m ³ 149 597 870.691(30) km 4.6 Ga 220(20) km/s

TABLE 56 (Continued) Astronomical constants.

Quantity	Sумвоц	Value	
around centre of galaxy			
Solar velocity	$v_{\odot \mathrm{b}}$	370.6(5) km/s	
against cosmic background			
Sun's surface gravity	g_{\odot}	$274 \mathrm{m/s}^2$	
Sun's lower photospheric pressure	P⊙	15 kPa	
Distance to Milky Way's centre		$8.0(5) \mathrm{kpc} = 26.1(1.6) \mathrm{kal}$	
Milky Way's age		13.6 Ga	
Milky Way's size		c. 10 ²¹ m or 100 kal	
Milky Way's mass		10^{12} solar masses, $c. 2 \cdot 10^{42}$ kg	
Most distant galaxy cluster known	SXDF-XCLJ	9.6 · 10 ⁹ al	Motio
	0218-0510		Motion Mou

Challenge 661 s Ref. 346

- a. Defining constant, from vernal equinox to vernal equinox; it was once used to define the second. (Remember: π seconds is about a nanocentury.) The value for 1990 is about 0.7 s less, corresponding to a slowdown of roughly 0.2 ms/a. (Watch out: why?) There is even an empirical formula for the change of the length of the year over time.
- b. Average distance Earth–Sun. The truly amazing precision of 30 m results from time averages of signals sent from Viking orbiters and Mars landers taken over a period of over twenty years.
- c. The shape of the Earth is described most precisely with the World Geodetic System. The last edition dates from 1984. For an extensive presentation of its background and its details, see the www.wgs84.com website. The International Geodesic Union refined the data in 2000. The radii and the flattening given here are those for the 'mean tide system'. They differ from those of the 'zero tide system' and other systems by about 0.7 m. The details constitute a science in itself.
- d. Measured centre to centre. To find the precise position of the Moon in the sky at a given date, see the www.fourmilab.ch/earthview/moon_ap_per.html page. For the planets, see the page www.fourmilab.ch/solar/solar.html and the other pages on the same site.
- e. Angles are defined as follows: 1 degree = $1^{\circ} = \pi/180 \text{ rad}$, 1 (first) minute = $1' = 1^{\circ}/60$, 1 second (minute) = 1'' = 1'/60. The ancient units 'third minute' and 'fourth minute', each 1/60th of the preceding, are not in use any more. ('Minute' originally means 'very small', as it still does in modern English.)

Some properties of nature at large are listed in the following table. (If you want a challenge, can you determine whether any property of the universe itself is listed?)

Challenge 662 s

TABLE 57 Astrophysical constants.

Symbol	Value			
Λ	$c. 1 \cdot 10^{-52} \mathrm{m}^{-2}$			
t_0	$4.333(53) \cdot 10^{17} \text{ s} = 13.73(0.17) \cdot 10^9 \text{ a}$			
(determined from space-time, via expansion, using general relativity)				
t_0	over $3.5(4) \cdot 10^{17}$ s = $11.5(1.5) \cdot 10^9$ a			
(determined from matter, via galaxies and stars, using quantum theory)				
H_0	$2.3(2) \cdot 10^{-18} \mathrm{s}^{-1} = 0.73(4) \cdot 10^{-10} \mathrm{a}^{-1}$			
$= h_0 \cdot 100 \mathrm{km/s} \mathrm{Mpc} = h_0 \cdot 1.0227 \cdot 10^{-10} \mathrm{a}^{-1}$				
h_0	0.71(4)			
	Λ t_0 expansion, using ge t_0 dies and stars, using H_0 $= h_0 \cdot 100 \text{km/s} \text{M}$			

TABLE 57 (Continued) Astrophysical constants.

QUANTITY	SYMBOL	Value
Deceleration parameter ^a	$q_0 = -(\ddot{a}/a)_0/H_0^2$	-0.66(10)
Universe's horizon distance ^a	$d_0 = 3ct_0$	$40.0(6) \cdot 10^{26} \mathrm{m} = 13.0(2) \mathrm{Gpc}$
Universe's topology		trivial up to 10^{26} m
Number of space dimensions		3, for distances up to 10^{26} m
Critical density	$\rho_{\rm c} = 3H_0^2/8\pi G$	$h_0^2 \cdot 1.87882(24) \cdot 10^{-26} \mathrm{kg/m}^3$
of the universe		$= 0.95(12) \cdot 10^{-26} \mathrm{kg/m^3}$
(Total) density parameter ^a	$\Omega_0 = \rho_0/\rho_c$	1.02(2)
Baryon density parameter ^a	$\Omega_{\mathrm{B0}} = \rho_{\mathrm{B0}}/\rho_{\mathrm{c}}$	0.044(4)
Cold dark matter density parameter ^a	$\Omega_{\mathrm{CDM0}} = \rho_{\mathrm{CDM0}}/\rho$	o _c 0.23(4)
Neutrino density parameter ^a	$\Omega_{v0} = \rho_{v0}/\rho_{\rm c}$	0.001 to 0.05
Dark energy density parameter ^a	$\Omega_{\rm X0} = \rho_{\rm X0}/\rho_{\rm c}$	0.73(4)
Dark energy state parameter	$w = p_{\rm X}/\rho_{\rm X}$	-1.0(2)
Baryon mass	$m_{\rm b}$	$1.67 \cdot 10^{-27} \mathrm{kg}$
Baryon number density		$0.25(1)/m^3$
Luminous matter density		$3.8(2) \cdot 10^{-28} \mathrm{kg/m^3}$
Stars in the universe	$n_{\rm s}$	$10^{22\pm1}$
Baryons in the universe	$n_{\rm b}$	$10^{81\pm1}$
Microwave background temperature ^b	T_0	2.725(1) K
Photons in the universe	n_{γ}	10 ⁸⁹
Photon energy density	$\rho_{\gamma} = \pi^2 k^4 / 15 T_0^4$	$4.6 \cdot 10^{-31} \text{ kg/m}^3$
Photon number density		$410.89 \text{/cm}^3 \text{ or } 400 \text{/cm}^3 (T_0/2.7 \text{K})^3$
Density perturbation amplitude	\sqrt{S}	$5.6(1.5) \cdot 10^{-6}$
Gravity wave amplitude	\sqrt{T}	$< 0.71 \sqrt{S}$
Mass fluctuations on 8 Mpc	σ_8	0.84(4)
Scalar index	n	0.93(3)
Running of scalar index	$dn/d \ln k$	-0.03(2)
Planck length	$l_{\rm Pl} = \sqrt{\hbar G/c^3}$ $t_{\rm Pl} = \sqrt{\hbar G/c^5}$	$1.62 \cdot 10^{-35} \mathrm{m}$
Planck time	$t_{\rm Pl} = \sqrt{\hbar G/c^5}$	$5.39 \cdot 10^{-44} \mathrm{s}$
Planck mass	$m_{\rm Pl} = \sqrt{\hbar c/G}$	21.8 μg
Instants in history ^a	$t_0/t_{ m Pl}$	$8.7(2.8) \cdot 10^{60}$
Space-time points	$N_0 = (R_0/l_{\rm Pl})^3 \cdot$	$10^{244\pm1}$
inside the horizon ^a	$(t_0/t_{ m Pl})$	
Mass inside horizon	M	$10^{54\pm1} \mathrm{kg}$

a. The index 0 indicates present-day values.

b. The radiation originated when the universe was 380 000 years old and had a temperature of about 3000 K; the fluctuations ΔT_0 which led to galaxy formation are today about $16 \pm 4 \,\mu\text{K} = 6(2) \cdot 10^{-6} \,T_0$.

Useful numbers

Ref. 318	π	$3.14159265358979323846264338327950288419716939937510_{5}$
	e	$2.71828182845904523536028747135266249775724709369995_{9}$
	γ	$0.57721566490153286060651209008240243104215933593992_3$
	ln 2	$0.69314718055994530941723212145817656807550013436025_{5}$
	ln 10	$2.30258509299404568401799145468436420760110148862877_2$
	$\sqrt{10}$	$3.16227766016837933199889354443271853371955513932521_6$



APPENDIX C

SOURCES OF INFORMATION ON MOTION

No place affords a more striking conviction of the vanity of human hopes than a public library.

Samuel Johnson

In a consumer society there are inevitably two kinds of slaves: the prisoners of addiction and the prisoners of envy.

Ivan Illich*

In the text, good books that introduce neighbouring domains are presented n the bibliography. The bibliography also points to journals and websites, n order to satisfy any additional curiosity about what is encountered in this adventure. All citations can also be found by looking up the author in the name index. To find additional information, either libraries or the internet can help.

In a library, review articles of recent research appear in journals such as Reviews of Modern Physics, Reports on Progress in Physics, Contemporary Physics and Advances in Physics. Good pedagogical introductions are found in the American Journal of Physics, the European Journal of Physics and Physik in unserer Zeit.

Overviews on research trends occasionally appear in magazines such as Physics World, Physics Today, Europhysics Journal, Physik Journal and Nederlands tijdschrift voor natuurkunde. For coverage of all the sciences together, the best sources are the magazines Nature, New Scientist, Naturwissenschaften, La Recherche and Science News.

Research papers on the foundations of motion appear mainly in Physics Letters B, Nuclear Physics B, Physical Review D, Physical Review Letters, Classical and Quantum Gravity, General Relativity and Gravitation, International Journal of Modern Physics and Modern Physics Letters. The newest results and speculative ideas are found in conference proceedings, such as the Nuclear Physics B Supplements. Research articles also appear in Fortschritte der Physik, European Physical Journal, La Rivista del Nuovo Cimento, Europhysics Letters, Communications in Mathematical Physics, Journal of Mathematical Physics, Foundations of Physics, International Journal of Theoretical Physics and Journal of Physics G.

But by far the simplest and most efficient way to keep in touch with ongoing research on motion and modern physics is to use the *internet*, the international computer network. To start using the web, ask a friend who knows.**

^{*} Ivan Illich (b. 1926 Vienna, d. 2002 Bremen), Austrian theologian and social and political thinker.

^{**} It is also possible to use the internet and to download files through FTP with the help of email only. But

In the last decade of the twentieth century, the internet expanded into a combination of library, business tool, discussion platform, media collection and, above all, addiction provider. Do not use it too much. Commerce, advertising and – unfortunately – addictive material for children, youth and adults, as well as crime of all kind are also an integral part of the web. With a personal computer, a modem and free browser software, one can look for information in millions of pages of documents or destroy one's professional career through addiction. The various parts of the documents are located in various computers around the world, but the user does not need to be aware of this.*

Most theoretical physics papers are available free of charge, as *preprints*, i.e., before official publication and checking by referees, at the arxiv.org website. A service for finding subsequent preprints that cite a given one is also available.

There are a few internet physics journals: One is Living Reviews in Relativity, found at www.livingreviews.org, the other is the New Journal of Physics, which can be found at the www.njp.org website.

On the internet, papers on the description of motion without time and space which appear *after* this text is published can be found via the Web of Science, a site accessible only from libraries. It allows one to search for all publications which *cite* a given paper.

Searching the web for authors, organizations, books, publications, companies or simple keywords using search engines can be a rewarding experience or an episode of addiction, depending entirely on yourself. A selection of interesting servers are given below.

TABLE 58 Some interesting sites on the world-wide web.

Торіс	Website address
General knowledge	
Wikipedia Usenet discssions	www.wikipedia.org groups.google.com

the tools change too often to give a stable guide here. Ask your friend.

Illich develops the idea that if such a system were informal – he then calls it a 'learning web' or 'opportunity web' – it would be superior to formal, state-financed institutions, such as conventional schools, for the development of mature human beings. These ideas are deepened in his following works, *Deschooling Our Lives*, Penguin, 1976, and *Tools for Conviviality*, Penguin, 1973.

Today, any networked computer offers *email* (electronic mail), FTP (file transfer to and from another computer), access to *usenet* (the discussion groups on specific topics, such as particle physics), and the *world-wide web*. (Roughly speaking, each of those includes the ones before.) In a rather unexpected way, all these facilities of the internet have transformed it into the backbone of the 'opportunity web' discussed by Illich. However, as in any school, it strongly depends on the user's discipline whether the internet actually does provide a learning web or an entry into addiction.

^{*} Several decades ago, the provocative book by IVAN ILLICH, *Deschooling Society*, Harper & Row, 1971, listed four basic ingredients for any educational system:

^{1.} access to *resources* for learning, e.g. books, equipment, games, etc. at an affordable price, for everybody, at any time in their life;

^{2.} for all who want to learn, access to *peers* in the same learning situation, for discussion, comparison, cooperation and competition;

^{3.} access to elders, e.g. teachers, for their care and criticism towards those who are learning;

^{4.} exchanges between students and *performers* in the field of interest, so that the latter can be models for the former. For example, there should be the possibility to listen to professional musicians and reading the works of specialist writers. This also gives performers the possibility to share, advertise and use their skills.

Торіс WEBSITE ADDRESS

Frequently asked questions on www.faqs.org

physics and other topics

Book collections www.ulib.org

books.google.com

Science News www.sciencenews.org Interesting science www.max-wissen.de Entertaining and professional thehappyscientist.com

science education by Robert

Krampf

Innovation in science and

technology

www.innovations-report.de

Physics

Learning physics with toys

from rubbish

www.arvindguptatoys.com

Official SI unit website www.bipm.fr

Unit conversion www.chemie.fu-berlin.de/chemistry/general/units.html

Particle data pdg.web.cern.ch Engineering data and formulae www.efunda.com

Information on relativity math.ucr.edu/home/baez/relativity.html

Research preprints

www.slac.stanford.edu/spires

Abstracts of papers in physics www.osti.gov

journals

www.aip.org/physnews/update

Physics news, daily www.innovations-report.de/berichte/physik.php

Physics problems by Yacov

KantorKantor, Yacov

Physics news, weekly

www.tau.ac.il/~kantor/QUIZ/

Physics problems by Henry

 $www.phy.duke.edu/{\sim}hsg/physics-challenges/challenges.html$

Greenside

Physics 'question of the week'

www.physics.umd.edu/lecdem/outreach/QOTW/active

Physics 'miniproblem' www.nyteknik.se/miniproblemet Physics stack exchange physics.stackexchange.com

'Ask the experts' www.sciam.com/askexpert_directory.cfm

Nobel Prize winners www.nobel.se/physics/laureates Pictures of physicists www.if.ufrj.br/famous/physlist.html

www.cern.ch Physics organizations

www.hep.net www.nikhef.nl

www.het.brown.edu/physics/review/index.html

Physics textbooks on the web www.plasma.uu.se/CED/Book

www.biophysics.org/education/resources.htm

	<
	2
	\leq
	2
	unta
	음.
)
	The
	D
	Ad
	é
	큳
	0
	of F
	f Ph
,	SYI
	CS.
	S
	0
	읔
	₫
	P
	SVB
	/aila
	20
	ë
	=
	99
	0
	ha
	D.E
	ē
	2
	S
	⋛
	ww.mo
	3
	읔
	otionmo
	≝
	S
	a
	ıtain.r
	net
	Ť
	\cap
,	ρΫ́
(글.
	Ħ
	hris
	15
	do
	Š
	S
	8
	= e
	Z
	2
	emb
	Φ,
	1997
	97
	Ĭ
	≤a
	7
	20
	_

Торіс	Website address
	www.lightandmatter.com
	www.motionmountain.net
Three beautiful French sets of notes on classical mechanics and particle theory	feynman.phy.ulaval.ca/marleau/notesdecours.htm
The excellent <i>Radical Freshman Physics</i> by David Raymond	www.physics.nmt.edu/~raymond/teaching.html
Physics course scripts from MIT	ocw.mit.edu/OcwWeb/Physics/index.html
Physics lecture scripts in German and English	www.akleon.de
'World lecture hall'	wlh.webhost.utexas.edu
Living Reviews in Relativity	www.livingreviews.org
Wissenschaft in die Schulen	www.wissenschaft-schulen.de
Mathematics	
'Math forum' internet resource collection	e mathforum.org/library
Biographies of mathematician	s www-history.mcs.st-andrews.ac.uk/BiogIndex.html
Purdue math problem of the week	www.math.purdue.edu/academics/pow
Macalester College maths problem of the week	mathforum.org/wagon
Mathematical formulae	dlmf.nist.gov
Weisstein's World of Mathematics	mathworld.wolfram.com
Functions	functions.wolfram.com
Symbolic integration	www.integrals.com
Algebraic surfaces	www.mathematik.uni-kl.de/~hunt/drawings.html
Astronomy	
ESA	sci.esa.int
NASA	www.nasa.gov
Hubble space telescope	hubble.nasa.gov
Sloan Digital Sky Survey	skyserver.sdss.org
The 'cosmic mirror'	www.astro.uni-bonn.de/~dfischer/mirror
Solar system simulator	space.jpl.nasa.gov
Observable satellites	liftoff.msfc.nasa.gov/RealTime/JPass/20
Astronomy picture of the day	antwrp.gsfc.nasa.gov/apod/astropix.html
The Earth from space	www.visibleearth.nasa.gov
From Stargazers to Starships	www.phy6.org/stargaze/Sintro.htm
Current solar data	www.n3kl.org/sun

>
0
ĕ.
9
Ž
0
\simeq
unta
ם
⋾
#
je He
~
d
ĕ
ntu
ĕ
9
7
ihysi
S
Ö
흑
file
P
رو
Vα
<u>=</u>
8
ĕ
=
(D)
Ø
9
\cap
ha
e,
۵
_
≶
ww.mo
?
3
2
omn
3
H
ta
Ξ.
ntain.ne
et
\bigcirc
Cop
二.
9
7
0
0
Ħ
hristo
Ö
ž
Sch
Ė
9
No
9
é
ğ
Ö
9
_
1997-
7
Mar
0
9

Торіс	WEBSITE ADDRESS
Specific topics	
Minerals	webmineral.com
	www.mindat.org
Geological Maps	onegeology.org
Optical illusions	www.sandlotscience.com
Petit's science comics	www.jp-petit.org
Physical toys	www.e20.physik.tu-muenchen.de/~cucke/toylinke.htm
Physics humour	www.dctech.com/physics/humor/biglist.php
Literature on magic	www.faqs.org/faqs/magic-faq/part2
Making paper aeroplanes	www.pchelp.net/paper_ac.htm
	www.ivic.qc.ca/~aleexpert/aluniversite/klinevogelmann.html
Small flying helicopters	pixelito.reference.be
Science curiosities	www.wundersamessammelsurium.info
Ten thousand year clock	www.longnow.org
Gesellschaft Deutscher	www.gdnae.de
Naturforscher und Ärzte	
Pseudoscience	suhep.phy.syr.edu/courses/modules/PSEUDO/pseudo_main. html
Crackpots	www.crank.net
Mathematical quotations	math.furman.edu/mwoodard/~mquot.html
The 'World Question Center'	www.edge.org/questioncenter.html
Plagiarism	www.plagiarized.com
Hoaxes	www.museumofhoaxes.com

Do you want to study physics without actually going to university? Nowadays it is possible to do so via email and internet, in German, at the University of Kaiserslautern.* In the near future, a nationwide project in Britain should allow the same for English-speaking students. As an introduction, use the latest update of this physics text!

Das Internet ist die offenste Form der geschlossenen Anstalt.**

Matthias Deutschmann

Si tacuisses, philosophus mansisses.***


After Boethius.



^{*} See the www.fernstudium-physik.de website.

^{** &#}x27;The internet is the most open form of a closed institution.'

^{*** &#}x27;If you had kept quiet, you would have remained a philosopher.' After the story Boethius (c. 480–c. 525) tells in *De consolatione philosophiae*, 2.7, 67 ff.



CHALLENGE HINTS AND SOLUTIONS

Never make a calculation before you know the answer.

John Wheeler's motto

John Wheeler wanted people to estimate, to try and to guess; but not saying the guess out loud. A correct guess reinforces the physics instinct, whereas a wrong one leads to the pleasure of surprise. Guessing is thus an important first step in solving every problem.

Teachers have other criteria to keep in mind. Good problems can be solved on different levels of difficulty, can be solved with words or with images or with formulae, activate knowledge, concern real world applications, and are open.

Challenge 1, page 9: Do not hesitate to be demanding and strict. The next edition of the text will benefit from it.

Challenge 2, page 16: There are many ways to distinguish real motion from an illusion of motion: for example, only real motion can be used to set something else into motion. In addition, the motion illusions of the figures show an important failure; nothing moves if the head and the paper remain fixed with respect to each other. In other words, the illusion only *amplifies* existing motion, it does not *create* motion from nothing.

Challenge 3, page 17: Without detailed and precise experiments, both sides can find examples to prove their point. Creation is supported by the appearance of mould or bacteria in a glass of water; creation is also supported by its opposite, namely traceless disappearance, such as the disappearance of motion. However, conservation is supported and creation falsified by all those investigations that explore assumed cases of appearance or disappearance in full detail.

Challenge 4, page 19: The amount of water depends on the shape of the bucket. The system chooses the option (tilt or straight) for which the centre of gravity is lowest.

Challenge 5, page 19: Political parties, sects, helping organizations and therapists of all kinds are typical for this behaviour.

Challenge 6, page 24: The issue is not yet completely settled for the motion of empty space, such as in the case of gravitational waves. Thus, the motion of empty space might be an exception. In any case, empty space is not made of small particles of finite size, as this would contradict the transversality of gravity waves.

Challenge 7, page 26: The circular definition is: objects are defined as what moves with respect to the background, and the background is defined as what stays when objects change. We shall return to this important issue several times in our adventure. It will require a certain amount of patience to solve it, though.

Challenge 8, page 27: Holes are not physical systems, because in general they cannot be tracked.

Challenge 9, page 27: See the discussion in volume IV, on quantum theory.

Page 369

Vol. IV, page 146




FIGURE 244
A soap
bubble while
bursting
(© Peter
Wienerroither).

Challenge 10, page 29: Hint: yes, there is such a point.

Challenge 11, page 30: See Figure 244 for an intermediate step. A bubble bursts at a point, and then the rim of the hole increases rapidly, until it disappears on the antipodes. During that process the remaining of the bubble keeps its spherical shape, as shown in the figure. For a film of the process, see www.youtube.com/watch?v=SpcXtmkk26Q. In other words, the final droplets that are ejected stem from the point of the bubble which is opposite to the point of puncture; they are never ejected from the centre of the bubble.

Challenge 12, page 30: A ghost can be a moving image; it cannot be a moving object, as objects cannot interpenetrate.

Vol. IV, page 117

Challenge 13, page 30: If something could stop moving, motion could disappear into nothing. For a precise proof, one would have to show that no atom moves any more. So far, this has never been observed: motion is conserved. (Nothing in nature can disappear into nothing.)

Challenge 14, page 30: This would indeed mean that space is infinite; however, it is impossible to observe that something moves "forever": nobody lives that long. In short, there is no way to prove that space is infinite in this way. In fact, there is no way to prove that space if infinite in any other way either.

Challenge 15, page 30: The necessary rope length is nh, where n is the number of wheels/pulleys. And yes, the farmer is indeed doing something sensible.

Challenge 16, page 30: How would you measure this?

Challenge 17, page 30: The number of reliable digits of a measurement result is a simple quantification of precision. More details can be found by looking up 'standard deviation' in the index.

Challenge 18, page 30: No; memory is needed for observation and measurements. This is the case for humans and measurement apparatus. Quantum theory will make this particularly clear.

Challenge 19, page 30: Note that you never have observed zero speed. There is always some measurement error which prevents one to say that something is zero. No exceptions!

Challenge 20, page 30: $(2^{64} - 1) = 18446744073700551615$ grains of wheat, with a grain weight of 40 mg, are 738 thousand million tons. Given a world harvest in 2006 of 606 million

tons, the grains amount to about 1200 years of the world's wheat harvests.

The grain number calculation is simplified by using the formula $1 + m + m^2 + m^3 + ...m^n = (m^{n+1}-1)/(m-1)$, that gives the sum of the so-called *geometric sequence*. (The name is historical and is used as a contrast to the *arithmetic sequence* 1 + 2 + 3 + 4 + 5 + ...n = n(n+1)/2.) Can you prove the two expressions?

The chess legend is mentioned first by Abu-l 'Abbas Ahmand Ibn Khallikan (1211–1282).. King Shiram and king Balhait, also mentioned in the legend, are historical figures that lived between the second and fourth century CE. The legend appears to have combined two different stories. Indeed, the calculation of grains appears already in the year 947, in the famous text *Meadows of Gold and Mines of Precious Stones* by Abu ul-Hasan 'Ali ibn Husayn ibn 'Ali ul-Mas'udi.

Challenge 21, page 30: In clean experiments, the flame leans inwards. But such experiments are not easy, and sometimes the flame leans outwards. Just try it. Can you explain both observations?

Challenge 22, page 30: Accelerometers are the simplest motion detectors. They exist in form of piezoelectric devices that produce a signal whenever the box is accelerated and can cost as little as one euro. Another accelerometer that might have a future is an interference accelerometer that makes use of the motion of an interference grating; this device might be integrated in silicon. Other, more precise accelerometers use gyroscopes or laser beams running in circles.

Velocimeters and position detectors can also detect motion; they need a wheel or at least an optical way to look out of the box. Tachographs in cars are examples of velocimeters, computer mice are examples of position detectors.

A cheap enough device would be perfect to measure the speed of skiers or skaters. No such device exists yet.

Challenge 23, page 31: The ball rolls (or slides) towards the centre of the table, as the table centre is somewhat nearer to the centre of the Earth than the border; then the ball shoots over, performing an oscillation around the table centre. The period is 84 min, as shown in challenge 355. (This has never been observed, so far. Why?)

Challenge 24, page 31: Only if the acceleration never vanishes. Accelerations can be felt. Accelerometers are devices that measure accelerations and then deduce the position. They are used in aeroplanes when flying over the atlantic. If the box does not accelerate, it is impossible to say whether it moves or sits still. It is even impossible to say in which direction one moves. (Close your eyes in a train at night to confirm this.)

Challenge 25, page 31: The block moves twice as fast as the cylinders, independently of their radius.

Challenge 26, page 31: This methods is known to work with other fears as well.

Challenge 27, page 31: Three couples require 11 passages. Two couples require 5. For four or more couples there is no solution. What is the solution if there are n couples and n-1 places on the boat?

Challenge 28, page 32: Hint: there is an infinite number of such shapes. These curves are called also *Reuleaux curves*. Another hint: The 20 p and 50 p coins in the UK have such shapes. And yes, other shapes than cylinders are also possible: take a twisted square bar, for example.

Challenge 29, page 32: Conservation, relativity and minimization are valid generally. In some rare processes in nuclear physics, motion invariance is broken, as is mirror invariance. Continuity is known not to be valid at smallest length and time intervals, but no experiments has yet probed those domains, so that it is still valid in practice.

Challenge 30, page 33: In everyday life, this is correct; what happens when quantum effects are taken into account?

Challenge 31, page 35: Take the average distance change of two neighbouring atoms in a piece

of quartz over the last million years. Do you know something still slower?

Challenge 32, page 36: There is only one way: compare the velocity to be measured with the speed of light. In fact, almost all physics textbooks, both for schools and for university, start with the definition of space and time. Otherwise excellent relativity textbooks have difficulties avoiding this habit, even those that introduce the now standard k-calculus (which is in fact the approach mentioned here). Starting with speed is the logically cleanest approach.

Challenge 33, page 36: There is no way to sense your own motion if you are in vacuum. No way in principle. This result is often called the *principle of relativity*.

In fact, there is a way to measure your motion in space (though not in vacuum): measure your speed with respect to the cosmic background radiation. So we have to be careful about what is implied by the question.

Challenge 34, page 36: The wing load W/A, the ratio between weight W and wing area A, is obviously proportional to the third root of the weight. (Indeed, $W \sim l^3$, $A \sim l^2$, l being the dimension of the flying object.) This relation gives the green trend line.

The wing load W/A, the ratio between weight W and wing area A, is, like all forces in fluids, proportional to the square of the cruise speed v: we have $W/A = v^2 0.38 \, \text{kg/m}^3$. The unexplained factor contains the density of air and a general numerical coefficient that is difficult to calculate. This relation connects the upper and lower horizontal scales in the graph.

As a result, the cruise speed scales as the *sixth root* of weight: $v \sim W^{1/6}$. In other words, an Airbus A380 is 750 000 million times heavier than a fruit fly, but only a hundred times as fast.

Challenge 35, page 40: Equivalently: do points in space exist? The final part of our ascent studies this issue in detail.

Vol. VI, page 59

Page 141

Challenge 36, page 41: All electricity sources must use the same phase when they feed electric power into the net. Clocks of computers on the internet must be synchronized.

Challenge 37, page 41: Note that the shift increases quadratically with time, not linearly.

Challenge 38, page 41: Natural time is measured with natural motion. Natural motion is the motion of light. Natural time is thus defined with the motion of light.

Challenge 39, page 42: Galileo measured time with a scale (and with other methods). His stopwatch was a water tube that he kept closed with his thumb, pointing into a bucket. To start the stopwatch, he removed his thumb, to stop it, he put it back on. The volume of water in the bucket then gave him a measure of the time interval. This is told in his famous book Galileo Galilei, *Discorsi e dimostrazioni matematiche intorno a due nuove scienze attenenti alla mecanica e i movimenti locali*, usually simply called the 'Discorsi', which he published in 1638 with Louis Elsevier in Leiden, in the Netherlands.

Challenge 40, page 47: There is no way to define a local time at the poles that is consistent with all neighbouring points. (For curious people, check the website www.arctic.noaa.gov/gallery_np. html.)

Challenge 42, page 48: The forest is full of light and thus of light rays: they are straight, as shown by the sunbeams in Figure 245.

Challenge 43, page 49: One pair of muscles moves the lens along the third axis by deforming the eye from prolate to spherical to oblate.

Challenge 44, page 49: You can solve this problem by trying to think in four dimensions. (Train using the well-known three-dimensional projections of four-dimensional cubes.) Try to imagine how to switch the sequence when two pieces cross. Note: it is usually *not* correct, in this domain, to use time instead of a fourth *spatial* dimension!

Challenge 45, page 51: Measure distances using light.



FIGURE 245 Sunbeams in a forest (© Fritz Bieri and Heinz Rieder).

Challenge 48, page 54: It is easier to work with the unit torus. Take the unit interval [0,1] and equate the end points. Define a set B in which the elements are a given real number b from the interval plus all those numbers who differ from that real by a rational number. The unit circle can be thought as the union of all the sets B. (In fact, every set B is a shifted copy of the rational numbers Q.) Now build a set A by taking one element from each set B. Then build the set family consisting of the set A and its copies A_q shifted by a rational A. The union of all these sets is the unit torus. The set family is countably infinite. Then divide it into A0 two countably infinite set families. It is easy to see that each of the two families can be renumbered and its elements shifted in such a way that each of the two families forms a unit torus.

Mathematicians say that there is no countably infinitely additive measure of \mathbb{R}^n or that sets such as A are non-measurable. As a result of their existence, the 'multiplication' of lengths is possible. Later on we shall explore whether bread or *gold* can be multiplied in this way.

Challenge 49, page 55: Hint: start with triangles.

Ref. 41

Challenge 50, page 55: An example is the region between the x-axis and the function which assigns 1 to every transcendental and 0 to every non-transcendental number.

Challenge 51, page 56: We use the definition of the function of the text. The dihedral angle of a regular tetrahedron is an irrational multiple of π , so the tetrahedron has a non-vanishing Dehn invariant. The cube has a dihedral angle of $\pi/2$, so the Dehn invariant of the cube is 0. Therefore, the cube is not equidecomposable with the regular tetrahedron.

Challenge 52, page 56: If you think you can show that empty space is continuous, you are wrong. Check your arguments. If you think you can prove the opposite, you *might* be right – but only if you already know what is explained in the final part of the text. If that is not the case, check your arguments.

Challenge 53, page 57: Obviously, we use light to check that the plumb line is straight, so the two definitions must be the same. This is the case because the field lines of gravity are also possible paths for the motion of light. However, this is not always the case; can you spot the exceptions?

Another way to check straightness is along the surface of calm water.

A third, less precise way, way is to make use of the straightness sensors on the brain. The human brain has a built-in faculty to determine whether an objects seen with the eyes is straight. There are special cells in the brain that fire when this is the case. Any book on vision perception tells more about this topic.




FIGURE 246 A simple way to measure bullet speeds.




FIGURE 247 How to make a hole in a postcard that allows stepping through it.

Challenge 54, page 58: The hollow Earth theory is correct if the distance formula is used consistently. In particular, one has to make the assumption that objects get smaller as they approach the centre of the hollow sphere. Good explanations of all events are found on www.geocities.com/inversedearth. Quite some material can be found on the internet, also under the names of celestrocentric system, inner world theory or concave Earth theory. There is no way to prefer one description over the other, except possibly for reasons of simplicity or intellectual laziness.

Challenge 56, page 58: A hint is given in Figure 246. For the measurement of the speed of light with almost the same method, see volume II, on page 18.

Challenge 57, page 58: A fast motorbike is faster: a driver can catch an arrow, a stunt that was shown on a television show in Germany around the year 2000.

Challenge 58, page 59: 72 stairs.

Challenge 61, page 59: See Figure 247.

Challenge 62, page 60: Within 1 per cent, one *fifth* of the height must be empty, and four fifths must be filled; the exact value follows from $\sqrt[3]{2} = 1.25992...$

Challenge 63, page 60: One pencil draws a line of between 20 and 80 km, if no lead is lost when sharpening. Numbers for the newly invented plastic, flexible pencils are unknown.

Challenge 64, page 60: The bear is white, because the obvious spot of the house is at the North pole. But there are infinitely many additional spots (without bears) near the South pole: can you




FIGURE 248 Two ways to lengthen a rope around the Earth.

find them?

Challenge 65, page 60: We call L the initial length of the rubber band, v the speed of the snail relative to the band and V the speed of the horse relative to the floor. The speed of the snail relative to the floor is given as

$$\frac{\mathrm{d}s}{\mathrm{d}t} = v + V \frac{s}{L + Vt} \ . \tag{121}$$

This is a so-called *differential equation* for the unknown snail position s(t). You can check – by simple insertion – that its solution is given by

$$s(t) = \frac{v}{V}(L + Vt)\ln(1 + Vt/L) . \tag{122}$$

Therefore, the snail reaches the horse at a time

$$t_{\text{reaching}} = \frac{L}{V} (e^{V/v} - 1) \tag{123}$$

which is finite for all values of L, V and v. You can check however, that the time is very large indeed, if realistic speed values are used.

Challenge 66, page 61: Colour is a property that applies only to objects, not to boundaries. In the mentioned case, only spots and backgrounds have colours. The question shows that it is easy to ask questions that make no sense also in physics.

Challenge 67, page 61: You can do this easily yourself. You can even find websites on the topic.

Challenge 69, page 61: Clocks with two hands: 22 times. Clocks with three hands: 2 times.

Challenge 70, page 61: For two hands, the answer is 143 times.

Challenge 71, page 61: The Earth rotates with 15 minutes per minute.

Challenge 72, page 61: You might be astonished, but no reliable data exist on this question. The highest speed of a throw measured so far seems to be a 45 m/s cricket bowl. By the way, much more data are available for speeds achieved with the help of rackets. The *c*. 70 m/s of fast badminton smashes seem to be a good candidate for record racket speed; similar speeds are achieved by golf balls.

Page 204




FIGURE 249 Leaving a parking space – the outer turning radius.

Challenge 73, page 61: A *spread out* lengthening by 1 m allows even many cats to slip through, as shown on the left side of Figure 248. But the right side of the figure shows a better way to use the extra rope length, as Dimitri Yatsenko points out: a *localized* lengthening by 1 mm then already yields a height of 1.25 m, allowing a child to walk through. In fact, a lengthening by 1 m performed in this way yields a peak height of 121 m!

Challenge 74, page 61: 1.8 km/h or 0.5 m/s.

Challenge 76, page 62: The different usage reflects the idea that we are able to determine our position by ourselves, but not the time in which we are. The section on determinism will show how wrong this distinction is.

Challenge 77, page 62: Yes, there is. However, this is not obvious, as it implies that space and time are not continuous, in contrast to what we learn in primary school. The answer will be found in the final part of this text.

Challenge 78, page 62: For a curve, use, at each point, the curvature radius of the circle approximating the curve in that point; for a surface, define two directions in each point and use two such circles along these directions.

Challenge 79, page 62: It moves about 1 cm in 50 ms.

Challenge 80, page 62: The surface area of the lung is between 100 and $200 \,\mathrm{m}^2$, depending on the literature source, and that of the intestines is between 200 and $400 \,\mathrm{m}^2$.

Challenge 81, page 63: A limit does not exist in classical physics; however, there is one in nature which appears as soon as quantum effects are taken into account.

Challenge 82, page 63: The final shape is a full cube without any hole.

Challenge 83, page 63: The required gap d is

$$d = \sqrt{(L-b)^2 - w^2 + 2w\sqrt{R^2 - (L-b)^2}} - L + b , \qquad (124)$$

as deduced from Figure 249. See also R. HOYLE, Requirements for a perfect s-shaped parallel parking maneuvre in a simple mathematical model, 2003.

Challenge 84, page 63: A smallest gap does not exist: any value will do! Can you show this?

Challenge 85, page 63: The following solution was proposed by Daniel Hawkins.

Assume you are sitting in car A, parked behind car B, as shown in Figure 250. There are two basic methods for exiting a parking space that requires the reverse gear: rotating the car to move the centre of rotation away from (to the right of) car B, and shifting the car downward to move the




FIGURE 250 Solving the car parking puzzle (© Daniel Hawkins).

centre of rotation away from (farther below) car B. The first method requires car A to be partially diagonal, which means that the method will not work for d less than a certain value, essentially the value given above, when no reverse gear is needed. We will concern ourselves with the second method (pictured), which will work for an infinitesimal d.

In the case where the distance d is less than the minimum required distance to turn out of the parking space without using the reverse gear for a given geometry L, w, b, R, an attempt to turn out of the parking space will result in the corner of car A touching car B at a distance T away from the edge of car B, as shown in Figure 250. This distance T is the amount by which car A must be translated downward in order to successfully turn out of the parking space.

The method to leave the parking space, shown in the top left corner of Figure 250, requires two phases to be successful: the initial turning phase, and the straightening phase. By turning and straightening out, we achieve a vertical shift downward and a horizontal shift left, while preserving the original orientation. That last part is key because if we attempted to turn until the corner of car A touched car B, car A would be rotated, and any attempt to straighten out would just follow the same arc backward to the initial position, while turning the wheel the other direction would rotate the car even more, as in the first method described above.

Our goal is to turn as far as we can and still be able to completely straighten out by time car A touches car B. To analyse just how much this turn should be, we must first look at the properties of a turning car.

Ackermann steering is the principle that in order for a car to turn smoothly, all four wheels must rotate about the same point. This was patented by Rudolph Ackermann in 1817. Some properties of Ackermann steering in relation to this problem are as follows:

- The back wheels stay in alignment, but the front wheels (which we control), must turn different amounts to rotate about the same centre.
- The centres of rotation for left and right turns are on opposite sides of the car
- For equal magnitudes of left and right turns, the centres of rotation are equidistant from the nearest edge of the car. Figure 250 makes this much clearer.
- All possible centres of rotation are on the same line, which also always passes through the back wheels.
- When the back wheels are "straight" (straight will always mean in the same orientation as the initial position), they will be vertically aligned with the centres of rotation.
- When the car is turning about one centre, say the one associated with the maximum left turn, then the potential centre associated with the maximum right turn will rotate along with the car. Similarly, when the cars turns about the right centre, the left centre rotates.

Now that we know the properties of Ackermann steering, we can say that in order to maximize the shift downward while preserving the orientation, we must turn left about the 1st centre such that the 2nd centre rotates a *horizontal* distance d, as shown in Figure 250. When this is achieved, we brake, and turn the steering wheel the complete opposite direction so that we are now turning right about the 2nd centre. Because we shifted leftward d, we will straighten out at the exact moment car A comes in contact with car B. This results in our goal, a downward shift m and leftward shift d while preserving the orientation of car A. A similar process can be performed in reverse to achieve another downward shift m and a m and a m while preserving its orientation. This can be done indefinitely, which is why it is possible to get out of a parking space with an infinitesimal d between car A and car B. To determine how many times this procedure (both sets of turning and straightening) must be performed, we must only divide T (remember T is the amount by which car A must be shifted downward in order to turn out of the parking spot normally) by 2m, the total downward shift for one iteration of the procedure. Symbolically,

$$n = \frac{T}{2m} \ . \tag{125}$$

In order to get an expression for n in terms of the geometry of the car, we must solve for T and




FIGURE 251 A simple drawing – one of the many possible one – that allows proving Pythagoras' theorem.




FIGURE 252 The trajectory of the middle point between the two ends of the hands of a clock.

2m. To simplify the derivations we define a new length x, also shown in Figure 250.

$$x = \sqrt{R^2 - (L - b)^2}$$

$$T = \sqrt{R^2 - (L - b + d)^2} - x + w$$
$$= \sqrt{R^2 - (L - b + d)^2} - \sqrt{R^2 - (L - b)^2} + w$$

$$m = 2x - w - \sqrt{(2x - w)^2 - d^2}$$

$$= 2\sqrt{R^2 - (L - b)^2} - w - \sqrt{(2\sqrt{R^2 - (L - b)^2} - w)^2 - d^2}$$

$$= 2\sqrt{R^2 - (L - b)^2} - w - \sqrt{4(R^2 - (L - b)^2) - 4w\sqrt{R^2 - (L - b)^2} + w^2 - d^2}$$

$$= 2\sqrt{R^2 - (L - b)^2} - w - \sqrt{4R^2 - 4(L - b)^2 - 4w\sqrt{R^2 - (L - b)^2} + w^2 - d^2}$$

We then get

$$n = \frac{T}{2m} = \frac{\sqrt{R^2 - (L - b + d)^2} - \sqrt{R^2 - (L - b)^2} + w}{4\sqrt{R^2 - (L - b)^2} - 2w - 2\sqrt{4R^2 - 4(L - b)^2} - 4w\sqrt{R^2 - (L - b)^2} + w^2 - d^2}} \; .$$

The value of n must always be rounded up to the next integer to determine how many times one must go backward and forward to leave the parking spot.

Challenge 86, page 63: Nothing, neither a proof nor a disproof.

Challenge 87, page 64: See volume II, on page 18. On extreme shutters, see also the discussion in Volume VI, on page 111.




FIGURE 253 The angles defined by the hands against the sky, when the arms are extended.

Challenge 88, page 64: A hint for the solution is given in Figure 251.

Challenge 89, page 64: Because they are or were liquid.

Challenge 90, page 64: The shape is shown in Figure 252; it has eleven lobes.

Challenge 91, page 65: The cone angle φ , the angle between the cone axis and the cone border (or equivalently, *half* the apex angle of the cone) is related to the solid angle Ω through the relation $\Omega = 2\pi(1 - \cos \varphi)$. Use the surface area of a spherical cap to confirm this result.

Challenge 93, page 65: See Figure 253.

Challenge 97, page 66: Hint: draw all objects involved.

Challenge 98, page 66: The curve is obviously called a *catenary*, from Latin 'catena' for chain. The formula for a catenary is $y = a \cosh(x/a)$. If you approximate the chain by short straight segments, you can make wooden blocks that can form an arch without any need for glue. The St. Louis arch is in shape of a catenary. A suspension bridge has the shape of a catenary before it is loaded, i.e., before the track is attached to it. When the bridge is finished, the shape is in between a catenary and a parabola.

Challenge 99, page 67: The inverse radii, or curvatures, obey $a^2+b^2+c^2+d^2=(1/2)(a+b+c+d)^2$. This formula was discovered by René Descartes. If one continues putting circles in the remaining spaces, one gets so-called circle packings, a pretty domain of recreational mathematics. They have many strange properties, such as intriguing relations between the coordinates of the circle centres and their curvatures.

Challenge 100, page 67: One option: use the three-dimensional analogue of Pythagoras's theorem. The answer is 9.

Challenge 101, page 68: There are two solutions. (Why?) They are the two positive solutions of $l^2 = (b+x)^2 + (b+b^2/x)^2$; the height is then given as h = b+x. The two solutions are 4.84 m and 1.26 m. There are closed formulas for the solutions; can you find them?

Challenge 102, page 68: The best way is to calculate first the height B at which the blue ladder touches the wall. It is given as a solution of $B^4 - 2hB^3 - (r^2 - b^2)B^2 + 2h(r^2 - b^2)B - h^2(r^2 - b^2) = 0$. Integer-valued solutions are discussed in Martin Gardner, *Mathematical Circus*, Spectrum, 1996.

Challenge 103, page 68: Draw a logarithmic scale, i.e., put every number at a distance corresponding to its natural logarithm. Such a device, called a *slide rule*, is shown in Figure 254. Slide




FIGURE 254 A high-end slide rule, around 1970 (© Jörn Lütjens).

rules were the precursors of electronic calculators; they were used all over the world in prehistoric times, i.e., until around 1970. See also the web page www.oughtred.org.

Challenge 104, page 68: Two more. Build yourself a model of the Sun and the Earth to verify this

Challenge 105, page 68: The Sun is exactly behind the back of the observer; it is setting, and the rays are coming from behind and reach deep into the sky in the direction opposite to that of the Sun.

Challenge 107, page 69: The volume is given by $V = \int A dx = \int_{-1}^{1} 4(1 - x^2) dx = 16/3$.

Challenge 108, page 69: Yes. Try it with a paper model.

Challenge 109, page 69: Problems appear when quantum effects are added. A two-dimensional universe would have no matter, since matter is made of spin 1/2 particles. But spin 1/2 particles do not exist in two dimensions. Can you find other reasons?

Challenge 110, page 70: Two dimensions do not allow ordering of events. To say 'before' and 'afterwards' becomes impossible. In everyday life and all domains accessible to measurement, time is surely one-dimensional.

Challenge 111, page 70: No experiment has ever found any hint. Can this be nevertheless? Probably not, as argued in the last volume of Motion Mountain.

Challenge 112, page 70: The best solution seems to be 23 extra lines. Can you deduce it? To avoid spoiling the fun of searching, no solution is given here. You can find solutions on blog.vixra.org/ 2010/12/26/a-christmas-puzzle.

Challenge 113, page 70: If you solve this so-called ropelength problem, you will become a famous mathematician. The length is known only with about 6 decimals of precision. No exact formula is known, and the exact shape of such ideal knots is unknown for all non-trivial knots. The problem is also unsolved for all non-trivial ideal *closed* knots, for which the two ends are glued together.

Challenge 114, page 73: From $x = gt^2/2$ you get the following rule: square the number of seconds, multiply by five and you get the depth in metres.

Challenge 115, page 73: Just experiment.

Challenge 116, page 74: The Academicians suspended one cannon ball with a thin wire just in front of the mouth of the cannon. When the shot was released, the second, flying cannon ball

flew through the wire, thus ensuring that both balls started at the same time. An observer from far away then tried to determine whether both balls touched the Earth at the same time. The experiment is not easy, as small errors in the angle and air resistance confuse the results.

Challenge 117, page 74: A parabola has a so-called focus or focal point. All light emitted from that point and reflected exits in the same direction: all light ray are emitted in parallel. The name 'focus' – Latin for fireplace – expresses that it is the hottest spot when a parabolic mirror is illuminated. Where is the focus of the parabola $y = x^2$? (Ellipses have two foci, with a slightly different definition. Can you find it?)

Challenge 118, page 75: The long jump record could surely be increased by getting rid of the sand stripe and by measuring the true jumping distance with a photographic camera; that would allow jumpers to run more closely to their top speed. The record could also be increased by a small inclined step or by a spring-suspended board at the take-off location, to increase the take-off angle.

Challenge 119, page 75: Walk or run in the rain, measure your own speed v and the angle from the vertical α with which the rain appears to fall. Then the speed of the rain is $v_{\text{rain}} = v/\tan \alpha$.

Challenge 121, page 76: Neglecting air resistance and approximating the angle by 45°, we get $v = \sqrt{dg}$, or about 3.8 m/s. This speed is created by a steady pressure build-up, using blood pressure, which is suddenly released with a mechanical system at the end of the digestive canal. The cited reference tells more about the details.

Challenge 122, page 76: On horizontal ground, for a speed v and an angle from the horizontal α , neglecting air resistance and the height of the thrower, the distance d is $d = v^2 \sin 2\alpha/g$.

Challenge 123, page 76: Check your calculation with the information that the 1998 world record is juggling with 9 balls.

Challenge 124, page 76: It is said so, as rain drops would then be ice spheres and fall with high speed.

Challenge 125, page 76: There are conflicting statements in the literature. But it is a fact that people have gone to hospital and even died because a falling bullet went straight through their head. (See S. MIRSKY, *It is high, it is far*, Scientific American p. 86, February 2004, or C. Tuijn, *Vallende kogels*, Nederlands tijdschrift voor natuurkunde 71, pp. 224–225, 2005.) In addition, the lead in bullets is bad for the environment.

Challenge 126, page 76: This is a true story. The answer can only be given if it is known whether the person had the chance to jump while running or not. In the case described by R. Cross, *Forensic physics 101: falls from a height*, American Journal of Physics 76, pp. 833–837, 2008, there was no way to run, so that the answer was: murder.

Challenge 127, page 76: For jumps of an animal of mass m the necessary energy E is given as E = mgh, and the work available to a muscle is roughly speaking proportional to its mass $W \sim m$. Thus one gets that the height h is independent of the mass of the animal. In other words, the specific mechanical energy of animals is around $1.5 \pm 0.7 \,\text{J/kg}$.

Challenge 128, page 77: Stones *never* follow parabolas: when studied in detail, i.e., when the change of g with height is taken into account, their precise path turns out to be an ellipse. This shape appears most clearly for long throws, such as throws around a sizeable part of the Earth, or for orbiting objects. In short, stones follow parabolas only if the Earth is assumed to be flat. If its curvature is taken into account, they follow ellipses.

Challenge 129, page 77: The set of all rotations around a point in a plane is indeed a vector space. What about the set of all rotations around *all* points in a plane? And what about the three-dimensional cases?

Challenge 132, page 78: The scalar product between two vectors a and b is given by

$$ab = ab\cos \triangleleft (a, b). \tag{126}$$

How does this differ from the vector product?

Challenge 135, page 81: A candidate for low acceleration of a physical system might be the accelerations measured by gravitational wave detectors. They are below 10^{-13} m/s².

Challenge 136, page 82: In free fall (when no air is present) or inside a space station orbiting the Earth, one is accelerated but does not feel anything. In fact, this indistinguishability or equivalence between acceleration and 'feeling nothing' was an essential step for Albert Einstein in his development of general relativity.

Challenge 137, page 82: Professor to student: What is the derivative of velocity? Acceleration! What is the derivative of acceleration? I don't know. *Jerk*! The fourth, fifth and sixth derivatives of position are sometimes called *snap*, *crackle* and *pop*.

Challenge 139, page 84: One can argue that any source of light must have finite size.

Challenge 141, page 85: What the unaided human eye perceives as a tiny black point is usually about $50 \, \mu m$ in diameter.

Challenge 142, page 85: See volume III, page 142.

Challenge 143, page 85: One has to check carefully whether the conceptual steps that lead us to extract the concept of point from observations are correct. It will be shown in the final part of the adventure that this is not the case.

Challenge 144, page 85: One can rotate the hand in a way that the arm makes the motion described. See also volume IV, page 114.

Challenge 145, page 85: Any number, without limit.

Challenge 146, page 85: The blood and nerve supply is not possible if the wheel has an axle. The method shown to avoid tangling up connections only works when the rotating part has *no* axle: the 'wheel' must float or be kept in place by other means. It thus becomes impossible to make a wheel *axle* using a single piece of skin. And if a wheel without an axle could be built (which might be possible), then the wheel would periodically run over the connection. Could such a axle-free connection realize a propeller?

By the way, it is still thinkable that animals have wheels on axles, if the wheel is a 'dead' object. Even if blood supply technologies like continuous flow reactors were used, animals could not make such a detached wheel grow in a way tuned to the rest of the body and they would have difficulties repairing a damaged wheel. Detached wheels cannot be grown on animals; they must be dead.

Challenge 147, page 87: The brain in the skull, the blood factories inside bones or the growth of the eye are examples.

Challenge 148, page 87: In 2007, the largest big wheels for passengers are around 150 m in diameter. The largest wind turbines are around 125 m in diameter. Cement kilns are the longest wheels: they can be over 300 m along their axis.

Challenge 149, page 87: Air resistance reduces the maximum distance, which is achieved for an angle of about $\pi/4 = 45^{\circ}$, from around $v^2/g = 91.7$ m down to around 50 m.

Challenge 153, page 90: One can also add the Sun, the sky and the landscape to the list.

Challenge 154, page 91: There is no third option. Ghosts, hallucinations, Elvis sightings, or extraterrestrials must all be objects or images. Also shadows are only special types of images.

Challenge 155, page 91: The issue was hotly discussed in the seventeenth century; even Galileo argued for them being images. However, they are objects, as they can collide with other objects,




FIGURE 255 The measured (black bars) and calculated behaviour (coloured lines) – more precisely, the dynamical eigenvalues – of a bicycle as a function of its speed (© Arend Schwab).

as the spectacular collision between Jupiter and the comet Shoemaker-Levy 9 in 1994 showed. In the meantime, satellites have been made to collide with comets and even to shoot at them (and hitting).

Challenge 156, page 92: The minimum speed is roughly the one at which it is possible to ride without hands. If you do so, and then *gently* push on the steering wheel, you can make the experience described above. Watch out: too strong a push will make you fall badly.

The *bicycle* is one of the most complex mechanical systems of everyday life, and it is still a subject of research. And obviously, the world experts are Dutch. An overview of the behaviour of a bicycle is given in Figure 255. The main result is that the bicycle is stable in the upright position at a range of medium speeds. Only at low and at large speeds must the rider actively steer to ensure upright position of the bicycle.

For more details, see J. P. Meijaard, J. M. Papadopoulos, A. Ruina & A. L. Schwab, Linearized dynamics equations for the balance and steer of a bicycle: a benchmark and review, Proceedings of the Royal Society A 463, pp. 1955–1982, 2007, and J. D. G. Kooijman, A. L. Schwab & J. P. Meijaard, Experimental validation of a model of an uncontrolled bicycle, Multibody System Dynamics 19, pp. 115–132, 2008. See also the audiophile.tam.cornell.edu/~als93/Bicycle/index.htm website.

Challenge 157, page 94: The weight decreased due to the evaporated water lost by sweating and, to a minor degree, due to the exhaled carbon bound in carbon dioxide.

Challenge 158, page 95: If the moving ball is not rotating, after the collision the two balls will

depart with a *right* angle between them.

Page 159

Challenge 159, page 95: Part of the energy is converted into heat; the rest is transferred as kinetic energy of the concrete block. As the block is heavy, its speed is small and easily stopped by the human body. This effect works also with anvils, it seems. In another common variation the person does not lie on nails, but on air: he just keeps himself horizontal, with head and shoulders on one chair, and the feet on a second one.

Challenge 160, page 96: Yes, the definition of mass works also for magnetism, because the precise condition is not that the interaction is central, but that the interaction realizes a more general condition that includes accelerations such as those produced by magnetism. Can you deduce the condition from the definition of mass as that quantity that keeps momentum conserved?

Challenge 161, page 97: Rather than using the inertial effects of the Earth, it is easier to deduce its mass from its gravitational effects.

Challenge 165, page 98: At first sight, relativity implies that tachyons have imaginary mass; however, the imaginary factor can be extracted from the mass–energy and mass–momentum relation, so that one can define a real mass value for tachyons; as a result, faster tachyons have smaller energy and smaller momentum. Both momentum and energy can be a negative number of any size.

Challenge 166, page 100: The leftmost situation has a tiny effect, the second makes the car role forward and backward, the right two pictures show ways to open wine bottles without bottle opener.

Challenge 167, page 100: Legs are never perfectly vertical; they would immediately glide away. Once the cat or the person is on the floor, it is almost impossible to stand up again.

Challenge 168, page 100: Momentum (or centre of mass) conservation would imply that the environment would be accelerated into the opposite direction. Energy conservation would imply that a huge amount of energy would be transferred between the two locations, melting everything in between. Teleportation would thus contradict energy and momentum conservation.

Challenge 169, page 101: The part of the tides due to the Sun, the solar wind, and the interactions between both magnetic fields are examples of friction mechanisms between the Earth and the Sun

Challenge 170, page 101: With the factor 1/2, increase of (physical) kinetic energy is equal to the (physical) work performed on a system: total energy is thus conserved only if the factor 1/2 is added

Challenge 172, page 103: It is a smart application of momentum conservation.

Challenge 173, page 103: Neither. With brakes on, the damage is higher, but still equal for both cars.

Challenge 174, page 104: Heating systems, transport engines, engines in factories, steel plants, electricity generators covering the losses in the power grid, etc. By the way, the richest countries in the world, such as Sweden or Switzerland, consume only half the energy per inhabitant as the USA. This waste is one of the reasons for the lower average standard of living in the USA.

Challenge 179, page 108: Just throw it into the air and compare the dexterity needed to make it turn around various axes.

Challenge 180, page 108: Use the definition of the moment of inertia and Pythagoras' theorem for every mass element of the body.

Challenge 181, page 108: Hang up the body, attaching the rope in two different points. The crossing point of the prolonged rope lines is the centre of mass.

Challenge 182, page 108: See Tables 19 and 20.

Challenge 183, page 108: Spheres have an orientation, because we can always add a tiny spot on their surface. This possibility is not given for microscopic objects, and we shall study this situation in the part on quantum theory.

Challenge 184, page 111: Yes, the ape can reach the banana. The ape just has to turn around its own axis. For every turn, the plate will rotate a bit towards the banana. Of course, other methods, like blowing at a right angle to the axis, peeing, etc., are also possible.

Challenge 185, page 111: Self-propelled linear motion contradicts the conservation of momentum; self-propelled change of orientation (as long as the motion stops again) does not contradict any conservation law. But the deep, final reason for the difference will be unveiled in the final part of our adventure.

Challenge 187, page 112: The points that move exactly along the radial direction of the wheel form a circle below the axis and above the rim. They are the points that are sharp in Figure 70 of page 112.

Challenge 188, page 112: Use the conservation of angular momentum around the point of contact. If all the wheel's mass is assumed in the rim, the final rotation speed is half the initial one; it is independent of the friction coefficient.

Challenge 190, page 113: Probably the 'rest of the universe' was meant by the writer. Indeed, a moving a part never shifts the centre of gravity of a closed system. But is the universe closed? Or a system? The last part of our adventure covers these issues.

Challenge 191, page 113: The human body is more energy-efficient at low and medium power output. The topic is still subject of research, as detailed in the cited reference. The critical slope is estimated to be around 16° for uphill walkers, but should differ for downhill walkers.

Challenge 192, page 114: Hint: an energy per distance is a force.

Challenge 193, page 114: The conservation of angular momentum saves the glass. Try it.

Challenge 194, page 114: First of all, MacDougall's experimental data is flawed. In the six cases MacDougall examined, he did not know the exact timing of death. His claim of a mass decrease cannot be deduced from his own data. Modern measurements on dying sheep, about the same mass as humans, have shown no mass change, but clear weight pulses of a few dozen grams when the heart stopped. This temporary weight decrease could be due to the expelling of air or moisture, to the relaxing of muscles, or to the halting of blood circulation. The question is not settled.

Challenge 196, page 115: Assuming a square mountain, the height h above the surrounding crust and the depth d below are related by

$$\frac{h}{d} = \frac{\rho_{\rm m} - \rho_{\rm c}}{\rho_{\rm c}} \tag{127}$$

where ρ_c is the density of the crust and ρ_m is the density of the mantle. For the density values given, the ratio is 6.7, leading to an additional depth of 6.7 km below the mountain.

Challenge 200, page 116: The behaviour of the spheres can only be explained by noting that elastic waves propagate through the chain of balls. Only the propagation of these elastic waves, in particular their reflection at the end of the chain, explains that the same number of balls that hit on one side are lifted up on the other. For long times, friction makes all spheres oscillate in phase. Can you confirm this?

Challenge 201, page 116: When the short cylinder hits the long one, two compression waves start to run from the point of contact through the two cylinders. When each compression wave arrives at the end, it is reflected as an expansion wave. If the geometry is well chosen, the expansion wave coming back from the short cylinder can continue into the long one (which is still in

his compression phase). For sufficiently long contact times, waves from the short cylinder can thus depose much of their energy into the long cylinder. Momentum is conserved, as is energy; the long cylinder is oscillating in length when it detaches, so that not all its energy is translational energy. This oscillation is then used to drive nails or drills into stone walls. In commercial hammer drills, length ratios of 1:10 are typically used.

Challenge 202, page 117: The momentum transfer to the wall is double when the ball rebounds perfectly.

Challenge 203, page 117: If the cork is in its intended position: take the plastic cover off the cork, put the cloth around the bottle (this is for protection reasons only) and repeatedly hit the bottle on the floor or a fall in an inclined way, as shown in Figure 64 on page 100. With each hit, the cork will come out a bit.

If the cork has fallen inside the bottle: put half the cloth inside the bottle; shake until the cork falls unto the cloth. Pull the cloth out: first slowly, until the cloth almost surround the cork, and then strongly.

Challenge 205, page 117: The atomic force microscope.

Challenge 207, page 118: Running man: $E \approx 0.5 \cdot 80 \text{ kg} \cdot (5 \text{ m/s})^2 = 1 \text{ kJ}$; rifle bullet: $E \approx 0.5 \cdot 0.04 \text{ kg} \cdot (500 \text{ m/s})^2 = 5 \text{ kJ}$.

Challenge 208, page 118: It almost doubles in size.

Challenge 209, page 118: At the highest point, the acceleration is $g \sin \alpha$, where α is the angle of the pendulum at the highest point. At the lowest point, the acceleration is v^2/l , where l is the length of the pendulum. Conservation of energy implies that $v^2 = 2gl(1 - \cos \alpha)$. Thus the problem requires that $\sin \alpha = 2(1 - \cos \alpha)$. This results in $\cos \alpha = 3/5$.

Challenge 210, page 118: One needs the mass change equation $dm/dt = \pi \rho_{\text{vapour}} r^2 |v|$ due to the vapour and the drop speed evolution $m \, dv/dt = mg - v \, dm/dt$. These two equations yield

$$\frac{\mathrm{d}v^2}{\mathrm{d}r} = \frac{2g}{C} - 6\frac{v^2}{r} \tag{128}$$

where $C = \rho_{\text{vapour}}/4\rho_{\text{water}}$. The trick is to show that this can be rewritten as

$$r\frac{d}{dr}\frac{v^2}{r} = \frac{2g}{C} - 7\frac{v^2}{r} \,. \tag{129}$$

For large times, all physically sensible solutions approach $v^2/r = 2g/7C$; this implies that for large times,

$$\frac{\mathrm{d}v}{\mathrm{d}t}\frac{v^2}{r} = \frac{g}{7} \quad \text{and} \quad r = \frac{gC}{14}t^2 \,. \tag{130}$$

About this famous problem, see for example, B. F. Edwards, J. W. Wilder & E. E. Scime, *Dynamics of falling raindrops*, European Journal of Physics 22, pp. 113–118, 2001, or A. D. Sokal, *The falling raindrop, revisited*, preprint at arxiv.org/abs/09080090.

Challenge 213, page 118: Weigh the bullet and shoot it against a mass hanging from the ceiling. From the mass and the angle it is deflected to, the momentum of the bullet can be determined.

Challenge 215, page 119: The curve described by the midpoint of a ladder sliding down a wall is a circle.

Challenge 216, page 119: The switched use the power that is received when the switch is pushed and feed it to a small transmitter that acts a high frequency remote control to switch on the light.

Challenge 217, page 119: A clever arrangement of bimetals is used. They move every time the temperature changes from day to night – and vice versa – and wind up a clock spring. The clock itself is a mechanical clock with low energy consumption.

Challenge 218, page 119: The weight of the lift does not change at all when a ship enters it. A twin lift, i.e., a system in which both lifts are mechanically or hydraulically connected, needs no engine at all: it is sufficient to fill the upper lift with a bit of additional water every time a ship enters it. Such ship lifts without engines at all used to exist in the part.

Challenge 220, page 121: This is not easy; a combination of friction and torques play a role. See for example the article J. Sauer, E. Schörner & C. Lennerz, *Real-time rigid body simulation of some classical mechanical toys*, 10th European Simulation and Symposium and Exhibition (ESS '98) 1998, pp. 93–98, or http://www.lennerz.de/paper_ess98.pdf.

Challenge 223, page 126: The Coriolis effect can be seen as the sum two different effects of equal magnitude. The first effect is the following: on a rotating background, velocity changes over time. What an inertial (nonrotating) observer sees as a *constant* velocity will be seen a velocity that *changes* over time by the rotating observer. The acceleration seen by the rotating observer is negative, and is proportional to the angular velocity and to the velocity.

The second effect is change of velocity in space. In a rotating frame of reference, different points have different velocities. The effect is negative, and proportional to the angular velocity and to the velocity.

In total, the Coriolis acceleration (or Coriolis effect) is thus $a_C = -2\omega \times v$.

Challenge 224, page 127: A *short* pendulum of length L that swings in two dimensions (with amplitude ρ and orientation φ) shows two additional terms in the Lagrangian \mathcal{L} :

$$\mathcal{L} = T - V = \frac{1}{2}m\dot{\rho}^2 \left(1 + \frac{\rho^2}{L^2}\right) + \frac{l_z^2}{2m\rho^2} - \frac{1}{2}m\omega_0^2 \rho^2 \left(1 + \frac{\rho^2}{4L^2}\right)$$
 (131)

where as usual the basic frequency is $\omega_0^2 = g/L$ and the angular momentum is $l_z = m\rho^2\dot{\phi}$. The two additional terms disappear when $L\to\infty$; in that case, if the system oscillates in an ellipse with semiaxes a and b, the ellipse is fixed in space, and the frequency is ω_0 . For *finite* pendulum length L, the frequency changes to

$$\omega = \omega_0 \left(1 - \frac{a^2 + b^2}{16 L^2} \right) \,. \tag{132}$$

The ellipse turns with a frequency

$$\Omega = \omega \frac{3}{8} \frac{ab}{L^2} \,. \tag{133}$$

These formulae can be derived using the least action principle, as shown by C. G. Gray, G. Karl & V. A. Novikov, *Progress in classical and quantum variational principles*, arxiv.org/abs/physics/0312071. In other words, a short pendulum in elliptical motion shows a precession even *without* the Coriolis effect. Since this precession frequency diminishes with $1/L^2$, the effect is small for long pendulums, where only the Coriolis effect is left over. To see the Coriolis effect in a short pendulum, one thus has to avoid that it starts swinging in an elliptical orbit by adding a mechanism that suppresses elliptical motion.

Challenge 225, page 128: The Coriolis acceleration is the reason for the deviation from the straight line. The Coriolis acceleration is due to the change of speed with distance from the rotation axis. Now think about a pendulum, located in Paris, swinging in the North-South direction with amplitude A. At the Southern end of the swing, the pendulum is further from the axis by $A\sin\varphi$, where φ is the latitude. At that end of the swing, the central support point overtakes the pendulum bob with a relative horizontal speed given by $v=2\pi A\sin\varphi/23\,\text{h56}$ min. The period of precession is given by $T_F=v/2\pi A$, where $2\pi A$ is the circumference $2\pi A$ of the envelope of the




FIGURE 256 Deducing the expression for the Sagnac effect.

pendulum's path (relative to the Earth). This yields $T_{\rm F} = 23\,{\rm h}56\,{\rm min/\sin\phi}$. Why is the value that appears in the formula not 24 h, but 23 h56 min?

Challenge 226, page 128: The axis stays fixed with respect to distant stars, not with respect to absolute space (which is an entity that cannot be observed at all).

Challenge 227, page 128: Rotation leads to a small frequency and thus colour changes of the circulating light.

Challenge 228, page 128: The weight changes when going east or when moving west due to the Coriolis acceleration. If the rotation speed is tuned to the oscillation frequency of the balance, the effect is increased by resonance. This trick was also used by Eötvös.

Challenge 229, page 128: The Coriolis acceleration makes the bar turn, as every moving body is deflected to the side, and the two deflections add up in this case. The direction of the deflection depends on whether the experiments is performed on the northern or the southern hemisphere.

Challenge 230, page 130: When rotated by π around an east–west axis, the Coriolis force produces a drift velocity of the liquid around the tube. It has the value

$$v = 2\omega r \sin \theta, \tag{134}$$

as long as friction is negligible. Here ω is the angular velocity of the Earth, θ the latitude and r the (larger) radius of the torus. For a tube with 1 m diameter in continental Europe, this gives a speed of about $6.3 \cdot 10^{-5}$ m/s.

The measurement can be made easier if the tube is restricted in diameter at one spot, so that the velocity is increased there. A restriction by an area factor of 100 increases the speed by the same factor. When the experiment is performed, one has to carefully avoid any other effects that lead to moving water, such as temperature gradients across the system.

Challenge 231, page 130: Imagine a circular light path (for example, inside a circular glass fibre) and two beams moving in opposite directions along it, as shown in Figure 256. If the fibre path rotates with rotation frequency Ω , we can deduce that, after one turn, the difference ΔL in path length is

$$\Delta L = 2R\Omega t = \frac{4\pi R^2 \Omega}{c} \ . \tag{135}$$

The phase difference is thus

$$\Delta \varphi = \frac{8\pi^2 R^2}{G \lambda} \Omega \tag{136}$$

if the refractive index is 1. This is the required formula for the main case of the Sagnac effect.

It is regularly suggested that the Sagnac effect can only be understood with help of general relativity; this is wrong. As just done, the effect is easily deduced from the invariance of the speed of light *c*. The effect is a consequence of special relativity.

Challenge 232, page 131: The metal rod is slightly longer on one side of the axis. When the wire keeping it up is burned with a candle, its moment of inertia decreases by a factor of 10^4 ; thus it starts to rotate with (ideally) 10^4 times the rotation rate of the Earth, a rate which is easily visible by shining a light beam on the mirror and observing how its reflection moves on the wall.

Challenge 233, page 137: The original result by Bessel was 0.3136", or 657.7 thousand orbital radii, which he thought to be 10.3 light years or 97.5 Pm.




FIGURE 257 How the night sky, and our galaxy in particular, looks in the near infrared (NASA false colour image).

Challenge 235, page 140: The galaxy forms a stripe in the sky. The galaxy is thus a flattened structure. This is even clearer in the infrared, as shown more clearly in Figure 257. From the flattening (and its circular symmetry) we can deduce that the galaxy must be rotating. Thus other matter must exist in the universe.

Challenge 237, page 142: See page 163.

Challenge 239, page 143: The scale reacts to your heartbeat. The weight is almost constant over time, except when the heart beats: for a short duration of time, the weight is somewhat lowered at each beat. Apparently it is due to the blood hitting the aortic arch when the heart pumps it upwards. The speed of the blood is about 0.3 m/s at the maximum contraction of the left ventricle. The distance to the aortic arch is a few centimetres. The time between the contraction and the reversal of direction is about 15 ms. And the measured weight is not even constant for a dead person, as air currents disturb the measurement.

Challenge 240, page 143: Use Figure 85 on page 126 for the second half of the trajectory, and think carefully about the first half.

Challenge 241, page 143: Hint: starting rockets at the Equator saves a lot of energy, thus of fuel and of weight.

Challenge 243, page 144: The flame leans towards the inside.

Challenge 245, page 144: For your exam it is better to say that centrifugal force does not exist. But since in each stationary system there is a force balance, the discussion is somewhat a red herring.

Challenge 247, page 144: Place the tea in cups on a board and attach the board to four long ropes that you keep in your hand.

Challenge 248, page 145: The ball leans in the direction it is accelerated to. As a result, one could imagine that the ball in a glass at rest pulls upwards because the floor is accelerated upwards. We will come back to this issue in the section of general relativity.

Challenge 249, page 145: The friction of the tides on Earth are the main cause.

Challenge 250, page 145: An earthquake with Richter magnitude of 12 is 1000 times the energy of the 1960 Chile quake with magnitude 10; the latter was due to a crack throughout the full 40 km of the Earth's crust along a length of 1000 km in which both sides slipped by 10 m with respect to each other. Only the impact of a meteorite could lead to larger values than 12.

Challenge 252, page 145: If a wedding ring rotates on an axis that is not a principal one, angular momentum and velocity are not parallel.

Challenge 253, page 146: Yes; it happens twice a year. To minimize the damage, dishes should be dark in colour.

Challenge 254, page 146: A rocket fired from the back would be a perfect defence against planes attacking from behind. However, when released, the rocket is effectively flying backwards with respect to the air, thus turns around and then becomes a danger to the plane that launched it. Engineers who did not think about this effect almost killed a pilot during the first such tests.

Challenge 255, page 146: Whatever the ape does, whether it climbs up or down or even lets himself fall, it remains at the same height as the mass. Now, what happens if there is friction at the wheel?

Challenge 256, page 146: Yes, if he moves at a large enough angle to the direction of the boat's motion.

Challenge 257, page 146: The moment of inertia is $\Theta = \frac{2}{5}mr^2$.

Challenge 258, page 146: The moments of inertia are equal also for the cube, but the values are $\Theta = \frac{1}{6}ml^2$. The efforts required to put a sphere and a cube into rotation are thus different.

Challenge 259, page 146: See the article by C. UCKE & H.-J. SCHLICHTING, *Faszinierendes Dynabee*, Physik in unserer Zeit 33, pp. 230–231, 2002.

Challenge 260, page 146: See the article by C. UCKE & H.-J. SCHLICHTING, *Die kreisende Büroklammer*, Physik in unserer Zeit 36, pp. 33–35, 2005.

Challenge 261, page 147: Yes. Can you imagine what happens for an observer on the Equator?

Challenge 262, page 147: A straight line at the zenith, and circles getting smaller at both sides. See an example on the website antwrp.gsfc.nasa.gov/apod/ap021115.html.

Challenge 264, page 148: The plane is described in the websites cited; for a standing human the plane is the vertical plane containing the two eyes.

Challenge 265, page 148: As said before, legs are simpler than wheels to grow, to maintain and to repair; in addition, legs do not require flat surfaces (so-called 'streets') to work.

Challenge 266, page 150: The staircase formula is an empirical result found by experiment, used by engineers world-wide. Its origin and explanation seems to be lost in history.

Challenge 267, page 150: Classical or everyday nature is right-left symmetric and thus requires an even number of legs. Walking on two-dimensional surfaces naturally leads to a minimum of four legs. Starfish, snails, slugs, clams, eels and snakes are among the most important exceptions for which the arguments are not valid.

Challenge 269, page 151: The length of the day changes with latitude. So does the length of a shadow or the elevation of stars at night, facts that are easily checked by telephoning a friend. Ships appear at the horizon first be showing only their masts. These arguments, together with the round shadow of the earth during a lunar eclipse and the observation that everything falls

Vol. III, page 250

downwards everywhere, were all given already by Aristotle, in his text On the Heavens. It is now known that everybody in the last 2500 years knew that the Earth is s sphere. The myth that many people used to believe in a flat Earth was put into the world – as rhetorical polemic – by Copernicus. The story then continued to be exaggerated more and more during the following centuries, because a new device for spreading lies had just been invented: book printing. Fact is that for 2500 years the vast majority of people knew that the Earth is a sphere.

Challenge 270, page 151: Robert Peary had forgotten that on the date he claimed to be at the North Pole, 6th of April 1909, the Sun is very low on the horizon, casting very long shadows, about ten times the height of objects. But on his photograph the shadows are much shorter. (In fact, the picture is taken in such a way to hide all shadows as carefully as possible.) Interestingly, he had even convinced the US congress to officially declare him the first man on the North Pole in 1911. (A rival crook had claimed to have reached it before Peary, but his photograph has the same mistake.) Peary also cheated on the travelled distances of the last few days; he also failed to mention that the last days he was pulled by his partner, Matthew Henson, because he was not able to walk any more. In fact Matthew Henson deserves more credit for that adventure than Peary. Henson, however, did not know that Peary cheated on the position they had reached.

Challenge 271, page 151: Yes, the effect has been measured for skyscrapers. Can you estimate the values?

Challenge 272, page 153: The tip of the velocity arrow, when drawn over time, produces a circle around the centre of motion.

Challenge 273, page 154: Draw a figure of the situation.

Challenge 274, page 154: Again, draw a figure of the situation.

Challenge 275, page 154: The value of the product GM for the Earth is $4.0 \cdot 10^{14} \,\mathrm{m}^3/\mathrm{s}^2$.

Challenge 276, page 155: All points can be reached for general inclinations; but when shooting horizontally in one given direction, only points on the first half of the circumference can be reached.

Challenge 277, page 156: On the moon, the gravitational acceleration is 1.6 m/s², about one sixth of the value on Earth. The surface values for the gravitational acceleration for the planets can be found on many internet sites.

Challenge 278, page 157: The Atwood machine is the answer: two almost equal masses m_1 and m_2 connected by a string hanging from a well-oiled wheel of negligible mass. The heavier one falls very slowly. Can show that the acceleration a of this 'unfree' fall is given by $a = g(m_1$ m_2)/($m_1 + m_2$)? In other words, the smaller the mass difference is, the slower the fall is.

Challenge 279, page 157: You should absolutely try to understand the origin of this expression. It allows understanding many important concepts of mechanics. The idea is that for small amplitudes, the acceleration of a pendulum of length l is due to gravity. Drawing a force diagram for a pendulum at a general angle α shows that

$$ma = -mg \sin \alpha$$

$$ml \frac{d^{2}\alpha}{dt^{2}} = -mg \sin \alpha$$

$$l \frac{d^{2}\alpha}{dt^{2}} = -g \sin \alpha .$$
(137)

For the mentioned small amplitudes (below 15°) we can approximate this to

$$l\frac{\mathrm{d}^2\alpha}{\mathrm{d}t^2} = -g\alpha \ . \tag{138}$$

This is the equation for a harmonic oscillation (i.e., a sinusoidal oscillation). The resulting motion is:

$$\alpha(t) = A\sin(\omega t + \varphi). \tag{139}$$

The amplitude A and the phase φ depend on the initial conditions; however, the oscillation frequency is given by the length of the pendulum and the acceleration of gravity (check it!):

$$\omega = \sqrt{\frac{l}{g}} \ . \tag{140}$$

(For arbitrary amplitudes, the formula is much more complex; see the internet or special mechanics books for more details.)

Challenge 280, page 157: Walking speed is proportional to l/T, which makes it proportional to $l^{1/2}$. The relation is also true for animals in general. Indeed, measurements show that the maximum walking speed (thus not the running speed) across all animals is given by

$$v_{\text{maxwalking}} = (2.2 \pm 0.2) \,\text{m}^{1/2}/\text{s} \,\sqrt{l}$$
 (141)

Challenge 283, page 159: The acceleration due to gravity is $a = Gm/r^2 \approx 5 \text{ nm/s}^2$ for a mass of 75 kg. For a fly with mass $m_{\rm fly} = 0.1 \, {\rm g}$ landing on a person with a speed of $v_{\rm fly} = 1 \, {\rm cm/s}$ and deforming the skin (without energy loss) by $d = 0.3 \, {\rm mm}$, a person would be accelerated by $a = (v^2/d)(m_{\rm fly}/m) = 0.4 \, {\rm \mu m/s}^2$. The energy loss of the inelastic collision reduces this value at least by a factor of ten.

Challenge 284, page 161: The calculation shows that a surprisingly high energy value is stored in thermal motion.

Challenge 287, page 162: The easiest way to see this is to picture gravity as a flux emanating from a sphere. This gives a $1/r^{d-1}$ dependence for the force and thus a $1/r^{d-2}$ dependence of the potential.

Challenge 289, page 163: Since the paths of free fall are ellipses, which are curves lying in a plane, this is obvious.

Challenge 290, page 165: The vector *OF* can be calculated by using OS = -(GmM/E)OP/OP and then translating the construction given in the figure into formulae. This exercise yields

$$OF = \frac{K}{mE} \tag{142}$$

where

$$K = p \times L - GMmx/x \tag{143}$$

is the so-called Runge–Lenz vector. (We have used x = OP for the position of the orbiting body, p for its momentum and L for its angular momentum. The Runge–Lenz vector K is *constant* along the orbit of a body, thus has the *same* value for any position x on the orbit. (Prove it by starting from $xK = xK\cos\theta$.) The Runge–Lenz vector is thus a *conserved* quantity in universal gravity. As a result, the vector OF is also constant in time. The Runge–Lenz vector is also often used in quantum mechanics, when calculating the energy levels of a hydrogen atom, as it appears in all problems with a 1/r potential. (In fact, the incorrect name 'Runge–Lenz vector' is due to Wolfgang Pauli; the discoverer of the vector was, in 1710, Jakob Hermann.)

Challenge 291, page 165: On orbits, see page 168.

Challenge 293, page 166: The low gravitational acceleration of the Moon, 1.6 m/s^2 , implies that gas molecules at usual temperatures can escape its attraction.




FIGURE 258 The Lagrangian points and the effective potential that produces them (NASA).

Challenge 295, page 168: A flash of light is sent to the Moon, where several Cat's-eyes have been deposited by the Lunokhod and Apollo missions. The measurement precision of the time a flash take to go and come back is sufficient to measure the Moon's distance change. For more details, see challenge 8.

Challenge 300, page 171: The Lagrangian points L4 and L5 are on the orbit, 60° before and behind the orbiting body. They are stable if the mass ratio of the central and the orbiting body is sufficiently large (above 24.9).

Challenge 301, page 171: The Lagrangian point L3 is located on the orbit, but precisely on the other side of the central body. The Lagrangian point L1 is located on the line connecting the planet with the central body, whereas L2 lies outside the orbit, on the same line. If R is the radius of the orbit, the distance between the orbiting body and the L1 and L2 point is $\sqrt[3]{m/3M}R$, giving around 4 times the distance of the Moon for the Sun-Earth system. L1, L2 and L3 are saddle points, but effectively stable orbits exist around them. Many satellites make use of these properties, including the famous WMAP satellite that measured the ripples of the big bang, which is located at the 'quiet' point L2, where the Sun, the Earth and the Moon are easily shielded and satellite temperature remains constant.

Challenge 302, page 173: This is a resonance effect, in the same way that a small vibration of a string can lead to large oscillation of the air and sound box in a guitar.

Challenge 304, page 175: The expression for the strength of tides, namely $2GM/d^3$, can be rewritten as $(8/3)\pi G\rho(R/d)^3$. Now, R/d is roughly the same for Sun and Moon, as every eclipse shows. So the density ρ must be much larger for the Moon. In fact, the ratio of the strengths (height) of the tides of Moon and Sun is roughly 7:3. This is also the ratio between the mass densities of the two bodies.

Challenge 305, page 175: The total angular momentum of the Earth and the Moon must remain constant.

Challenge 309, page 178: Unfortunately, the myth of 'passive gravitational mass' is spread by many books. Careful investigation shows that it is measured in exactly the same way as inertial mass. The two concepts are identical.



FIGURE 259 The famous 'vomit comet', a KC-135, performing a parabolic flight (NASA).

Challenge 310, page 179: Either they fell on inclined snowy mountain sides, or they fell into high trees, or other soft structures. The record was over 7 km of survived free fall. A recent case made the news in 2007 and is told in www.bbc.co.uk/jersey/content/articles/2006/12/20/michael_holmes_fall_feature.shtml.

Challenge 312, page 180: For a few thousand Euros, you can experience zero-gravity in a parabolic flight, such as the one shown in Figure 259. (Many 'photographs' of parabolic flights found on the internet are in fact computer graphics. What about this one?)

How does zero-gravity *feel*? It feels similar to floating under water, but without the resistance of the water. It also feels like the time in the air when one is diving into water. However, for cosmonauts, there is an additional feeling; when they rotate their head rapidly, the sensors for orientation in our ear are not reset by gravity. Therefore, for the first day or two, most cosmonauts have feelings of vertigo and of nausea, the so-called *space sickness*. After that time, the body adapts and the cosmonaut can enjoy the situation thoroughly.

Challenge 313, page 180: The centre of mass of a broom falls with the usual acceleration; the end thus falls faster.

Challenge 314, page 180: Just use energy conservation for the two masses of the jumper and the string. For more details, including the comparison of experimental measurements and theory, see N. Dubelaar & R. Brantjes, *De valversnelling bij bungee-jumping*, Nederlands tijdschrift voor natuurkunde 69, pp. 316–318, October 2003.

Challenge 315, page 180: About 1 ton.

Challenge 316, page 180: About 5 g.

Challenge 317, page 180: Your weight is roughly constant; thus the Earth must be round. On a flat Earth, the weight would change from place to place, depending on your distance from the border.

Challenge 318, page 180: Nobody ever claimed that the centre of mass is the same as the centre of gravity! The attraction of the Moon is negligible on the surface of the Earth.

Challenge 320, page 181: That is the mass of the Earth. Just turn the table on its head.

Challenge 322, page 182: The Moon will be about 1.25 times as far as it is now. The Sun then will slow down the Earth–Moon system rotation, this time due to the much smaller tidal friction from the Sun's deformation. As a result, the Moon will return to smaller and smaller distances to





FIGURE 260 The analemma photographed, at local noon, from January to December 2002, at the Parthenon on Athen's Acropolis, and a precision sundial (© Anthony Ayiomamitis, Stefan Pietrzik).

Earth. However, the Sun will have become a red giant by then, after having swallowed both the Earth and the Moon.

Challenge 324, page 182: As Galileo determined, for a swing (half a period) the ratio is $\sqrt{2}/\pi$. (See challenge 279). But not more than two, maybe three decimals of π can be determined in this way.

Challenge 325, page 182: Momentum conservation is not a hindrance, as any tennis racket has the same effect on the tennis ball.

Challenge 326, page 182: In fact, in velocity space, elliptic, parabolic and hyperbolic motions Ref. 148 are all described by circles. In all cases, the hodograph is a circle.

Challenge 327, page 183: This question is old (it was already asked in Newton's times) and deep. One reason is that stars are kept apart by rotation around the galaxy. The other is that galaxies are kept apart by the momentum they got in the big bang. Without the big bang, all stars would have collapsed together. In this sense, the big bang can be deduced from the attraction of gravitation and the immobile sky at night. We shall find out later that the darkness of the night sky gives a second argument for the big bang.

Challenge 328, page 184: The choice is clear once you notice that there is no section of the orbit which is concave towards the Sun. Can you show this?

Challenge 330, page 184: The escape velocity, from Earth, to leave the solar system – without help of the other planets – is 42 km/s. However, if help by the other planets is allowed, it can be less than half that value (why?).

If the escape velocity from a body were the speed of light, the body would be a black hole; not even light could escape. Black holes are discussed in detail in the volume on relativity.

Challenge 331, page 184: Using a maximal jumping height of $h=0.5\,\mathrm{m}$ on Earth and an estimated asteroid density of $\rho=3\,\mathrm{Mg/m^3}$, we get a maximum radius of $R^2=3gh/4\pi G\rho$, or $R\approx 2.4\,\mathrm{km}$.

Challenge 332, page 185: A handle of two bodies.

Challenge 335, page 185: The shape of an analemma at local noon is shown in Figure 260. The vertical extension of the analemma is due to the obliquity, i.e., the tilt of the Earth's axis (it is twice 23.45°). The horizontal extension is due to the combination of the obliquity and of the ellipticity of the orbit around the Sun. Both effects change the speed of the Earth along its orbit, leading to changes of the position of the Sun at local noon during the course of the year. The asymmetrical position of the central crossing point The shape of the analemma is also built into the shadow

Vol. II, page 240

pole of precision sundials.

Challenge 336, page 186: Capture of a fluid body is possible if it is split by tidal forces.

Challenge 337, page 186: The tunnel would be an elongated ellipse in the plane of the Equator, reaching from one point of the Equator to the point at the antipodes. The time of revolution would not change, compared to a non-rotating Earth. See A. J. Simonson, *Falling down a hole through the Earth*, Mathematics Magazine 77, pp. 171–188, June 2004.

Challenge 339, page 187: The centre of mass of the solar system can be as far as twice the radius from the centre of the Sun; it thus can be outside the Sun.

Challenge 340, page 188: First, during northern summer time the Earth moves faster around the Sun than during northern winter time. Second, shallow Sun's orbits on the sky give longer days because of light from when the Sun is below the horizon.

Challenge 341, page 188: Apart from the visibility of the Moon, no effect on humans has ever been detected. Gravitational effects – including tidal effects – electrical effects, magnetic effects and changes in cosmic rays are all swamped by other effects. Indeed the gravity of passing trucks, factory electromagnetic fields, the weather and solar activity changes have larger influences on humans than the Moon. The locking of the menstrual cycle to the moon phase is a visual effect.

Challenge 342, page 188: Distances were difficult to measure. It is easy to observe a planet that is before the Sun, but it is hard to check whether a planet is behind the Sun.

Challenge 343, page 188: See the mentioned reference.

Challenge 344, page 188: True.

Challenge 345, page 189: For each pair of opposite shell elements (drawn in yellow), the two attractions compensate.

Challenge 346, page 189: There is no practical way; if the masses on the shell could move, along the surface (in the same way that charges can move in a metal) this might be possible, provided that enough mass is available.

Challenge 349, page 189: Yes, one could, and this has been thought of many times, including by Jules Verne. The necessary speed depends on the direction of the shot with respect of the rotation of the Earth.

Challenge 350, page 189: Never. The Moon points always towards the Earth. The Earth changes position a bit, due to the ellipticity of the Moon's orbit. Obviously, the Earth shows phases.

Challenge 352, page 190: What counts is *local* verticality; with respect to it, the river always flows downhill.

Challenge 353, page 190: There are no such bodies, as the chapter of general relativity will show. **Challenge 355**, page 192: The oscillation is a purely sinusoidal, or harmonic oscillation, as the restoring force increases linearly with distance from the centre of the Earth. The period T for a

homogeneous Earth is $T = 2\pi \sqrt{R^3/GM} = 84 \text{ min.}$

Challenge 356, page 192: The period is the same for all such tunnels and thus in particular it is the same as the 84 min period that is valid also for the pole to pole tunnel. See for example, R. H. Romer, *The answer is forty-two – many mechanics problems, only one answer*, Physics Teacher 41, pp. 286–290, May 2003.

Challenge 358, page 192: There is no simple answer: the speed depends on the latitude and on other parameters.

Challenge 359, page 192: The centrifugal force must be equal to the gravitational force. Call the constant linear mass density d and the unknown length l. Then we have $GMd\int_R^{R+l} \mathrm{d}r/r^2 = \omega^2 d\int_R^{R+l} r \,\mathrm{d}r$. This gives $GMdl/(R^2+Rl) = (2Rl+l^2)\omega^2 d/2$, yielding $l = 0.14\,\mathrm{Gm}$.

Challenge 361, page 193: The inner rings must rotate faster than the outer rings. If the rings were solid, they would be torn apart. But this reasoning is true only if the rings are inside a certain limit, the so-called *Roche limit*. The Roche limit is that radius at which gravitational force F_g and tidal force F_t cancel on the surface of the satellite. For a satellite with mass m and radius r, orbiting a central mass M at distance d, we look at the forces on a small mass μ on its surface. We get the condition $Gm\mu/r^2 = 2GM\mu r/d^3$. With a bit of algebra, one gets the Roche limit

$$d_{\text{Roche}} = R \left(2 \frac{\rho_M}{\rho_m} \right)^{1/3} . \tag{144}$$

Below that distance from a central mass M, fluid satellites cannot exist. The calculation shown here is only an approximation; the actual Roche limit is about two times that value.

Challenge 365, page 197: In reality muscles keep an object above ground by continuously lifting and dropping it; that requires energy and work.

Challenge 366, page 197: The electricity consumption of a rising escalator indeed increases when the person on it walks upwards. By how much?

Challenge 367, page 197: Knowledge is power. Time is money. Now, power is defined as work per time. Inserting the previous equations and transforming them yields

$$money = \frac{work}{knowledge}, \qquad (145)$$

which shows that the less you know, the more money you make. That is why scientists have low salaries.

Challenge 370, page 200: Yes, because side wind increases the effective speed v in air due to vector addition, and because air resistance is (roughly) proportional to v^2 .

Challenge 371, page 200: The lack of static friction would avoid that the fluid stays attached to the body; the so-called boundary layer would not exist. One then would have to wing effect.

Challenge 373, page 201: True?

Challenge 375, page 202: From $dv/dt = g - v^2(1/2c_wA\rho/m)$ and using the abbreviation $c = 1/2c_wA\rho$, we can solve for v(t) by putting all terms containing the variable v on one side, all terms with t on the other, and integrating on both sides. We get $v(t) = \sqrt{gm/c} \tanh \sqrt{cg/m} t$.

Challenge 376, page 203: For extended deformable bodies, the intrinsic properties are given by the mass density – thus a function of space and time – and the state is described by the density of kinetic energy, linear and angular momentum, as well as by its stress and strain distributions.

Challenge 377, page 203: Electric charge.

Challenge 378, page 203: The phase space has 3N position coordinates and 3N momentum coordinates.

Vol. III, page 107

Challenge 379, page 203: The light mill is an example.

Challenge 380, page 204: No in both cases. If you have found reasons to answer yes, you overlooked something. Just go into more details and check whether the concepts you used apply to the universe. Also define carefully what you mean by 'universe'.

Challenge 382, page 205: A system showing energy or matter motion faster than light would imply that for such systems there are observers for which the order between cause and effect are reversed. A space-time diagram (and a bit of exercise from the section on special relativity) shows this.




FIGURE 261 The mechanism inside the south-pointing carriage.

Challenge 383, page 206: If reproducibility would not exist, we would have difficulties in checking observations; also reading the clock is an observation. The connection between reproducibility and time shall become important in the final part of our adventure.

Challenge 384, page 207: Even if surprises were only rare, each surprise would make it impossible to define time just before and just after it.

Challenge 387, page 207: Of course; moral laws are summaries of what others think or will do about personal actions.

Challenge 388, page 209: The fastest glide path between two points, the *brachistochrone*, turns out to be the *cycloid*, the curve generated by a point on a wheel that is rolling along a horizontal plane.

The proof can be found in many ways. The simplest is by Johann Bernoulli and is given on en. wikipedia.org/wiki/Brachistochrone_problem.

Challenge 392, page 210: Figure 261 shows the most credible reconstruction of a south-pointing carriage.

Challenge 393, page 210: The water is drawn up along the sides of the spinning egg. The fastest way to empty a bottle of water is to spin the water while emptying it.

Challenge 395, page 210: The right way is the one where the chimney falls like a V, not like an inverted V. See challenge 313 on falling brooms for inspiration on how to deduce the answer. It turns out that the chimney breaks (if it is not fastened to the base) at a height between half or two thirds of the total, depending at the angle at which this happens. For a complete solution of the problem, see the excellent paper G. Vareschi & K. Kamiya, *Toy models for the falling chimney*, AMerican Journal of Physics 71, pp. 1025–1031, 2003.

Challenge 403, page 220: In one dimension, the expression F = ma can be written as $-dV/dx = md^2x/dt^2$. This can be rewritten as $d(-V)/dx - d/dt[d/d\dot{x}(\frac{1}{2}m\dot{x}^2)] = 0$. This can be expanded to

 $\partial/\partial x(\frac{1}{2}m\dot{x}^2 - V(x)) - d/[\partial/\partial \dot{x}(\frac{1}{2}m\dot{x}^2 - V(x))] = 0$, which is Lagrange's equation for this case.

Challenge 405, page 220: Do not despair. Up to now, nobody has been able to imagine a universe (that is not necessarily the same as a 'world') different from the one we know. So far, such attempts have always led to logical inconsistencies.

Challenge 407, page 221: The two are equivalent since the equations of motion follow from the principle of minimum action and at the same time the principle of minimum action follows from the equations of motion.

Challenge 409, page 222: For gravity, all three systems exist: rotation in galaxies, pressure in planets and the Pauli pressure in stars. Against the strong interaction, the Pauli principle acts in nuclei and neutron stars; in neutron stars maybe also rotation and pressure complement the Pauli pressure. But for the electromagnetic interaction there are no composites other than our everyday matter, which is organized by the Pauli principle alone.

Challenge 411, page 225: Angular momentum is the change with respect to angle, whereas rotational energy is again the change with respect to time, as all energy is.

Challenge 412, page 225: Not in this way. A small change can have a large effect, as every switch shows. But a small change in the brain must be communicated outside, and that will happen roughly with a $1/r^2$ dependence. That makes the effects so small, that even with the most sensitive switches – which for thoughts do not exist anyway – no effects can be realized.

Challenge 414, page 226: This is a wrong question. T - U is not minimal, only its average is.

Challenge 415, page 226: No. A system tends to a minimum potential only if it is dissipative. One could, however, deduce that conservative systems oscillate around potential minima.

Challenge 416, page 227: The relation is

$$\frac{c_1}{c_2} = \frac{\sin \alpha_1}{\sin \alpha_2} \ . \tag{146}$$

The particular speed ratio between air (or vacuum, which is almost the same) and a material gives the *index of refraction n*:

$$n = \frac{c_1}{c_0} = \frac{\sin \alpha_1}{\sin \alpha_0} \tag{147}$$

Challenge 417, page 227: The principle for the growth of trees is simply the minimum of potential energy, since the kinetic energy is negligible. The growth of vessels inside animal bodies is minimized for transport energy; that is again a minimum principle. The refraction of light is the path of shortest time; thus it minimizes change as well, if we imagine light as moving entities moving without any potential energy involved.

Challenge 418, page 227: Special relativity requires that an invariant measure of the action exist. It is presented later in the walk.

Challenge 419, page 227: The universe is not a physical system. This issue will be discussed in detail later on.

Challenge 420, page 227: Use either the substitution $u = \tan t/2$ or use the historical trick

$$\sec \varphi = \frac{1}{2} \left(\frac{\cos \varphi}{1 + \sin \varphi} + \frac{\cos \varphi}{1 - \sin \varphi} \right) . \tag{148}$$

Challenge 421, page 227: A skateboarder in a cycloid has the same oscillation time independently of the oscillation amplitude. But a half-pipe needs to have vertical ends, in order to avoid jumping outside it. A cycloid never has a vertical end.

Vol. VI, page 99

Challenge 423, page 229: We talk to a person because we know that somebody understands us. Thus we assume that she somehow sees the same things we do. That means that observation is partly viewpoint-independent. Thus nature is symmetric.

Challenge 424, page 233: Memory works because we recognize situations. This is possible because situations over time are similar. Memory would not have evolved without this reproducibility.

Challenge 425, page 234: Taste differences are not fundamental, but due to different viewpoints and – mainly – to different experiences of the observers. The same holds for feelings and judgements, as every psychologist will confirm.

Challenge 426, page 235: The integers under addition form a group. Does a painter's set of oil colours with the operation of mixing form a group?

Challenge 428, page 235: There is only one symmetry operation: a rotation about π around the central point. That is the reason that later on the group D_4 is only called the approximate symmetry group of Figure 163.

Challenge 434, page 239: Scalar is the magnitude of any vector; thus the speed, defined as v = |v|, is a scalar, whereas the velocity v is not. Thus the length of any vector (or pseudovector), such as force, acceleration, magnetic field, or electric field, is a scalar, whereas the vector itself is not a scalar.

Challenge 437, page 240: The charge distribution of an extended body can be seen as a sum of a charge, a charge dipole, a charge quadrupole, a charge octupole, etc. The quadrupole is described by a tensor.

Compare: The inertia against motion of an extended body can be seen as sum of a mass, a mass dipole, a mass quadrupole, a mass octupole, etc. The mass quadrupole is described by the moment of inertia.

Challenge 441, page 242: The conserved charge for rotation invariance is angular momentum.

Challenge 444, page 245: An oscillation has a period in time, i.e., a discrete time translation symmetry. A wave has both discrete time and discrete space translation symmetry.

Challenge 445, page 245: Motion reversal is a symmetry for any closed system; despite the observations of daily life, the statements of thermodynamics and the opinion of several famous physicists (who form a minority though) all ideally closed systems are reversible.

Challenge 446, page 245: The symmetry group is a Lie group and called U(1), for 'unitary group in 1 dimension'.

Challenge 447, page 245: The surprising answer is no.

Challenge 448, page 245: There is no such thing as a 'perfect' symmetry.

Challenge 452, page 249: Just insert x(t) into the Lagrangian L=0, the minimum possible value for a system that transforms all kinetic energy into potential energy and vice versa.

Challenge 460, page 258: The potential energy is due to the 'bending' of the medium; a simple displacement produces no bending and thus contains no energy. Only the gradient captures the bending idea.

Challenge 462, page 259: The phase changes by π .

Challenge 464, page 260: Waves can be damped to extremely low intensities. If this is not possible, the observation is not a wave.

Challenge 465, page 260: The way to observe diffraction and interference with your naked fingers is told on page 92 in volume III.

Challenge 472, page 272: Skiers scrape snow from the lower side of each bump towards the upper side of the next bump. This leads to an upward motion of ski bumps.

Challenge 473, page 272: If the distances to the loudspeaker is a few metres, and the distance to the orchestra is 20 m, as for people with enough money, the listener at home hears it first.

Challenge 475, page 272: In general, the body moves along an ellipse (as for planets around the Sun) but with the fixed point as centre. In contrast to planets, where the Sun is in a *focus* of the ellipse and there is a perihelion and an apohelion, such a body moves *symmetrically* around the *centre* of the ellipse. In special cases, the body moves back and forward along a straight segment.

Challenge 478, page 273: The sound of thunder or of car traffic gets lower and lower in frequency with increasing distance.

Challenge 480, page 273: Neither; both possibilities are against the properties of water: in surface waves, the water molecules move in circles.

Challenge 481, page 273: Swimmers are able to cover 100 m in 48 s, or slightly better than 2 m/s. (Swimmer with fins achieve just over 3 m/s.) With a body length of about 1.9 m, the critical speed is 1.7 m/s. That is why short distance swimming depends on training; for longer distances the technique plays a larger role, as the critical speed has not been attained yet. The formula also predicts that on the 1500 m distance, a 2 m tall swimmer has a potential advantage of over 45 s on one with body height of 1.8 m. In addition, longer swimmers have an additional advantage: they swim shorter distances in pools (why?). It is thus predicted that successful long-distance swimmers will get taller and taller over time. This is a pity for a sport that so far could claim to have had champions of all sizes and body shapes, in contrast to many other sports.

Challenge 483, page 275: To reduce noise reflection and thus hall effects. They effectively diffuse the arriving wave fronts.

Challenge 485, page 275: Waves in a river are never elliptical; they remain circular.

Challenge 486, page 275: The lens is a cushion of material that is 'transparent' to sound. The speed of sound is faster in the cushion than in the air, in contrast to a glass lens, where the speed of light is slower in the glass. The shape is thus different: the cushion must look like a biconcave lens.

Challenge 487, page 275: Experiments show that the sound does not depend on air flows (find out how), but does depend on external sound being present. The sound is due to the selective amplification by the resonances resulting from the geometry of the shell shape.

Challenge 488, page 275: The Sun is always at a different position than the one we observe it to be. What is the difference, measured in angular diameters of the Sun?

Challenge 489, page 276: The $3 \times 3 \times 3$ cube has a rigid system of three perpendicular axes, on which a square can rotate at each of the 6 ends. The other squares are attaches to pieces moving around theses axes. The $4 \times 4 \times 4$ cube is different though; just find out. The limit on the segment number seems to be 6, so far. A $7 \times 7 \times 7$ cube requires varying shapes for the segments. But more than $5 \times 5 \times 5$ is not found in shops. On the other hand, the website www.oinkleburger.com/ Cube/applet allows playing with virtual cubes up to $100 \times 100 \times 100$, and more.

Challenge 491, page 277: An overview of systems being tested at present can be found in K.- U. Graw, *Energiereservoir Ozean*, Physik in unserer Zeit 33, pp. 82–88, Februar 2002. See also *Oceans of electricity – new technologies convert the motion of waves into watts*, Science News 159, pp. 234–236, April 2001.

Challenge 492, page 277: In everyday life, the assumption is usually justified, since each spot can be approximately represented by an atom, and atoms can be followed. The assumption is questionable in situations such as turbulence, where not all spots can be assigned to atoms, and most of all, in the case of motion of the vacuum itself. In other words, for gravity waves, and in particular for the quantum theory of gravity waves, the assumption is not justified.

Challenge 498, page 283: There are many. One would be that the transmission and thus reflection coefficient for waves would almost be independent of wavelength.

Challenge 499, page 284: A drop with a diameter of 3 mm would cover a surface of 7.1 m^2 with a 2 nm film.

Challenge 501, page 287: The critical height for a column of material is given by $h_{\text{crit}}^4 = \frac{\beta}{4\pi g} m \frac{E}{\rho^2}$, where $\beta \approx 1.9$ is the constant determined by the calculation when a column buckles under its own weight.

Challenge 503, page 288: One possibility is to describe particles as clouds; another is given in the last part of the text.

Challenge 505, page 294: Check your answers with the delightful text by P. Goldrich, S. Mahajan & S. Phinney, Order-of-Magnitude Physics: Understanding the World with Dimensional Analysis, Educated Guesswork, and White Lies, available on the internet.

Challenge 506, page 294: Glass shatters, glass is elastic, glass shows transverse sound waves, glass does not flow (in contrast to what many books state), not even on scale of centuries, glass molecules are fixed in space, glass is crystalline at small distances, a glass pane supported at the ends does not hang through.

Challenge 507, page 294: No metal wire allows building such a long wire. Only the idea of carbon nanotubes has raised the hope again; some dream of wire material based on them, stronger than any material known so far. However, no such material is known yet. The system faces many dangers, such as fabrication defects, lightning, storms, meteorites and space debris. All would lead to the breaking of the wires – if such wires will ever exist. But the biggest of all dangers is the lack of cash to build it.

Challenge 509, page 295: A medium-large earthquake would be generated.

Challenge 510, page 295: A stalactite contains a thin channel along its axis through which the water flows, whereas a stalagmite is massive throughout.

Challenge 511, page 295: About 1 part in a thousand.

Challenge 512, page 296: Even though the iron core of the Earth formed by collecting the iron from colliding asteroids which then sunk into the centre of the Earth, the scheme will not work today: in its youth, the Earth was much more liquid than today. The iron will most probably not sink. In addition, there is no known way to build a measurement probe that can send strong enough sound waves for this scheme. The temperature resistance is also an issue, but this may be solvable.

Challenge 514, page 298: Atoms are not infinitely hard, as quantum theory shows. Atoms are more similar to deformable clouds.

Vol. IV, page 67

Challenge 517. page 305: The constant k follows from the conservation of energy and that of mass: $k = \sqrt{2/(\rho(A_1^2/A_2^2 - 1))}$. The cross sections are denoted by A and the subscript 1 refers to any point far from the constriction, and the subscript 2 to the constriction.

Challenge 518, page 308: Some people notice that in some cases friction is too high, and start sucking at one end of the tube to get the flow started; while doing so, they can inhale or swallow gasoline, which is poisonous.

Challenge 524, page 309: The blood pressure in the feet of a standing human is about 27 kPa, double the pressure at the heart.

Challenge 525, page 310: Calculation gives $N = J/j = 0.0001 \,\text{m}^3/\text{s}/(7 \,\mu\text{m}^2 0.0005 \,\text{m/s})$, or about $6 \cdot 10^9$; in reality, the number is much larger, as most capillaries are closed at a given instant. The reddening of the face shows what happens when all small blood vessels are opened at the same time.

Challenge 526, page 310: Throwing the stone makes the level fall, throwing the water or the piece of wood leaves it unchanged.

Challenge 527, page 310: The ship rises higher into the sky. (Why?)

Challenge 531, page 310: The pumps worked in suction; but air pressure only allows 10 m of height difference for such systems.

Challenge 532, page 310: This argument is comprehensible only when one remembers that 'twice the amount' means 'twice as many molecules'.

Challenge 533, page 310: The alcohol is frozen and the chocolate is put around it.

Challenge 534, page 311: The author suggested in an old edition that a machine should be based on the same machines that throw the clay pigeons used in the sports of trap shooting and skeet. In the meantime, Lydéric Bocquet and Christophe Clanet have built such a machine, but using a different design; a picture can be found on the website lpmcn.univ-lyon1.fr/%7Elbocquet.

Challenge 535, page 311: The third component of *air* is the noble gas argon, making up about 1%. A longer list of components is given in Table 59.

Challenge 536, page 311: The *pleural cavity* between the lungs and the thorax is permanently below atmospheric pressure, usually 5 mbar, but even 10 mbar at inspiration. A hole in it, formed for example by a bullet, a sword or an accident, leads to the collapse of the lung – the so-called *pneumothorax* – and often to death. Open chest operations on people have became possible only after the surgeon Ferdinand Sauerbruch learned in 1904 how to cope with the problem. Nowadays however, surgeons keep the lung under *higher* than atmospheric pressure until everything is sealed again.

Challenge 537, page 311: The fountain shown in the figure is started by pouring water into the uppermost container. The fountain then uses the air pressure created by the water flowing downwards.

Challenge 538, page 312: Yes. The bulb will not resist two such cars though.

Challenge 539, page 312: Radon is about 8 times as heavy as air; it is he densest gas known. In comparison, Ni(CO) is 6 times, $SiCl_4$ 4 times heavier than air. Mercury vapour (obviously also a gas) is 7 times heavier than air. In comparison, bromine vapour is 5.5 times heavier than air.

Challenge 541, page 312: None.

Challenge 543, page 312: He brought the ropes into the cabin by passing them through liquid mercury.

Challenge 544, page 312: The pressure destroys the lung. Snorkeling is only possible at the water surface, not below the water!

Challenge 546, page 313: There are no official solutions for these questions; just check your assumptions and calculations carefully. The internet is full of such calculations.

Challenge 547, page 313: The soap flows down the bulb, making it thicker at the bottom and thinner at the top, until it reaches the thickness of two molecular layers. Later, it bursts.

Challenge 548, page 313: The temperature leads to evaporation of the involved liquid, and the vapour prevents the direct contact between the two non-gaseous bodies.

Challenge 549, page 313: For this to happen, friction would have to exist on the microscopic scale and energy would have to disappear.

Challenge 550, page 314: The longer funnel is empty before the short one. (If you do not believe it, try it out.) In the case that the amount of water in the funnel outlet can be neglected, one can use energy conservation for the fluid motion. This yields the famous Bernoulli equation $p/\rho + gh + v^2/2 = \text{const}$, where p is pressure, ρ the density of water, and g is 9.81 m/s². Therefore, the

TABLE 59 Gaseous composition of dry air, at present time^a (sources: NASA, IPCC).

GAS	Symbol	VOLUME PART ^b
Nitrogen	N_2	78.084 %
Oxygen	O_2	20.946 %
Argon	Ar	0.934 %
Carbon dioxide (in large part due to human pollution)	CO_2	387 ppm
Neon	Ne	18.18 ppm
Helium	He	5.24 ppm
Methane (mostly due to human pollution)	CH_4	1.79 ppm
Krypton	Kr	1.14 ppm
Hydrogen	H_2	0.55 ppm
Nitrous oxide (mostly due to human pollution)	N_2O	0.3 ppm
Carbon monoxide (partly due to human pollution)	CO	0.1 ppm
Xenon	Xe	0.087 ppm
Ozone (strongly influenced by human pollution)	O_3	0 to 0.07 ppm
Nitrogen dioxide (mostly due to human pollution)	NO_2	0.02 ppm
Iodine	I_2	0.01 ppm
Ammonia (mostly due to human pollution)	NH_3	traces
Radon	Ra	traces
Halocarbons and other fluorine compounds (all being humans pollutants)	20 types	0.0012 ppm
Mercury, other metals, sulfur compounds, other organic compounds (all being human pollutants)	numerous	concentration varies

a. Wet air can contain up to 4% water vapour, depending on the weather. Apart from gases, air can contain water droplets, ice, sand, dust, pollen, spores, volcanic ash, forest fire ash, fuel ash, smoke particles, meteoroids and cosmic ray particles. During the history of the Earth, the gaseous composition varied strongly.

speed v is higher for greater lengths h of the thin, straight part of the funnel: the longer funnel empties first.

But this is strange: the formula gives a simple free fall relation, as the *air* pressure is the same above and below and disappears from the calculation. The expression for the speed is thus independent of whether a tube is present or not. The real reason for the faster emptying of the tube is thus that a tube forces more water to flow out than the lack of a tube. Without tube, the diameter of the water flow *diminishes* during fall. With tube, it stays *constant*. This difference leads to the faster emptying for longer tubes.

Alternatively, you can look at the *water* pressure value *inside* the funnel. You will discover that the water pressure is lowest at the start of the exit tube. This internal water pressure is lower for longer tubes and sucks out the water faster in those cases.

Challenge 551, page 314: The eyes of fish are positioned in such a way that the pressure reduction by the flow is compensated by the pressure increase of the stall. By the way, their heart is positioned in such a way that it is helped by the underpressure.

b. The abbreviation *ppm* means 'part per million'.

Challenge 553, page 314: This feat has been achieved for lower mountains, such as the Monte Bianco in the Alps. At present however, there is no way to safely hover at the high altitudes of the Himalayas.

Challenge 555, page 314: Press the handkerchief in the glass, and lower the glass into the water with the opening first, while keeping the opening horizontal. This method is also used to lower people below the sea. The paper ball in the bottle will fly towards you. Blowing into a funnel will keep the ping-pong ball tightly into place, and the more so the stronger you blow. Blowing through a funnel towards a candle will make it lean towards you.

Challenge 563, page 324: In 5000 million years, the present method will stop, and the Sun will become a red giant. But it will burn for many more years after that.

Challenge 565, page 325: The answer depends on the size of the balloons, as the pressure is not a monotonous function of the size. If the smaller balloon is not too small, the smaller balloon wins

Challenge 568, page 326: Measure the area of contact between tires and street (all four) and then multiply by 200 kPa, the usual tire pressure. You get the weight of the car.

Challenge 572, page 329: If the average square displacement is proportional to time, the liquid is made of smallest particles. This was confirmed by the experiments of Jean Perrin. The next step is to deduce the number of these particles from the proportionality constant. This constant, defined by $\langle d^2 \rangle = 4Dt$, is called the diffusion constant (the factor 4 is valid for random motion in two dimensions). The diffusion constant can be determined by watching the motion of a particle under the microscope.

We study a Brownian particle of radius a. In two dimensions, its square displacement is given by

$$\langle d^2 = \rangle \frac{4kT}{\mu} t , \qquad (149)$$

where k is the Boltzmann constant and T the temperature. The relation is deduced by studying the motion of a particle with drag force $-\mu v$ that is subject to random hits. The linear drag coefficient μ of a sphere of radius a is given by

$$\mu = 6\pi \eta a , \qquad (150)$$

where η is the kinematic viscosity. In other words, one has

$$k = \frac{6\pi\eta a}{4T} \frac{\langle d^2 \rangle}{t} \ . \tag{151}$$

All quantities on the right can be measured, thus allowing to determine the Boltzmann constant k. Since the ideal gas relation shows that the ideal gas constant R is related to the Boltzmann constant by $R = N_{\rm A}k$, the Avogadro constant $N_{\rm A}$ that gives the number of molecules in a mole is also found in this way.

Challenge 577, page 336: The possibility of motion inversion for all observed phenomena is indeed a fundamental property of nature. It has been confirmed for all interactions and all experiments every performed. Independent of this is the fact, that realizing the inversion might be extremely hard, because inverting the motion of many atoms is often not feasible.

Challenge 583, page 339: We will find out later that the universe is not a physical system; thus the concept of entropy does not apply to it. Thus the universe is neither isolated nor closed.

Challenge 585, page 340: Egg white starts to harden at lower temperature than yolk, but for complete hardening, the opposite is true. White hardens *completely* at 80°C, egg yolk hardens considerably at 66 to 68°C. Cook an egg at the latter temperature, and the feat is possible; the white

Vol. VI, page 99



FIGURE 262 A candle on Earth and in microgravity (NASA).

remains runny, but does not remain transparent, though. Note again that the cooking time plays no role, only the precise temperature value.

Challenge 587, page 340: Yes, the effect is easily noticeable.

Challenge 589, page 341: Hot air is less dense and thus wants to rise.

Challenge 590, page 341: Keep the paper wet.

Challenge 592, page 341: The air had to be dry.

Challenge 593, page 341: In general, it is impossible to draw a line through three points. Since absolute zero and the triple point of water are fixed in magnitude, it was practically a sure bet that the boiling point would not be at precisely 100°C.

Challenge 594, page 342: No, as a water molecule is heavier than that. However, if the water is allowed to be dirty, it is possible. What happens if the quantum of action is taken into account?

Challenge 595, page 342: The danger is not due to the amount of energy, but due to the time in which it is available.

Challenge 596, page 342: The internet is full of solutions.

Challenge 598, page 343: Only if it is a closed system. Is the universe closed? Is it a system? This is discussed in the final part of the mountain ascent.

Challenge 601, page 343: For such small animals the body temperature would fall too low. They could not eat fast enough to get the energy needed to keep themselves warm.

Challenge 610, page 344: It is about 10^{-9} that of the Earth.

Challenge 612, page 344: The thickness of the folds in the brain, the bubbles in the lung, the density of blood vessels and the size of biological cells.

Challenge 613, page 345: The mercury vapour above the liquid gets saturated.

Challenge 614, page 345: A dedicated NASA project studied this question. Figure 262 gives an example comparison. You can find more details on their website.

Challenge 615, page 345: The risks due to storms and the financial risks are too high.

Challenge 616, page 345: The vortex inside the tube is cold near its axis and hot in the regions away from the axis. Through the membrane in the middle of the tube (shown in Figure 226 on page 345) the air from the axis region is sent to one end and the air from the outside region to the other end. The heating of the outside region is due to the work that the air rotating inside has to do on the air outside to get a rotation that consumes angular momentum. For a detailed explanation, see the beautiful text by MARK P. SILVERMAN, And Yet it Moves: Strange Systems and Subtle Questions in Physics, Cambridge University Press, 1993, p. 221.

Challenge 620, page 346: In the case of water, a few turns mixes the ink, and turning backwards increases the mixing. In the case of glycerine, a few turns *seems* to mix the ink, and turning backwards undoes the mixing.

Challenge 621, page 346: Put them in clothes.

Challenge 623, page 346: Negative temperatures are a conceptual crutch definable only for systems with a few discrete states; they are not real temperatures, because they do not describe equilibrium states, and indeed never apply to systems with a continuum of states.

Challenge 624, page 348: This is also true for the shape of human bodies, the brain control of human motion, the growth of flowers, the waves of the sea, the formation of clouds, the processes leading to volcano eruptions, etc.

Challenge 626, page 354: See the puzzle about the motion of ski moguls.

Challenge 631, page 357: First, there are many more butterflies than tornadoes. Second, tornadoes do not rely on small initial disturbances for their appearance. Third, the belief in the butterfly 'effect' completely neglects an aspect of nature that is essential for self-organization: friction and dissipation. The butterfly 'effect', assumed that it existed, would require that dissipation in the air should have completely unrealistic properties. This is not the case in the atmosphere. But most important of all, there is no experimental basis for the 'effect': it has never been observed. Thus it does not exist.

Challenge 641, page 368: No. Nature does not allow more than about 20 digits of precision, as we will discover later in our walk. That is not sufficient for a standard book. The question whether such a number can be part of its own book thus disappears.

Challenge 642, page 368: All three statements are hogwash. A drag coefficient implies that the cross area of the car is known to the same precision. This is actually extremely difficult to measure and to keep constant. In fact, the value 0.375 for the Ford Escort was a cheat, as many other measurements showed. The fuel consumption is even more ridiculous, as it implies that fuel volumes and distances can be measured to that same precision. Opinion polls are taken by phoning at most 2000 people; due to the difficulties in selecting the right representative sample, that gives a precision of at most 3 %.

Challenge 644, page 369: Space-time is defined using matter; matter is defined using space-time

Challenge 645, page 369: Fact is that physics has been based on a circular definition for hundreds of years. Thus it is possible to build even an exact science on sand. Nevertheless, the elimination of the circularity is an important aim.

Challenge 646, page 370: Every measurement is a comparison with a standard; every comparison requires light or some other electromagnetic field. This is also the case for time measurements.

Challenge 647, page 370: Every mass measurement is a comparison with a standard; every comparison requires light or some other electromagnetic field.

Challenge 648, page 370: Angle measurements have the same properties as length or time measurements.

Challenge 653, page 387: About 10 µg.

Challenge 654, page 388: Probably the quantity with the biggest variation is mass, where a prefix for $1 \text{ eV}/c^2$ would be useful, as would be one for the total mass in the universe, which is about 10^{90} times larger.

Challenge 655, page 389: The formula with n-1 is a better fit. Why?

Challenge 658, page 390: No! They are much too precise to make sense. They are only given as an illustration for the behaviour of the Gaussian distribution. Real measurement distributions are not Gaussian to the precision implied in these numbers.

Challenge 659, page 390: About 0.3 m/s. It is *not* 0.33 m/s, it is *not* 0.333 m/s and it is *not* any longer strings of threes!

Challenge 661, page 395: The slowdown goes *quadratically* with time, because every new slowdown adds to the old one!

Challenge 662, page 395: No, only properties of parts of the universe are listed. The universe Vol. VI, page 103 itself has no properties, as shown in the last volume..

Challenge 663, page 448: For example, speed inside materials is slowed, but between atoms, light still travels with vacuum speed.



Aiunt enim multum legendum esse, non multa. Plinius, *Epistulae.**

- For a history of science in antiquity, see Lucio Russo, *La rivoluzione dimenticata*, Feltrinelli 1996, also available in several other languages. Cited on page 15.
- 2 If you want to catch up secondary school physics, the clearest and shortest introduction world-wide is a free school text, available in English and many other languages, written by a researcher who has dedicated all his life to the teaching of physics in secondary school, together with his university team: Friedrich Herrmann, *The Karlsruhe Physics Course*, free to download in English, Spanish, Russian and Italian at www.physikdidaktik. uni-karlsruhe.de. It is one of the few secondary school texts that captivates, surprises and challenges even professional physicists. This can be said even more of the wonderfully daring companion book Friedrich Herrmann & Georg Job, *Altlasten der Physik*, Aulis Verlag Deubner, 2002, whose content is available for free, in German, on the same site.

A beautiful book explaining physics and its many applications in nature and technology vividly and thoroughly is Paul G. Hewitt, John Suchocki & Leslie A. Hewitt, Conceptual Physical Science, Bejamin/Cummings, 1999.

A book famous for its passion for curiosity is RICHARD P. FEYNMAN, ROBERT B. LEIGHTON & MATTHEW SANDS, *The Feynman Lectures on Physics*, Addison Wesley, 1977.

A lot can be learned about motion from quiz books. One of the best is the well-structured collection of beautiful problems that require no mathematics, written by Jean-Marc Lévy-Leblond, *La physique en questions – mécanique*, Vuibert, 1998.

Another excellent quiz collection is YAKOV PERELMAN, *Oh, la physique*, Dunod, 2000, a translation from the Russian original.

A good problem book is W. G. Rees, *Physics by Example: 200 Problems and Solutions*, Cambridge University Press, 1994.

A good history of physical ideas is given in the excellent text by DAVID PARK, *The How and the Why*, Princeton University Press, 1988.

An excellent introduction into physics is ROBERT POHL, *Pohl's Einführung in die Physik*, Klaus Lüders & Robert O. Pohl editors, Springer, 2004, in two volumes with CDs. It is a new edition of a book that is over 70 years old; but the didactic quality, in particular of the experimental side of physics, is unsurpassed.

^{* &#}x27;Read much, but not anything.' Ep. 7, 9, 15. Gaius Plinius Secundus (b. 23/4 Novum Comum, d. 79 Vesuvius eruption), Roman writer, especially famous for his large, mainly scientific work *Historia naturalis*, which has been translated and read for almost 2000 years.

Another excellent Russian physics problem book, the so-called *Saraeva*, seems to exist only as Portuguese translation: B.B. BÚJOVTSEV, V.D. KRÍVCHENKOV, G.YA. MIÁK-ISHEV & I.M. SARÁEVA *Problemas seleccionados de física elemental*, Mir, 1979.

Another good physics problem book is GIOVANNI TONZIG, *Cento errori di fisica pronti per l'uso*, Sansoni, third edition, 2006. See also his www.giovannitonzig.it website. Cited on pages 15, 111, 189, and 275.

- An overview of motion illusions can be found on the excellent website www.michaelbach. de/ot. The complex motion illusion figure is found on www.michaelbach.de/ot/mot_rotsnake/index.html; it is a slight variation of the original by Kitaoka Akiyoshi at www.ritsumei.ac.jp/~akitaoka/rotsnake.gif, published as A. KITAOKA & H. ASHIDA, Phenomenal characteristics of the peripheral drift illusion, Vision 15, pp. 261–262, 2003. A famous scam is to pretend that the illusion is due to or depends on stress. Cited on page 16.
- These and other fantastic illusions are also found in AKIYOSHI KITAOKA, *Trick Eyes*, Barnes & Noble, 2005. Cited on page 16.
- A well-known principle in the social sciences states that, given a question, for every possible answer, however weird it may seem, there is somebody and often a whole group who holds it as his opinion. One just has to go through literature (or the internet) to confirm this.

About group behaviour in general, see R. Axelrod, *The Evolution of Cooperation*, Harper Collins, 1984. The propagation and acceptance of ideas, such as those of physics, are also an example of human cooperation, with all its potential dangers and weaknesses. Cited on page 16.

All the known texts by Parmenides and Heraclitus can be found in Jean-Paul Dumont, Les écoles présocratiques, Folio-Gallimard, 1988. Views about the non-existence of motion have also been put forward by much more modern and much more contemptible authors, such as in 1710 by Berkeley. Cited on page 17.

Vol. III, page 260

- An example of people worried by Zeno is given by WILLIAM MCLAUGHLIN, *Resolving Zeno's paradoxes*, Scientific American pp. 66–71, November 1994. The actual argument was not about a hand slapping a face, but about an arrow hitting the target. See also Ref. 61. Cited on page 17.
- 8 The full text of *La Beauté* and the other poems from *Les fleurs du mal*, one of the finest books of poetry ever written, can be found at the hypermedia.univ-paris8.fr/bibliotheque/Baudelaire/Spleen.html website. Cited on page 18.
- 9 The most famous text is Jearl Walker, *The Flying Circus of Physics*, Wiley, 1975. For more interesting physical effects in everyday life, see Erwein Flachsel, *Hundertfünfzig Physikrätsel*, Ernst Klett Verlag, 1985. The book also covers several clock puzzles, in puzzle numbers 126 to 128. Cited on page 19.
- A concise and informative introduction into the history of classical physics is given in the first chapter of the book by FLOYD KARKER RICHTMYER, EARLE HESSE KENNARD & JOHN N. COOPER, Introduction to Modern Physics, McGraw-Hill, 1969. Cited on page 19.
- An introduction into perception research is BRUCE GOLDSTEIN, *Perception*, Books/Cole, 5th edition, 1998. Cited on pages 21 and 25.
- A good overview over the arguments used to prove the existence of god from motion is given by MICHAEL BUCKLEY, *Motion and Motion's God*, Princeton University Press, 1971. The intensity of the battles waged around these failed attempts is one of the tragicomic chapters of history. Cited on page 21.

- **13** THOMAS AQUINAS, *Summa Theologiae* or *Summa Theologica*, 1265–1273, online in Latin at www.newadvent.org/summa, in English on several other servers. Cited on page 21.
- **14** For an exploration of 'inner' motions, see the beautiful text by RICHARD SCHWARTZ, *Internal Family Systems Therapy*, The Guilford Press, 1995. Cited on page 21.
- 15 For an authoritative description of proper motion development in babies and about how it leads to a healthy character see EMMI PIKLER, *Laßt mir Zeit Die selbstständige Bewegungsentwicklung des Kindes bis zum freien Gehen*, Pflaum Verlag, 2001, and her other books. See also the website www.pikler.org. Cited on page 21.
- **16** See e.g. the fascinating text by DAVID G. CHANDLER, *The Campaigns of Napoleon The Mind and Method of History's Greatest Soldier*, Macmillan, 1966. Cited on page 21.
- 17 RICHARD MARCUS, *American Roulette*, St Martin's Press, 2003, a thriller and a true story. Cited on page 21.
- A good and funny book on behaviour change is the well-known text by RICHARD BANDLER, Using Your Brain for a Change, Real People Press, 1985. See also RICHARD BANDLER & JOHN GRINDER, Frogs into princes Neuro Linguistic Programming, Eden Grove Editions, 1990. Cited on pages 21 and 31.
- A beautiful book about the mechanisms of human growth from the original cell to full size is Lewis Wolpert, *The Triumph of the Embryo*, Oxford University Press, 1991. Cited on page 21.
- On the topic of grace and poise, see e.g. the numerous books on the Alexander technique, such as M. Gelb, Body Learning An Introduction to the Alexander Technique, Aurum Press, 1981, and Richard Brennan, Introduction to the Alexander Technique, Little Brown and Company, 1996. Among others, the idea of the Alexander technique is to return to the situation that the muscle groups for sustainment and those for motion are used only for their respective function, and not vice versa. Any unnecessary muscle tension, such as neck stiffness, is a waste of energy due to the use of sustainment muscles for movement and of motion muscles for sustainment. The technique teaches the way to return to the natural use of muscles.

Motion of animals was discussed extensively already in the seventeenth century by G. Borelli, *De motu animalium*, 1680. An example of a more modern approach is J. J. Collins & I. Stewart, *Hexapodal gaits and coupled nonlinear oscillator models*, Biological Cybernetics 68, pp. 287–298, 1993. See also I. Stewart & M. Golubitsky, *Fearful Symmetry*, Blackwell, 1992. Cited on pages 22 and 112.

- 21 The results on the development of children mentioned here and in the following have been drawn mainly from the studies initiated by Jean Piaget; for more details on child development, see later on. At www.piaget.org you can find the website maintained by the Jean Piaget Society. Cited on pages 24, 39, and 41.
- The reptilian brain (eat? flee? ignore?), also called the R-complex, includes the brain stem, the cerebellum, the basal ganglia and the thalamus; the old mammalian (emotions) brain, also called the limbic system, contains the amygdala, the hypothalamus and the hippocampus; the human (and primate) (rational) brain, called the neocortex, consists of the famous grey matter. For images of the brain, see the atlas by John Nolte, *The Human Brain: An Introduction to its Functional Anatomy*, Mosby, fourth edition, 1999. Cited on page 25.
- The lower left corner film can be reproduced on a computer after typing the following lines in the Mathematica software package: Cited on page 25.

« Graphics'Animation' Nxpixels=72; Nypixels=54; Nframes=Nxpixels 4/3;

Vol. III, page 201

But our motion detection system is much more powerful than the example shown in the lower left corners. The following, different film makes the point.

Similar experiments, e.g. using randomly changing random patterns, show that the eye perceives motion even in cases where all Fourier components of the image are practically zero; such image motion is called *drift-balanced* or *non-Fourier* motion. Several examples are presented in J. Zanker, *Modelling human motion perception I: Classical stimuli*, Naturwissenschaften 81, pp. 156–163, 1994, and J. Zanker, *Modelling human motion perception II: Beyond Fourier motion stimuli*, Naturwissenschaften 81, pp. 200–209, 1994.

- 24 All fragments from Heraclitus are from John Mansley Robinson, *An Introduction to Early Greek Philosophy*, Houghton Muffin 1968, chapter 5. Cited on page 26.
- An introduction to Newton the alchemist are the two books by BETTY JO TEETER DOBBS, The Foundations of Newton's Alchemy, Cambridge University Press, 1983, and The Janus Face of Genius, Cambridge University Press, 1992. Newton is found to be a sort of highly intellectual magician, desperately looking for examples of processes where gods interact with the material world. An intense but tragic tale. A good overview is provided by R. G. KEESING, Essay Review: Newton's Alchemy, Contemporary Physics 36, pp. 117–119, 1995.

Newton's infantile theology, typical for god seekers who grew up without a father, can be found in the many books summarizing the letter exchanges between Clarke, his secretary, and Leibniz, Newton's rival for fame. Cited on page 33.

An introduction to the story of classical mechanics, which also destroys a few of the myths surrounding it – such as the idea that Newton could solve differential equations or that he introduced the expression F = ma – is given by CLIFFORD A. TRUESDELL, *Essays in the History of Mechanics*, Springer, 1968. Cited on pages 33, 165, and 196.

tain – The Adventure of Physics 🛮 pdf file available free of charge at www.motionmountain.net - Copyright © Christoph Schiller November 1997–March 2012

27 C. LIU, Z. DUTTON, C. H. BEHROOZI & L. VESTERGAARD HAU, Observation of coherent optical information storage in an atomic medium using halted light pulses, Nature 409, pp. 490–493, 2001. There is also a comment on the paper by E. A. CORNELL, Stopping light in its track, 409, pp. 461–462, 2001. However, despite the claim, the light pulses of course have not been halted. Can you give at least two reasons without even reading the paper, and maybe a third after reading it?

Challenge 663 s

- The work was an improvement on the previous experiment where a group velocity of light of 17 m/s had been achieved, in an ultracold gas of sodium atoms, at nanokelvin temperatures. This was reported by L. Vestergaard Hau, S. E. Harris, Z. Dutton & C. H. Behroozi, *Light speed reduction to 17 meters per second in an ultracold atomic gas*, Nature 397, pp. 594–598, 1999. Cited on page 35.
- **28** RAINER FLINDT, *Biologie in Zahlen Eine Datensammlung in Tabellen mit über 10.000 Einzelwerten*, Spektrum Akademischer Verlag, 2000. Cited on page 35.
- Two jets with that speed have been observed by I. F. MIRABEL & L. F. RODRÍGUEZ, *A superluminal source in the Galaxy*, Nature 371, pp. 46–48, 1994, as well as the comments on p. 18. Cited on page 35.
- A beautiful introduction to the slowest motions in nature, the changes in landscapes, is Detlev Busche, Jürgen Kempf & Ingrid Stengel, Landschaftsformen der Erde Bildatlas der Geomorphologie, Primus Verlag, 2005. Cited on page 36.
- 31 To build your own sundial, see the pretty and short ARNOLD ZENKERT, *Faszination Sonnenuhr*, VEB Verlag Technik, 1984. See also the excellent and complete introduction into this somewhat strange world at the www.sundials.co.uk website. Cited on page 41.
- An introduction to the sense of time as a result of clocks in the brain is found in R. B. IVRY & R. Spencer, *The neural representation of time*, Current Opinion in Neurobiology 14, pp. 225–232, 2004. The chemical clocks in our body are described in John D. Palmer, *The Living Clock*, Oxford University Press, 2002, or in A. Ahlgren & F. Halberg, *Cycles of Nature: An Introduction to Biological Rhythms*, National Science Teachers Association, 1990. See also the www.msi.umn.edu/~halberg/introd website. Cited on page 42.

Vol. III, page 219

- This has been shown among others by the work of Anna Wierzbicka that is discussed in more detail in one of the subsequent volumes. The passionate best seller by the Chomskian author Steven Pinker, *The Language Instinct How the Mind Creates Language*, Harper Perennial, 1994, also discusses issues related to this matter, refuting amongst others on page 63 the often repeated false statement that the *Hopi* language is an exception. Cited on page 42.
- For more information, see the excellent and freely downloadable books on biological clocks by Wolfgang Engelmann on the website www.uni-tuebingen.de/plantphys/bioclox. Cited on page 44.
- B. GÜNTHER & E. MORGADO, Allometric scaling of biological rhythms in mammals, Biological Research 38, pp. 207–212, 2005. Cited on page 44.
- Aristotle rejects the idea of the flow of time in chapter IV of his *Physics*. See the full text on the classics.mit.edu/Aristotle/physics.4.iv.html website. Cited on page 47.
- Perhaps the most informative of the books about the 'arrow of time' is Hans Dieter Zeh, The Physical Basis of the Direction of Time, Springer Verlag, 4th edition, 2001. It is still the best book on the topic. Most other texts exist – have a look on the internet – but lack clarity of ideas.

A typical conference proceeding is J. J. Halliwell, J. Pérez-Mercader & Wojciech H. Zurek, *Physical Origins of Time Asymmetry*, Cambridge University Press, 1994. Cited on page 48.

On the issue of absolute and relative motion there are many books about few issues. Examples are Julian Barbour, Absolute or Relative Motion? Vol. 1: A Study from the Machian Point of View of the Discovery and the Structure of Spacetime Theories, Cambridge University Press, 1989, Julian Barbour, Absolute or Relative Motion? Vol. 2: The Deep Structure of General Relativity, Oxford University Press, 2005, or John Earman, World Enough and Spacetime: Absolute vs Relational Theories of Spacetime, MIT Press, 1989. A speculative solution on the alternative between absolute and relative motion is presented in volume VI. Cited on page 53.

Vol. VI, page 59

- Goastlines and other fractals are beautifully presented in Heinz-Otto Peitgen, Hartmut Jürgens & Dietmar Saupe, Fractals for the Classroom, Springer Verlag, 1992, pp. 232–245. It is also available in several other languages. Cited on page 54.
- R. Dougherty & M. Foreman, Banach-Tarski decompositions using sets with the property of Baire, Journal of the American Mathematical Society 7, pp. 75–124, 1994. See also Alan L. T. Paterson, Amenability, American Mathematical Society, 1998, and Robert M. French, The Banach-Tarski theorem, The Mathematical Intelligencer 10, pp. 21–28, 1998. Finally, there are the books by Bernard R. Gelbaum & John M. H. Olmsted, counter-examples in Analysis, Holden-Day, 1964, and their Theorems and counter-examples in Mathematics, Springer, 1993. Cited on page 56.
- The beautiful but not easy text is STAN WAGON, *The Banach Tarski Paradox*, Cambridge University Press, 1993. Cited on pages 56 and 407.
- About the shapes of salt water bacteria, see the corresponding section in the interesting book by Bernard Dixon, *Power Unseen How Microbes Rule the World*, W.H. Freeman, 1994. The book has about 80 sections, in which as many microorganisms are vividly presented. Cited on page 57.
- OLAF MEDENBACH & HARRY WILK, Zauberwelt der Mineralien, Sigloch Edition, 1977. It combines beautiful photographs with an introduction into the science of crystals, minerals and stones. About the largest crystals, see P. C. RICKWOOD, The largest crystals, 66, pp. 885–908, 1981, also available on www.minsocam.org/MSA/collectors_corner/arc/large_crystals.htm. For an impressive example, the Naica cave in Mexico, see www.naica.com.mx/ingles/index.htm Cited on page 57.
- 44 See the websites www.weltbildfrage.de/3frame.htm and www.lhup.edu/~dsimanek/hollow/morrow.htm. Cited on page 57.
- The smallest distances are probed in particle accelerators; the distance can be determined from the energy of the particle beam. In 1996, the value of 10⁻¹⁹ m (for the upper limit of the size of quarks) was taken from the experiments described in F. ABE & al., *Measurement of dijet angular distributions by the collider detector at Fermilab*, Physical Review Letters 77, pp. 5336–5341, 1996. Cited on page 63.
- These puzzles are taken from the puzzle collection at www.mathematische-basteleien.de. Cited on page 67.
- 47 ALEXANDER K. DEWDNEY, The Planiverse Computer Contact with a Two-dimensional World, Poseidon Books/Simon & Schuster, 1984. See also Edwin A. Abbott, Flatland: A romance of many dimensions, 1884. Several other fiction authors had explored the option of a two-dimensional universe before, always answering, incorrectly, in the affirmative. Cited on page 69.
- **48** J. Bohr & K. Olsen, *The ancient art of laying rope*, preprint at arxiv.org/abs/1004.0814 Cited on page 69.

49 An overview and references can be found at www.pbrc.hawaii.edu/~petra/animal_olympians. html. Cited on page 70.

- P. PIERANSKI, S. PRZYBYL & A. STASIAK, *Tight open knots*, European Physical Journal E 6, pp. 123–128, 2001, preprint at arxiv.org/abs/physics/0103016. Cited on page 70.
- On the world of fireworks, see the frequently asked questions list of the usenet group rec.pyrotechnics, or search the web. A simple introduction is the article by J. A. Conkling, *Pyrotechnics*, Scientific American pp. 66–73, July 1990. Cited on page 72.
- There is a whole story behind the variations of g. It can be discovered in Chuji Tsuboi, Gravity, Allen & Unwin, 1979, or in Wolfgang Torge, Gravimetry, de Gruyter, 1989, or in Milan Burša & Karel Pěč, The Gravity Field and the Dynamics of the Earth, Springer, 1993. The variation of the height of the soil by up to 0.3 m due to the Moon is one of the interesting effects found by these investigations. Cited on pages 73 and 173.
- 53 STILLMAN DRAKE, *Galileo: A Very Short Introduction*, Oxford University Press, 2001. Cited on page 73.
- **54** Andrea Frova, *La fisica sotto il naso 44 pezzi facili*, Biblioteca Universale Rizzoli, Milano, 2001. Cited on page 74.
- On the other hands, other sciences enjoy studying usual paths in all detail. See, for example, Heini Hediger, editor, *Die Straßen der Tiere*, Vieweg & Sohn, 1967. Cited on page 74.
- **56** H. K. Eriksen, J. R. Kristiansen, Ø. Langangen & I. K. Wehus, *How fast could Usain Bolt have run? A dynamical study*, American Journal of Physics 77, pp. 224–228, 2009. Cited on page 75.
- This was discussed in the *Frankfurter Allgemeine Zeitung*, 2nd of August, 1997, at the time of the world athletics championship. The values are for the fastest part of the race of a $100 \,\mathrm{m}$ sprinter; the exact values cited were called the running speed world records in 1997, and were given as $12.048 \,\mathrm{m/s} = 43.372 \,\mathrm{km/h}$ by Ben Johnson for men, and $10.99 \,\mathrm{m/s} = 39.56 \,\mathrm{km/h}$ for women. Cited on page 75.
- Long jump data and literature can be found in three articles all entitled *Is a good long jumper a good high jumper?*, in the American Journal of Physics 69, pp. 104–105, 2001. In particular, world class long jumpers run at 9.35 ± 0.15 m/s, with vertical take-off speeds of 3.35 ± 0.15 m/s, giving take-off angles of about (only) 20°. A new technique for achieving higher take-off angles would allow the world long jump record to increase dramatically. Cited on page 75.
- The study of shooting faeces (i.e., shit) and its mechanisms is a part of modern biology. The reason that caterpillars do this was determined by M. Weiss, *Good housekeeping: why do shelter-dwelling caterpillars fling their frass?*, Ecology Letters 6, pp. 361–370, 2003, who also gives the present record of 1.5 m for the 24 mg pellets of *Epargyreus clarus*. The picture of the flying frass is from S. Caveney, H. McLean & D. Surry, *Faecal firing in a skipper caterpillar is pressure-driven*, The Journal of Experimental Biology 201, pp. 121–133, 1998. Cited on page 75.
- 60 H. C. Bennet-Clark, Scale effects in jumping animals, pp. 185–201, in T. J. Pedley, editor, Scale Effects in Animal Locomotion, Academic Press, 1977. Cited on page 76.
- The arguments of Zeno can be found in Aristotle, *Physics*, VI, 9. It can be found translated in almost any language. The classics.mit.edu/Aristotle/physics.6.vi.html website provides an online version in English. Cited on pages 80 and 445.
- 62 See, for exaple, K. V. KUMAR & W. T. NORFLEET, Issues of human acceleration tolerance

after long-duration space flights, NASA Technical Memorandum 104753, pp. 1–55, 1992, available at ntrs.nasa.gov. Cited on page 81.

63 Etymology can be a fascinating topic, e.g. when research discovers the origin of the German word 'Weib' ('woman', related to English 'wife'). It was discovered, via a few texts in Tocharian – an extinct Indo-European language from a region inside modern China – to mean originally 'shame'. It was used for the female genital region in an expression meaning 'place of shame'. With time, this expression became to mean 'woman' in general, while being shortened to the second term only. This connection was discovered by the German linguist Klaus T. Schmidt; it explains in particular why the word is not feminine but neutral, i.e., why it uses the article 'das' instead of 'die'. Julia Simon, private communication.

Etymology can also be simple and plain fun, for example when one discovers in the *Oxford English Dictionary* that 'testimony' and 'testicle' have the same origin; indeed in Latin the same word 'testis' was used for both concepts. Cited on pages 82 and 93.

- An overview of the latest developments is given by J. T. Armstrong, D. J. Hunter, K. J. Johnston & D. Mozurkewich, Stellar optical interferometry in the 1990s, Physics Today pp. 42–49, May 1995. More than 100 stellar diameters were known already in 1995. Several dedicated powerful instruments are being planned. Cited on page 84.
- A good biology textbook on growth is ARTHUR F. HOPPER & NATHAN H. HART, Foundations of Animal Development, Oxford University Press, 2006. Cited on page 85.
- This is discussed for example in C. L. Stong, *The amateur scientist how to supply electric power to something which is turning*, Scientific American pp. 120–125, December 1975. It also discusses how to make a still picture of something rotating simply by using a few prisms, the so-called *Dove prisms*. Other examples of attaching something to a rotating body are given by E. Rieflin, *Some mechanisms related to Dirac's strings*, American Journal of Physics 47, pp. 379–381, 1979. Cited on page 85.
- **67** James A. Young, *Tumbleweed*, Scientific American **264**, pp. 82–87, March 1991. The tumbleweed is in fact quite rare, except in Hollywood westerns, where all directors feel obliged to give it a special appearance. Cited on page 86.
- About N. decemspinosa, see R. L. Caldwell, A unique form of locomotion in a stomatopod backward somersaulting, Nature 282, pp. 71–73, 1979, and R. Full, K. Earls, M. Wong & R. Caldwell, Locomotion like a wheel?, Nature 365, p. 495, 1993. About rolling caterpillars, see J. Brackenbury, Caterpillar kinematics, Nature 330, p. 453, 1997, and J. Brackenbury, Fast locomotion in caterpillars, Journal of Insect Physiology 45, pp. 525–533, 1999. More images around legs can be found on rjf9.biol.berkeley.edu/twiki/bin/view/PolyPEDAL/LabPhotographs. Cited on page 87.
- This spider, of the genus *Cebrennus*, has been described by Ingo Rechenberg from Berlin. See the video at www.youtube.com/watch?v=Aayb_h31RyQ. Cited on page 87.
- The first experiments to prove the rotation of the flagella were by M. SILVERMAN & M. I. SIMON, Flagellar rotation and the mechanism of bacterial motility, Nature 249, pp. 73–74, 1974. For some pretty pictures of the molecules involved, see K. Namba, A biological molecular machine: bacterial flagellar motor and filament, Wear 168, pp. 189–193, 1993, or the website www.nanonet.go.jp/english/mailmag/2004/01la.html. The present record speed of rotation, 1700 rotations per second, is reported by Y. Magariyama, S. Sugiyama, K. Muramoto, Y. Maekawa, I. Kawagishi, Y. Imae & S. Kudo, Very fast flagellar rotation, Nature 371, p. 752, 1994.

More on bacteria can be learned from DAVID DUSENBERY, *Life at a Small Scale*, Scientific American Library, 1996. Cited on page 87.

71 M. P. Brenner, S. Hilgenfeldt & D. Lohse, Single bubble sonoluminescence, Reviews of Modern Physics 74, pp. 425–484, 2002. Cited on page 89.

- 72 K. R. Weninger, B. P. Barber & S. J. Putterman, Pulsed Mie scattering measurements of the collapse of a sonoluminescing bubble, Physical Review Letters 78, pp. 1799–1802, 1997. Cited on page 89.
- On shadows, see the agreeable popular text by ROBERTO CASATI, Alla scoperta dell'ombra Da Platone a Galileo la storia di un enigma che ha affascinato le grandi menti dell'umanità, Oscar Mondadori, 2000, and his websites located at www.shadowmill.com and roberto. casati.free.fr/casati/roberto.htm. Cited on page 90.
- 74 There is also the beautiful book by Penelope Farrant, *Colour in Nature*, Blandford, 1997. Cited on page 90.
- 75 The 'laws' of cartoon physics can easily be found using any search engine on the internet. Cited on page 90.
- For the curious, an overview of the illusions used in the cinema and in television, which lead to some of the strange behaviour of images mentioned above, is given in Bernard Wilkie, *The Technique of Special Effects in Television*, Focal Press, 1993, and his other books, or in the *Cinefex* magazine. On digital cinema techniques, see Peter C. Slansky, editor, *Digitaler film dgitales Kino*, UVK Verlag, 2004. Cited on page 91.
- **77** AETIUS, Opinions, I, XXIII, 3. See JEAN-PAUL DUMONT, *Les écoles présocratiques*, Folio Essais, Gallimard, p. 426, 1991. Cited on page 91.
- **78** GIUSEPPE FUMAGALLI, *Chi l'ha detto?*, Hoepli, 1983. It is from Pappus of Alexandria's opus *Synagoge*, book VIII, 19. Cited on pages 92 and 203.
- **79** See www.straightdope.com/classics/a5_262.html and the more dubious en.wikipedia.org/wiki/Guillotine. Cited on page 94.
- For the role and chemistry of adenosine triphosphate (ATP) in cells and in living beings, see any chemistry book, or search the internet. The uncovering of the mechanisms around ATP has led to Nobel Prizes in Chemistry in 1978 and in 1997. Cited on page 100.
- A picture of this unique clock can be found in the article by A. GARRETT, *Perpetual motion a delicious delirium*, Physics World pp. 23–26, December 1990. Cited on page 101.
- A Shell study estimated the world's total energy consumption in 2000 to be 500 EJ. The US Department of Energy estimated it to be around 416 EJ. We took the lower value here. A discussion and a breakdown into electricity usage (14 EJ) and other energy forms, with variations per country, can be found in S. Benka, *The energy challenge*, Physics Today 55, pp. 38–39, April 2002, and in E. J. Monitz & M. A. Kenderdine, *Meeting energy challenges: technology and policy*, Physics Today 55, pp. 40–46, April 2002. Cited on pages 104 and 105
- For an overview, see the paper by J. F. MULLIGAN & H. G. HERTZ, An unpublished lecture by Heinrich Hertz: 'On the energy balance of the Earth', American Journal of Physics 65, pp. 36–45, 1997. Cited on page 105.
- For a beautiful photograph of this feline feat, see the cover of the journal and the article of J. Darius, *A tale of a falling cat*, Nature 308, p. 109, 1984. Cited on page 111.
- NATTHI L. SHARMA, A new observation about rolling motion, European Journal of Physics 17, pp. 353–356, 1996. Cited on page 112.
- **86** C. Singh, *When physical intuition fails*, American Journal of Physics 70, pp. 1103–1109, 2002. Cited on page 112.

453

- 87 SERGE GRACOVETSKY, *The Spinal Engine*, Springer Verlag, 1990. It is now also knon that human gait is chaotic. This is explained by M. Perc, *The dynamics of human gait*, European Journal of Physics 26, pp. 525–534, 2005. Cited on page 112.
- **88** M. LLOBERA & T. J. SLUCKIN, *Zigzagging: theoretical insights on climbing strategies*, Journal of Theoretical Biology 249, pp. 206–217, 2007. Cited on page 113.
- This description of life and death is called the concept of *maximal metabolic scope*. Look up details in your favourite library. A different phrasing is the one by M. YA. AZBEL, *Universal biological scaling and mortality*, Proceedings of the National Academy of Sciences of the USA 91, pp. 12453–12457, 1994. He explains that every atom in an organism consumes, on average, 20 oxygen molecules per life-span. Cited on page 114.
- 90 Duncan MacDougall, Hypothesis concerning soul substance together with experimental evidence of the existence of such substance, American Medicine 2, pp. 240–243, April 1907, and Duncan MacDougall, Hypothesis concerning soul substance, American Medicine 2, pp. 395–397, July 1907. Reading the papers shows that the author has little practice in performing reliable weight and time measurements. Cited on page 114.
- A good roulette prediction story from the 1970s is told by Thomas A. Bass, *The Eudaemonic Pie* also published under the title *The Newtonian Casino*, Backinprint, 2000. An overview up to 1998 is given in the paper Edward O. Thorp, *The invention of the first wearable computer*, *Proceedings of the Second International Symposium on Wearable Computers* (ISWC 1998), 19-20 October 1998, Pittsburgh, Pennsylvania, USA (IEEE Computer Society), pp. 4–8, 1998, downloadable at csdl.computer.org/comp/proceedings/iswc/1998/9074/00/9074toc.htm. Cited on page 115.
- **92** This and many other physics surprises are described in the beautiful lecture script by Josef Zweck, *Physik im Alltag*, the notes of his lectures held in 1999/2000 at the Universität Regensburg. Cited on pages 116 and 121.
- **93** The equilibrium of ships, so important in car ferries, is an interesting part of shipbuilding; an introduction was already given by Leonhard Euler, *Scientia navalis*, 1749. Cited on page 117.
- 94 THOMAS HEATH, Aristarchus of Samos the Ancient Copernicus, Dover, 1981, reprinted from the original 1913 edition. Aristarchus' treaty is given in Greek and English. Aristarchus was the first proposer of the heliocentric system. Aristarchus had measured the length of the day (in fact, by determining the number of days per year) to the astonishing precision of less than one second. This excellent book also gives an overview of Greek astronomy before Aristarchus, explained in detail for each Greek thinker. Aristarchus' text is also reprinted in Aristarchus, On the sizes and the distances of the Sun and the Moon, c. 280 BCE in MICHAEL J. CROWE, Theories of the World From Antiquity to the Copernican Revolution, Dover, 1990, especially on pp. 27–29. Cited on page 123.
- See for example the videos on the Coriolis effect at techtv.mit.edu/videos/3722 and techtv. mit.edu/videos/3714, or search for videos on youtube.com. Cited on page 124.
- The influence of the Coriolis effect on icebergs was studied most thoroughly by the Swedish physicist turned oceanographer Walfrid Ekman (1874–1954); the topic was suggested by the great explorer Fridtjof Nansen, who also made the first observations. In his honour, one speaks of the Ekman layer, Ekman transport and Ekman spirals. Any text on oceanography or physical geography will give more details about them. Cited on page 127.
- An overview of the effects of the Coriolis acceleration $a = -2\omega \times v$ in the rotating frame is given by Edward A. Desloge, *Classical Mechanics*, Volume 1, John Wiley & Sons, 1982. Even the so-called *Gulf Stream*, the current of warm water flowing from the Caribbean to the North Sea, is influenced by it. Cited on page 127.

The original publication is by A. H. Shapiro, *Bath-tub vortex*, Nature 196, pp. 1080–1081, 1962. He also produced two films of the experiment. The experiment has been repeated many times in the northern and in the southern hemisphere, where the water drains clockwise; the first southern hemisphere test was L.M. Trefethen & al., *The bath-tub vortex in the southern hemisphere*, Nature 201, pp. 1084–1085, 1965. A complete literature list is found in the letters to the editor of the American Journal of Physics 62, p. 1063, 1994. Cited on page 127.

- The tricks are explained by RICHARD CRANE, Short Foucault pendulum: a way to eliminate the precession due to ellipticity, American Journal of Physics 49, pp. 1004–1006, 1981, and particularly in RICHARD CRANE, Foucault pendulum wall clock, American Journal of Physics 63, pp. 33–39, 1993. The Foucault pendulum was also the topic of the thesis of Heike Kamerling Onnes, Nieuwe bewijzen der aswenteling der aarde, Universiteit Groningen, 1879. Cited on page 127.
- **100** The reference is J. G. Hagen, *La rotation de la terre : ses preuves mécaniques anciennes et nouvelles*, Sp. Astr. Vaticana Second. App. Rome, 1910. His other experiment is published as J. G. Hagen, *How Atwood's machine shows the rotation of the Earth even quantitatively*, International Congress of Mathematics, Aug. 1912. Cited on page 128.
- **101** The original papers are A. H. Compton, *A laboratory method of demonstrating the Earth's rotation*, Science 37, pp. 803–806, 1913, A. H. Compton, *Watching the Earth revolve*, Scientific American Supplement no. 2047, pp. 196–197, 1915, and A. H. Compton, *A determination of latitude, azimuth and the length of the day independent of astronomical observations*, Physical Review (second series) 5, pp. 109–117, 1915. Cited on page 130.
- The G-ring in Wettzell is so precise, with a resolution of less than 10⁻⁸, that it has detected the motion of the poles. For details, see K. U. Schreiber, A. Velikoseltsev, M. Rothacher, T. Kluegel, G. E. Stedman & D. L. Wiltshire, Direct measurement of diurnal polar motion by ring laser gyroscopes, Journal of Geophysical Research 109 B, p. 06405, 2004, an a review article at T. Klügel, W. Schlüter, U. Schreiber & M. Schneider, Großringlaser zur kontinuierlichen Beobachtung der Erdrotation, Zeitschrift für Vermessungswesen 130, pp. 99–108, February 2005. Cited on page 130.
- **103** R. Anderson, H. R. Bilger & G. E. Stedman, *The Sagnac-effect: a century of Earth-rotated interferometers*, American Journal of Physics **62**, pp. 975–985, 1994.
 - See also the clear and extensive paper by G. E. STEDMAN, Ring laser tests of fundamental physics and geophysics, Reports on Progress in Physics 60, pp. 615–688, 1997. Cited on page 130.
- 104 About the length of the day, see the maia.usno.navy.mil website, or the books by K. Lambeck, *The Earth's Variable Rotation: Geophysical Causes and Consequences*, Cambridge University Press, 1980, and by W. H. Munk & G. J. F. Macdonald, *The Rotation of the Earth*, Cambridge University Press, 1960. About a modern ring laser set-up, see www. wettzell.ifag.de. Cited on pages 131 and 175.
- 105 H. Bucka, Zwei einfache Vorlesungsversuche zum Nachweis der Erddrehung, Zeitschrift für Physik 126, pp. 98–105, 1949, and H. Bucka, Zwei einfache Vorlesungsversuche zum Nachweis der Erddrehung. II. Teil, Zeitschrift für Physik 128, pp. 104–107, 1950. Cited on page 131.
- T. M. Demko, Late proterozoic and paleozoic tides, retreat of the moon, and rotation of the Earth, Science 273, pp. 100–104, 5 July 1996. They deduce from tidal sediment analysis that days were only 18 to 19 hours long in the Proterozoic, i.e., 900 million years ago; they assume that the year was 31 million seconds long from then to today. See also C. P. Sonett

- & M. A. Chan, Neoproterozoic Earth-Moon dynamics rework of the 900 MA Big Cotton-wood canyon tidal laminae, Geophysical Research Letters 25, pp. 539–542, 1998. Another determination was by G. E. Williams, Precambrian tidal and glacial clastic deposits: implications for precambrian Earth-Moon dynamics and palaeoclimate, Sedimentary Geology 120, pp. 55–74, 1998. Using a geological formation called *tidal rhythmites*, he deduced that about 600 million years ago there were 13 months per year and a day had 22 hours. Cited on page 131.
- **107** The story of this combination of history and astronomy is told in RICHARD STEPHENSON, Historical Eclispes and Earth's Rotation, Cambridge University Press, 1996. Cited on page 132.
- **108** B. F. Chao, *Earth Rotational Variations excited by geophysical fluids*, IVS 2004 General Meeting proceedings/ pages 38-46. Cited on page 132.
- **109** On the rotation and history of the solar system, see S. Brush, *Theories of the origin of the solar system 1956–1985*, Reviews of Modern Physics **62**, pp. 43–112, 1990. Cited on page 132.
- 110 The website hpiers.obspm.fr/eop-pc shows the motion of the Earth's axis over the last ten years. The International Latitude Service founded by Küstner is now part of the International Earth Rotation Service; more information can be found on the www.iers.org website. The latest idea is that two-thirds of the circular component of the polar motion, which in the USA is called 'Chandler wobble' after the person who attributed to himself the discovery by Küstner, is due to fluctuations of the ocean pressure at the bottom of the oceans and one-third is due to pressure changes in the atmosphere of the Earth. This is explained by R. S. Gross, *The excitation of the Chandler wobble*, Geophysical Physics Letters 27, pp. 2329–2332, 2000. Cited on page 133.
- **111** S. B. LAMBERT, C. BIZOUARD & V. DEHANT, Rapid variations in polar motion during the 2005-2006 winter season, Geophysical Research Letters 33, p. L13303, 2006. Cited on page 133.
- For more information about Alfred Wegener, see the (simple) text by Klaus Rohrbach, Alfred Wegener Erforscher der wandernden Kontinente, Verlag Freies Geistesleben, 1993; about plate tectonics, see the www.scotese.com website. About earthquakes, see the www.geo.ed.ac.uk/quakexe/quakes and the www.iris.edu/seismon website. See the vulcan.wr. usgs.gov and the www.dartmouth.edu/~volcano websites for information about volcanoes. Cited on page 135.
- J. Jouzel & al., Orbital and millennial Antarctic climate variability over the past 800,000 years, Science 317, pp. 793–796, 2007, takes the data from isotope concentrations in ice cores. In contrast, J. D. Hays, J. Imbrie & N. J. Shackleton, Variations in the Earth's orbit: pacemaker of the ice ages, Science 194, pp. 1121–1132, 1976, confirmed the connection with orbital parameters by literally digging in the mud that covers the ocean floor in certain places. Note that the web is full of information on the ice ages. Just look up 'Milankovitch' in a search engine. Cited on pages 138 and 139.
- **114** R. Humphreys & J. Larsen, *The sun's distance above the galactic plane*, Astronomical Journal 110, pp. 2183–2188, November 1995. Cited on page 140.
- **115** C. L. Bennet, M. S. Turner & M. White, *The cosmic rosetta stone*, Physics Today **50**, pp. 32–38, November 1997. Cited on page 140.
- 116 On http://www.geoffreylandis.com/vacuum.html you can read a description of what happened. See also the http://www.geoffreylandis.com/ebullism.html and imagine.gsfc.nasa. gov/docs/ask_astro/answers/970603.html websites. They all give details on the effects of vacuum on humans. Cited on page 144.

- **117** R. McN. Alexander, Leg design and jumping technique for humans, other vertebrates and insects, Philosophical Transactions of the Royal Society in London B 347, pp. 235–249, 1995. Cited on page 148.
- **118** J. W. Glasheen & T. A. McMahon, *A hydrodynamic model of locomotion in the basilisk lizard*, Nature **380**, pp. 340–342, For pictures, see also New Scientist, p. 18, 30 March 1996, or Scientific American, pp. 48–49, September 1997, or the website by the author at rjf2. biol.berkeley.edu/Full_Lab/FL_Personnel/J_Glasheen/J_Glasheen.html.
 - Several shore birds also have the ability to run over water, using the same mechanism. Cited on page 148.
- **119** A. Fernandez-Nieves & F. J. de las Nieves, *About the propulsion system of a kayak and of Basiliscus basiliscus*, European Journal of Physics 19, pp. 425–429, 1998. Cited on page 149.
- Y. S. Song, S. H. Suhr & M. Sitti, Modeling of the supporting legs for designing biomimetic water strider robot, Proceedings of the IEEE International Conference on Robotics and Automation, Orlando, USA, 2006. S. H. Suhr, Y. S. Song, S. J. Lee & M. Sitti, Biologically inspired miniature water strider robot, Proceedings of the Robotics: Science and Systems I, Boston, USA, 2005. See also the website www.me.cmu.edu/facultyl/sitti/nano/projects/waterstrider. Cited on page 149.
- **121** M. WITTLINGER, R. WEHNER & H. WOLF, *The ant odometer: stepping on stilts and stumps*, Science 312, pp. 1965–1967, 2006. Cited on page 150.
- **122** P. G. WEYAND, D. B. STERNLIGHT, M. J. BELLIZZI & S. WRIGHT, Faster top running speeds are achieved with greater ground forces not more rapid leg movements, Journal of Applied Physiology **89**, pp. 1991–1999, 2000. Cited on page 150.
- The material on the shadow discussion is from the book by ROBERT M. PRYCE, Cook and Peary, Stackpole Books, 1997. See also the details of Peary's forgeries in Wally Herbert, The Nose of Laurels, Doubleday 1989. The sad story of Robert Peary is also told in the centenary number of National Geographic, September 1988. Since the National Geographic Society had financed Peary in his attempt and had supported him until the US Congress had declared him the first man at the Pole, the (partial) retraction is noteworthy. (The magazine then changed its mind again later on, to sell more copies, and now again claims that Peary reached the North Pole.) By the way, the photographs of Cook, who claimed to have been at the North Pole even before Peary, have the same problem with the shadow length. Both men have a history of cheating about their 'exploits'. As a result, the first man at the North Pole was probably Roald Amundsen, who arrived there a few years later, and who was also the first man at the South Pole. Cited on page 151.
- **124** The story is told in M. NAUENBERG, *Hooke, orbital motion, and Newton's Principia*, American Journal of Physics 62, 1994, pp. 331–350. Cited on page 153.
- More details are given by D. RAWLINS, in *Doubling your sunsets or how anyone can measure the Earth's size with wristwatch and meter stick*, American Journal of Physics 47, 1979, pp. 126–128. Another simple measurement of the Earth radius, using only a sextant, is given by R. O'KEEFE & B. GHAVIMI-ALAGHA, in *The World Trade Center and the distance to the world's center*, American Journal of Physics 60, pp. 183–185, 1992. Cited on page 154.
- 126 More details on astronomical distance measurements can be found in the beautiful little book by A. VAN HELDEN, *Measuring the Universe*, University of Chicago Press, 1985, and in NIGEL HENBEST & HEATHER COOPER, *The Guide to the Galaxy*, Cambridge University Press, 1994. Cited on page 154.
- 127 A lot of details can be found in M. Jammer, Concepts of Mass in Classical and Modern Physics, reprinted by Dover, 1997, and in Concepts of Force, a Study in the Foundations of

Mechanics, Harvard University Press, 1957. These eclectic and thoroughly researched texts provide numerous details and explain various philosophical viewpoints, but lack clear statements and conclusions on the accurate description of nature; thus are not of help on fundamental issues.

Jean Buridan (*c.* 1295 to *c.* 1366) criticizes the distinction of sublunar and translunar motion in his book *De Caelo*, one of his numerous works. Cited on page 154.

- **128** D. TOPPER & D. E. VINCENT, *An analysis of Newton's projectile diagram*, European Journal of Physics 20, pp. 59–66, 1999. Cited on page 155.
- **129** The absurd story of the metre is told in the historical novel by Ken Alder, *The Measure of All Things: The Seven-Year Odyssey and Hidden Error that Transformed the World*, The Free Press, 2003. Cited on page 157.
- **130** H. CAVENDISH, *Experiments to determine the density of the Earth*, Philosophical Transactions of the Royal Society 88, pp. 469–526, 1798. In fact, the first value of the gravitational constant *G* found in the literature is only from 1873, by Marie-Alfred Cornu and Jean-Baptistin Baille, who used an improved version of Cavendish's method. Cited on page 159.
- About the measurement of spatial dimensions via gravity and the failure to find any hint for a number different from three see the review by E. G. ADELBERGER, B. R. HECKEL & A. E. Nelson, *Tests of the gravitational inverse-square law*, Annual Review of Nuclear and Particle Science 53, pp. 77–121, 2003, also arxiv.org/abs/hep-ph/0307284, or the review by J. A. Hewett & M. Spiropulu, *Particle physics probes of extra spacetime dimensions*, Annual Review of Nuclear and Particle Science 52, pp. 397–424, 2002, arxiv.org/abs/hep-ph/0205106. Cited on page 162.
- 132 There are many books explaining the origin of the precise shape of the Earth, such as the pocket book S. Anders, *Weil die Erde rotiert*, Verlag Harri Deutsch, 1985. Cited on page 162.
- **133** The shape of the Earth is described most precisely with the World Geodetic System. For an extensive presentation, see the www.eurocontrol.be/projects/eatchip/wgs84/start.html website. See also the website of the *International Earth Rotation Service* at hpiers.obspm.fr. Cited on page 162.
- **134** G. HECKMAN & M. VAN HAANDEL, *De vele beweijzen van Kepler's wet over ellipsenbanen: een nieuwe voor 'het Boek'?*, Nederlands tijdschrift voor natuurkunde 73, pp. 366–368, November 2007. Cited on page 165.
- **135** W. K. HARTMAN, R. J. PHILLIPS & G. J. TAYLOR, editors, *Origin of the Moon*, Lunar and Planetary Institute, 1986. Cited on page 168.
- 136 If you want to read about the motion of the Moon in all its fascinating details, have a look at Martin C. Gutzwiller, *Moon–Earth–Sun: the oldest three body problem*, Reviews of Modern Physics 70, pp. 589–639, 1998. Cited on page 168.
- **137** DIETRICH NEUMANN, *Physiologische Uhren von Insekten Zur Ökophysiologie lunarperiodisch kontrollierter Fortpflanzungszeiten*, Naturwissenschaften **82**, pp. 310–320, 1995. Cited on page 168.
- 138 The origin of the duration of the menstrual cycle is not yet settled; however, there are explanations on how it becomes synchronized with other cycles. For a general explanation see ARKADY PIKOVSKY, MICHAEL ROSENBLUM & JÜRGEN KURTHS, Synchronization: A Universal Concept in Nonlinear Science, Cambridge University Press, 2002. Cited on page 168.

- **139** J. LASKAR, F. JOUTEL & P. ROBUTEL, *Stability of the Earth's obliquity by the moon*, Nature **361**, pp. 615–617, 1993. However, the question is not completely settled, and other opinions exist. Cited on page 168.
- **140** NEIL F. Comins, What if the Moon Did not Exist? Voyages to Earths that Might Have Been, Harper Collins, 1993. Cited on page 168.
- PAUL A. WIEGERT, KIMMO A. INNANEN & SEPPO MIKKOLA, An asteroidal companion to the Earth, Nature 387, pp. 685–686, 12 June 1997, together with the comment on pp. 651–652. Details on the orbit and on the fact that Lagrangian points do not always form equilateral triangles can be found in F. NAMOUNI, A. A. CHRISTOU & C. D. MURRAY, Coorbital dynamics at large eccentricity and inclination, Physical Review Letters 83, pp. 2506–2509, 1999. Cited on page 171.
- **142** SIMON NEWCOMB, Astronomical Papers of the American Ephemeris 1, p. 472, 1882. Cited on page 172.
- 143 For an animation of the tides, have a look at www.jason.oceanobs.com/html/applications/marees/m2_atlantique_fr.html. Cited on page 172.
- **144** A beautiful introduction is the classic G. FALK & W. RUPPEL, *Mechanik, Relativität, Gravitation ein Lehrbuch*, Springer Verlag, Dritte Auflage, 1983. Cited on page 173.
- **145** J. SOLDNER, Berliner Astronomisches Jahrbuch auf das Jahr 1804, 1801, p. 161. Cited on page 177.
- The equality was first tested with precision by R. VON EÖTVÖS, Annalen der Physik & Chemie 59, p. 354, 1896, and by R. VON EÖTVÖS, V. PEKÁR, E. FEKETE, Beiträge zum Gesetz der Proportionalität von Trägheit und Gravität, Annalen der Physik 4, Leipzig 68, pp. 11–66, 1922. He found agreement to 5 parts in 10⁹. More experiments were performed by P. G. ROLL, R. KROTKOW & R. H. DICKE, The equivalence of inertial and passive gravitational mass, Annals of Physics (NY) 26, pp. 442–517, 1964, one of the most interesting and entertaining research articles in experimental physics, and by V. B. BRAGINSKY & V. I. PANOV, Soviet Physics JETP 34, pp. 463–466, 1971. Modern results, with errors less than one part in 10¹², are by Y. Su & al., New tests of the universality of free fall, Physical Review D50, pp. 3614–3636, 1994. Several experiments have been proposed to test the equality in space to less than one part in 10¹⁶. Cited on page 178.
- 147 H. EDELMANN, R. NAPIWOTZKI, U. HEBER, N. CHRISTLIEB & D. REIMERS, HE 0437-5439: an unbound hyper-velocity B-type star, The Astrophysical Journal 634, pp. L181–L184, 2005. Cited on page 182.
- **148** This is explained for example by D.K. FIRPIĆ & I.V. ANIÇIN, *The planets, after all, may run only in perfect circles but in the velocity space!*, European Journal of Physics 14, pp. 255–258, 1993. Cited on pages 182 and 430.
- **149** See L. Hodges, *Gravitational field strength inside the Earth*, American Journal of Physics 59, pp. 954–956, 1991. Cited on page 183.
- **150** P. MOHAZZABI & M. C. JAMES, *Plumb line and the shape of the Earth*, American Journal of Physics **68**, pp. 1038–1041, 2000. Cited on page 185.
- **151** From Neil de Gasse Tyson, *The Universe Down to Earth*, Columbia University Press, 1994. Cited on page 186.
- **152** G. D. QUINLAN, *Planet X: a myth exposed*, Nature 363, pp. 18–19, 1993. Cited on page 186.
- 153 See en.wikipedia.org/wiki/90377_Sedna. Cited on page 186.
- **154** See R. Matthews, *Not a snowball's chance ...*, New Scientist 12 July 1997, pp. 24–27. The original claim is by Louis A. Frank, J. B. Sigwarth & J. D. Craven, *On the influx of*

- *small comets into the Earth's upper atmosphere*, parts I and II, Geophysical Research Letters 13, pp. 303–306, pp. 307–310, 1986. The latest observations have disproved the claim. Cited on page 186.
- 155 The ray form is beautifully explained by J. Evans, *The ray form of Newton's law of motion*, American Journal of Physics **61**, pp. 347–350, 1993. Cited on page 188.
- This is a small example from the beautiful text by MARK P. SILVERMAN, And Yet It Moves: Strange Systems and Subtle Questions in Physics, Cambridge University Press, 1993. It is a treasure chest for anybody interested in the details of physics. Cited on page 189.
- 157 G.-L. Lesage, *Lucrèce Newtonien*, Nouveaux mémoires de l'Académie Royale des Sciences et Belles Lettres pp. 404–431, 1747, or www3.bbaw.de/bibliothek/digital/struktur/03-nouv/1782/jpg-0600/00000495.htm. See also en.wikipedia.org/wiki/Le_Sage's_theory_of_gravitation. In fact, the first to propose the idea of gravitation as a result of small particles pushing masses around was Nicolas Fatio de Duillier in 1688. Cited on page 189.
- J. LASKAR, A numerical experiment on the chaotic behaviour of the solar system, Nature 338, pp. 237–238, 1989, and J. LASKAR, The chaotic motion of the solar system A numerical estimate of the size of the chaotic zones, Icarus 88, pp. 266–291, 1990. The work by Laskar was later expanded by Jack Wisdom, using specially built computers, following only the planets, without taking into account the smaller objects. For more details, see G. J. Sussman & J. Wisdom, Chaotic Evolution of the Solar System, Science 257, pp. 56–62, 1992. Today, such calculations can be performed on your home PC with computer code freely available on the internet. Cited on page 190.
- B. Dubrulle & F. Graner, Titius-Bode laws in the solar system. 1: Scale invariance explains everything, Astronomy and Astrophysics 282, pp. 262–268, 1994, and Titius-Bode laws in the solar system. 2: Build your own law from disk models, Astronomy and Astrophysics 282, pp. 269–276, 1994. Cited on page 191.
- **160** M. LECAR, *Bode's Law*, Nature **242**, pp. 318–319, 1973, and M. HENON, *A comment on "The resonant structure of the solar system" by A.M. Molchanov*, Icarus **11**, pp. 93–94, 1969. Cited on page 191.
- **161** Cassius Dio, *Historia Romana*, c. 220, book 37, 18. For an English translation, see the site penelope.uchicago.edu/Thayer/E/Roman/Texts/Cassius_Dio/37*.html. Cited on page 191.
- **162** A. J. Simoson, *Falling down a hole through the Earth*, Mathematics Magazine 77, pp. 171–188, 2004. See also A. J. Simoson, *The gravity of Hades*, 75, pp. 335–350, 2002. Simoson speculates whether the spirographics swirls in the are due to such an effect. Cited on page 192.
- M. Bevis, D. Alsdorf, E. Kendrick, L. P. Fortes, B. Forsberg, R. Malley & J. Becker, Seasonal fluctuations in the mass of the Amazon River system and Earth's elastic response, Geophysical Research Letters 32, p. L16308, 2005. Cited on page 192.
- D. HESTENES, M. WELLS & G. SWACKHAMER, Force concept inventory, Physics Teacher 30, pp. 141–158, 1982. The authors developed tests to check the understanding of the concept of physical force in students; the work has attracted a lot of attention in the field of physics teaching. Cited on page 197.
- **165** For a general overview on friction, from physics to economics, architecture and organizational theory, see N. ÅKERMAN, editor, *The Necessity of Friction Nineteen Essays on a Vital Force*, Springer Verlag, 1993. Cited on page 200.
- **166** See M. HIRANO, K. SHINJO, R. KANECKO & Y. MURATA, Observation of superlubricity by scanning tunneling microscopy, Physical Review Letters 78, pp. 1448–1451, 1997. See

- also the discussion of their results by Serge Fayeulle, *Superlubricity: when friction stops*, Physics World pp. 29–30, May 1997. Cited on page 200.
- **167** Donald Ahrens, *Meteorology Today: An Introduction to the Weather, Climate, and the Environment*, West Publishing Company, 1991. Cited on page 201.
- 168 This topic is discussed with lucidity by J. R. MUREIKA, What really are the best 100 m performances?, Athletics: Canada's National Track and Field Running Magazine, July 1997. It can also be found as arxiv.org/abs/physics/9705004, together with other papers on similar topics by the same author. Cited on page 201.
- **169** F. P. BOWDEN & D. TABOR, *The Friction and Lubrication of Solids*, Oxford University Press, Part I, revised edition, 1954, and part II, 1964. Cited on page 202.
- **170** A powerful book on human violence is James Gilligan, *Violence Our Deadly Epidemic and its Causes*, Grosset/Putnam, 1992. Cited on page 202.
- 171 The main tests of randomness of number series among them the gorilla test can be found in the authoritative paper by G. MARSAGLIA & W. W. TSANG, Some difficult-to-pass tests of randomness, Journal of Statistical Software 7, p. 8, 2002. It can also be downloaded from the www.jstatsoft.org/v07/i03/tuftests.pdf website. Cited on page 205.
- **172** For one aspect of the issue, see for example the captivating book by Bert Hellinger, *Zweierlei Glück*, Carl Auer Systeme Verlag, 1997. The author explains how to live serenely and with the highest possible responsibility for one's actions, by reducing entanglements with the destiny of others. He describes a powerful technique to realise this goal.
 - A completely different viewpoint is given by Aung San Suu Kyi, *Freedom from Fear*, Penguin, 1991. One of the bravest women on Earth, she won the Nobel Peace Price in 1991. Cited on page 207.
- **173** HENRIK WALTER, *Neurophilosophie der Willensfreiheit*, Mentis Verlag, Paderborn 1999. Also available in English translation. Cited on page 207.
- 174 GIUSEPPE FUMAGALLI, Chi l'ha detto?, Hoepli, 1983. Cited on page 208.
- 175 See the tutorial on the Peaucellier-Lipkin linkage by D.W. Henderson and D. Taimina found on kmoddl.library.cornell.edu/tutorials/11/index.php. The internet contains many other pages on the topic. Cited on page 209.
- **176** The beautiful story of the south-pointing carriage is told in Appendix B of James Foster & J. D. Nightingale, *A Short Course in General Relativity*, Springer Verlag, 2nd edition, 1998. Such carriages have existed in China, as told by the great sinologist Joseph Needham, but their construction is unknown. The carriage described by Foster and Nightingale is the one reconstructed in 1947 by George Lancaster, a British engineer. Cited on page 209.
- **177** See for example Z. Ghahramani, *Building blocks of movement*, Nature 407, pp. 682–683, 2000. Researchers in robot control are also interested in such topics. Cited on page 210.
- **178** G. GUTIERREZ, C. FEHR, A. CALZADILLA & D. FIGUEROA, Fluid flow up the wall of a spinning egg, American Journal of Physics 66, pp. 442–445, 1998. Cited on page 210.
- **179** A historical account is given in Wolfgang Yourgray & Stanley Mandelstam, *Variational Principles in Dynamics and Quantum Theory*, Dover, 1968. Cited on pages 213 and 221.
- **180** C. G. Gray & E. F. Taylor, *When action is not least*, American Journal of Physics 75, pp. 434–458, 2007. Cited on page 217.
- **181** MAX PÄSLER, *Prinzipe der Mechanik*, Walter de Gruyter & Co., 1968. Cited on page 219.
- **182** The relations between possible Lagrangians are explained by Herbert Goldstein, *Classical Mechanics*, 2nd edition, Addison-Wesley, 1980. Cited on page 219.

183 The Hemingway statement is quoted by Marlene Dietrich in AARON E. HOTCHNER, *Papa Hemingway*, Random House, 1966, in part 1, chapter 1. Cited on page 220.

- **184** C. G. Gray, G. Karl & V. A. Novikov, From Maupertius to Schrödinger. Quantization of classical variational principles, American Journal of Physics 67, pp. 959–961, 1999. Cited on page 221.
- **185** J. A. Moore, *An innovation in physics instruction for nonscience majors*, American Journal of Physics 46, pp. 607–612, 1978. Cited on page 222.
- 186 See e.g. Alan P. Boss, *Extrasolar planets*, Physics Today 49, pp. 32–38. September 1996. The most recent information can be found at the 'Extrasolar Planet Encyclopaedia' maintained at www.obspm.fr/planets by Jean Schneider at the Observatoire de Paris. Cited on page 225.
- **187** A good review article is by DAVID W. HUGHES, *Comets and Asteroids*, Contemporary Physics 35, pp. 75–93, 1994. Cited on page 225.
- **188** G. B. West, J. H. Brown & B. J. Enquist, A general model for the origin of allometric scaling laws in biology, Science 276, pp. 122–126, 4 April 1997, with a comment on page 34 of the same issue. The rules governing branching properties of blood vessels, of lymph systems and of vessel systems in plants are explained. For more about plants, see also the paper G. B. West, J. H. Brown & B. J. Enquist, A general model for the structure and allometry of plant vascular systems, Nature 400, pp. 664–667, 1999. Cited on page 226.
- **189** J. R. BANAVAR, A. MARTIN & A. RINALDO, Size and form in efficient transportation networks, Nature 399, pp. 130–132, 1999. Cited on page 226.
- 190 N. MOREIRA, New striders new humanoids with efficient gaits change the robotics land-scape, Science News Online 6th of August, 2005. Cited on page 227.
- 191 WERNER HEISENBERG, Der Teil und das Ganze, Piper, 1969. Cited on page 229.
- **192** See the clear presenttion by E. H. LOCKWOOD & R. H. MACMILLAN, *Geometric Symmetry*, Cambridge University Press, 1978. Cited on page 229.
- **193** JOHN MANSLEY ROBINSON, *An Introduction to Early Greek Philosophy*, Houghton Muffin 1968, chapter 5. Cited on page 230.
- **194** See e.g. B. Bower, *A child's theory of mind*, Science News 144, pp. 40–41. Cited on page 234.
- 195 The most beautiful book on this topic is the text by Branko Grünbaum & G. C. Shephard, *Tilings and Patterns*, W.H. Freeman and Company, New York, 1987. It has been translated into several languages and republished several times. Cited on page 235.
- 196 U. NIEDERER, The maximal kinematical invariance group of the free Schrödinger equation, Helvetica Physica Acta 45, pp. 802–810, 1972. See also the introduction by O. Jahn & V. V. Sreedhar, The maximal invariance group of Newton's equations for a free point particle, arxiv.org/abs/math-ph/0102011. Cited on page 241.
- 197 The story is told in the interesting biography of Einstein by A. PAIS, 'Subtle is the Lord...'
 The Science and the Life of Albert Einstein, Oxford University Press, 1982. Cited on page 242.
- **198** W. ZÜRN & R. WIDMER-SCHNIDRIG, *Globale Eigenschwingungen der Erde*, Physik Journal 1, pp. 49–55, 2002. Cited on page 250.
- **199** N. GAUTHIER, What happens to energy and momentum when two oppositely-moving wave pulses overlap?, American Journal of Physics 71, pp. 787–790, 2003. Cited on page 258.

- **200** An informative and modern summary of present research about the ear and the details of its function is www.physicsweb.org/article/world/15/5/8. Cited on page 261.
- **201** S. Adachi, *Principles of sound production in wind instruments*, Acoustical Science and Technology 25, pp. 400–404, 2004. Cited on page 263.
- 202 The literature on tones and their effects is vast. For example, people have explored the differences and effects of various intonations in great detail. Several websites, such as bellsouthpwp.net/j/d/jdelaub/jstudio.htm, allow listening to music played with different intonations. People have even researched whether animals use just or chromatic intonation. (See, for example, K. Leutwyler, *Exploring the musical brain*, Scientific American January 2001.) There are also studies of the effects of low frequencies, of beat notes, and of many other effects on humans. However, many studies mix serious and non-serious arguments. It is easy to get lost in them. Cited on page 265.
- 203 M. FATEMI, P. L. OGBURN & J. F. GREENLEAF, Fetal stimulation by pulsed diagnostic ultrasound, Journal of Ultrasound in Medicine 20, pp. 883–889, 2001. See also M. FATEMI, A. ALIZAD & J. F. GREENLEAF, Characteristics of the audio sound generated by ultrasound imaging systems, Journal of the Acoustical Society of America 117, pp. 1448–1455, 2005. Cited on pages 266 and 267.
- **204** I know a female gynecologist who, during her own pregnancy, imaged her child *every day* with her ultrasound machine. The baby was born with strong hearing problems that did not go away. Cited on page 267.
- **205** R. Mole, *Possible hazards of imaging and Doppler ultrasound in obstetrics*, Birth Issues in Perinatal Care 13, pp. 29–37, 2007. Cited on page 267.
- 206 A. L. Hodgkin & A. F. Huxley, A quantitative description of membrane current and its application to conduction and excitation in nerve, Journal of Physiology 117, pp. 500–544, 1952. This famous paper of theoretical biology earned the authors the Nobel Prize in Medicine in 1963. Cited on page 268.
- **207** T. FILIPPOV, *The Versatile Soliton*, Springer Verlag, 2000. See also J. S. Russel, *Report of the Fourteenth Meeting of the British Association for the Advancement of Science*, Murray, London, 1844, pp. 311–390. Cited on pages 268 and 271.
- **208** R. S. WARD, Solitons and other extended field configurations, preprint at arxiv.org/abs/hep-th/0505135. Cited on page 271.
- **209** D. B. Bahr, W. T. Pfeffer & R. C. Browning, *The surprising motion of ski moguls*, Physics Today **62**, pp. 68–69, November 2009. Cited on page 271.
- **210** N. J. Zabusky & M. D. Kruskal, *Interaction of solitons in a collisionless plasma and the recurrence of initial states*, Physical Review Letters 15, pp. 240–243, 1965. Cited on page 271.
- **211** O. Muskens, *De kortste knal ter wereld*, Nederlands tijdschrift voor natuurkunde pp. 70–73, 2005. Cited on page 271.
- **212** E. Heller, *Freak waves: just bad luck, or avoidable?*, Europhysics News pp. 159–161, September/October 2005, downloadable at www.europhysicsnews.org. Cited on page 272.
- 213 See the beautiful article by D. AARTS, M. SCHMIDT & H. LEKKERKER, *Directe visuele waarneming van thermische capillaire golven*, Nederlands tijdschrift voor natuurkunde 70, pp. 216–218, 2004. Cited on page 273.
- 214 For more about the ocean sound channel, see the novel by Tom Clancy, *The Hunt for Red October*. See also the physics script by R. A. Muller, *Government secrets of the oceans, atmosphere, and UFOs*, web.archive.org/web/*/muller.lbl.gov/teaching/Physics10/chapters/9-SecretsofUFOs.html 2001. Cited on page 274.

B. WILSON, R. S. BATTY & L. M. DILL, Pacific and Atlantic herring produce burst pulse sounds, Biology Letters 271, number S3, 7 February 2004. Cited on page 274.

- **216** See for example the article by G. Fritsch, *Infraschall*, Physik in unserer Zeit 13, pp. 104–110, 1982. Cited on page 277.
- 217 Wavelet transformations were developed by the French mathematicians Alex Grossmann, Jean Morlet and Thierry Paul. The basic paper is A. GROSSMANN, J. MORLET & T. PAUL, Integral transforms associated to square integrable representations, Journal of Mathematical Physics 26, pp. 2473–2479, 1985. For a modern introduction, see STÉPHANE MALLAT, A Wavelet Tour of Signal Processing, Academic Press, 1999. Cited on page 277.
- **218** P. Manogg, *Knall und Superknall beim Überschallflug*, Der mathematische und naturwissenschaftliche Unterricht 35, pp. 26–33, 1982. Cited on page 278.
- **219** See the excellent introduction by L. ELLERBREOK & VAN DEN HOORN, *In het kielzog van Kelvin*, Nederlands tijdschrift voor natuurkunde 73, pp. 310–313, 2007. Cited on page 279.
- **220** JAY INGRAM, *The Velocity of Honey And More Science of Everyday Life*, Viking, 2003. See also W. W. L. Au & J. A. Simmons, *Echolocation in dolphins and bats*, Physics Today 60, pp. 40–45, 2007. Cited on page 280.
- M. Boiti, J.-P. Leon, L. Martina & F. Pempinelli, Scattering of localized solitons in the plane, Physics Letters A 132, pp. 432–439, 1988, A. S. Fokas & P. M. Santini, Coherent structures in multidimensions, Physics Review Letters 63, pp. 1329–1333, 1989, J. Hietarinta & R. Hirota, Multidromion solutions to the Davey-Stewartson equation, Physics Letters A 145, pp. 237–244, 1990. Cited on page 280.
- The sound frequency change with bottle volume is explained on hyperphysics.phy-astr.gsu. edu/Hbase/Waves/cavity.html. Cited on page 281.
- **223** A passionate introduction is Neville H. Fletcher & Thomas D. Rossing, *The Physics of Musical Instruments*, second edition, Springer 2000. Cited on page 281.
- 224 M. Ausloos & D. H. Berman, *Multivariate Weierstrass–Mandelbrot function*, Proceedings of the Royal Society in London A 400, pp. 331–350, 1985. Cited on page 283.
- 225 Catechism of the Catholic Church, Part Two, Section Two, Chapter One, Article 3, statements 1376, 1377 and 1413, found at www.vatican.va/archive/ENG0015/__P41.HTM or www.vatican.va/archive/ITA0014/__P40.HTM with their official explanations on www.vatican.va/archive/compendium_ccc/documents/archive_2005_compendium-ccc_en.html and www.vatican.va/archive/compendium_ccc/documents/archive_2005_compendium-ccc_it. html. Cited on page 285.
- 226 The original text of the 1633 conviction of Galileo can be found on it.wikisource.org/wiki/ Sentenza_di_condanna_di_Galileo_Galilei. Cited on page 285.
- The retraction that Galileo was forced to sign in 1633 can be found on it.wikisource.org/wiki/Abiura_di_Galileo_Galilei. Cited on page 285.
- **228** M. Artigas, *Un nuovo documento sul caso Galileo: EE 291*, Acta Philosophica 10, pp. 199–214, 2001. Cited on page 286.
- 229 Most of these points are made, directly or indirectly, in the book by Annibale Fantoli, Galileo: For Copernicanism and for the Church, Vatican Observatory Publications, second edition, 1996, and by George Coyne, director of the Vatican observatory, in his speeches and publications, for example in G. Coyne, Galileo: for Copernicanism and for the church, Zwoje 3/36, 2003, found at www.zwoje-scrolls.com/zwoje36/text05p.htm. Cited on page 286.

- **230** T. A. McMahob & J. Tyler Bonner, Form und Leben Konstruktion vom Reißbrett der Natur, Spektrum Verlag, 1985. Cited on page 287.
- **231** G. W. Koch, S. C. Sillett, G. M. Jennings & S. D. Davis, *The limits to tree height*, Nature 428, pp. 851–854, 2004. Cited on page 287.
- A simple article explaining the tallness of trees is A. MINEYEV, Trees worthy of Paul Bunyan, Quantum pp. 4–10, January–February 1994. (Paul Bunyan is a mythical giant lumberjack who is the hero of the early frontier pioneers in the United States.) Note that the transport of liquids in trees sets no limits on their height, since water is pumped up along tree stems (except in spring, when it is pumped up from the roots) by evaporation from the leaves. This works almost without limits because water columns, when nucleation is carefully avoided, can be put under tensile stresses of over 100 bar, corresponding to 1000 m. See also P. Nobel, Plant Physiology, Academic Press, 2nd Edition, 1999. An artificial tree though extremely small using the same mechanism was built and studied by T. D. Wheeler & A. D. Stroock, The transpiration of water at negative pressures in a synthetic tree, Nature 455, pp. 208–212, 2008. See also N. M. Holbrook & M. A. Zwieniecki, Transporting water to the top of trees, Physics Today pp. 76–77, 2008. Cited on pages 287 and 304.
- **233** Such information can be taken from the excellent overview article by M. F. ASHBY, *On the engineering properties of materials*, Acta Metallurgica 37, pp. 1273–1293, 1989. The article explains the various general criteria which determine the selection of materials, and gives numerous tables to guide the selection. Cited on page 288.
- 234 See the beautiful paper by S. E. VIRGO, *Loschmidt'S number*, Science Progress 27, pp. 634–649, 1933. It is also available iin HTML format on the internet. Cited on pages 289 and 291.
- **235** For a photograph of a single barium *atom* named Astrid see Hans Dehmelt, *Experiments with an isolated subatomic particle at rest*, Reviews of Modern Physics **62**, pp. 525–530, 1990. For another photograph of a barium *ion*, see W. Neuhauser, M. Hohenstatt, P. E. Toschek & H. Dehmelt, *Localized visible Ba*⁺ *mono-ion oscillator*, Physical Review A **22**, pp. 1137–1140, 1980. See also the photograph on page 293. Cited on page 292.
- **236** Holograms of atoms were first produced by Hans-Werner Fink & al., *Atomic resolution in lens-less low-energy electron holography*, Physical Review Letters 67, pp. 1543–1546, 1991. Cited on page 292.
- **237** A single-atom laser was built in 1994 by K. An, J. J. Childs, R. R. Dasari & M. S. Feld, *Microlaser: a laser with one atom in an optical resonator*, Physical Review Letters 73, p. 3375, 1994. Cited on page 292.
- 238 The photograph on the left of Figure 194 on page 293 is the first image that showed subatomic structures (visible as shadows on the atoms). It was published by F. J. GIESSIBL, S. HEMBACHER, H. BIELEFELDT & J. MANNHART, Subatomic features on the silicon (111)-(7x7) surface observed by atomic force microscopy, Science 289, pp. 422 425, 2000. Cited on page 294.
- **239** See for example C. Schiller, A. A. Koomans, van Rooy, C. Schönenberger & H. B. Elswijk, *Decapitation of tungsten field emitter tips during sputter sharpening*, Surface Science Letters **339**, pp. L925–L930, 1996. Cited on page 294.
- **240** U. WEIERSTALL & J. C. H. SPENCE, An STM with time-of-flight analyzer for atomic species identification, MSA 2000, Philadelphia, Microscopy and Microanalysis 6, Supplement 2, p. 718, 2000. Cited on page 294.
- **241** P. Krehl, S. Engemann & D. Schwenkel, The puzzle of whip cracking uncovered by a correlation of whip-tip kinematics with shock wave emission, Shock Waves 8, pp. 1–9, 1998.

The authors used high-speed cameras to study the motion of the whip. A new aspect has been added by A. Goriely & T. McMillen, *Shape of a cracking whip*, Physical Review Letters 88, p. 244301, 2002. This article focuses on the tapered shape of the whip. However, the neglection of the tuft – a piece at the end of the whip which is required to make it crack – in the latter paper shows that there is more to be discovered still. Cited on page 297.

- **242** Z. Sheng & K. Yamafuji, Realization of a Human Riding a Unicycle by a Robot, Proceedings of the 1995 IEEE International Conference on Robotics and Automation, Vol. 2, pp. 1319 1326, 1995. Cited on page 298.
- **243** On human unicycling, see JACK WILEY, *The Complete Book of Unicycling*, Lodi, 1984, and SEBASTIAN HOEHER, *Einradfahren und die Physik*, Reinbeck, 1991. Cited on page 298.
- **244** W. Thomson, *Lecture to the Royal Society of Edinburgh*, 18 February 1867, Proceedings of the Royal Society in Edinborough 6, p. 94, 1869. Cited on page 298.
- S. T. THORODDSEN & A. Q. SHEN, *Granular jets*, Physics of Fluids 13, pp. 4–6, 2001, and A. Q. SHEN & S. T. THORODDSEN, *Granular jetting*, Physics of Fluids 14, p. S3, 2002, Cited on page 298.
- 246 M. J. Hancock & J. W. M. Bush, Fluid pipes, Journal of Fluid Mechanics 466, pp. 285–304, 2002. A. E. Hosoi & J. W. M. Bush, Evaporative instabilities in climbing films, Journal of Fluid Mechanics 442, pp. 217–239, 2001. J. W. M. Bush & A. E. Hasha, On the collision of laminar jets: fluid chains and fishbones, Journal of Fluid Mechanics 511, pp. 285–310, 2004. Cited on page 302.
- **247** The present record for negative pressure in water was achieved by Q. Zheng, D. J. Durben, G. H. Wolf & C. A. Angell, *Liquids at large negative pressures: water at the homogeneous nucleation limit*, Science 254, pp. 829–832, 1991. Cited on page 304.
- **248** H. Maris & S. Balibar, *Negative pressures and cavitation in liquid helium*, Physics Today 53, pp. 29–34, 2000. Cited on page 304.
- **249** The present state of our understanding of turbulence is described by G. Falkovich & K. P. Sreenivasan, *Lessons from hydrodynamic turbulence*, Physics Today **59**, pp. 43–49, 2006. Cited on page 306.
- 250 K. Weltner, A comparison of explanations of aerodynamical lifting force, American Journal of Physics 55, pp. 50–54, 1987, K. Weltner, Aerodynamic lifting force, The Physics Teacher 28, pp. 78–82, 1990. See also the user.uni-frankfurt.de/~weltner/Flight/PHYSIC4. htm and the www.av8n.com/how/htm/airfoils.html websites. Cited on page 306.
- **251** S. GEKLE, I. R. PETERS, J. M. GORDILLO, D. VAN DER MEER & D. LOHSE, Supersonic air flow due to solid-liquid impact, Physicl Review Letters 104, p. 024501, 2010. Films of the effect can be found at physics.aps.org/articles/v3/4. Cited on page 308.
- 252 See www.esa.int/esaCP/SEMER89U7TG index 0.html. Cited on page 308.
- **253** For a fascinating account of the passion and the techniques of apnoea diving, see UMBERTO PELIZZARI, *L'Homme et la mer*, Flammarion, 1994. Palizzari cites and explains the saying by Enzo Maiorca: 'The first breath you take when you come back to the surface is like the first breath with which you enter life.' Cited on page 308.
- 254 Lydéric Bocquet, *The physics of stone skipping*, American Journal of Physics 17, pp. 150–155, 2003. The present recod holder is Kurt Steiner, with 40 skips. See pastoneskipping.com/steiner.htm and www.stoneskipping.com. The site www.yeeha.net/nassa/guin/g2.html is by the a previous world record holder, Jerdome Coleman-McGhee. Cited on page 310.
- 255 S. F. KISTLER & L. E. SCRIVEN, The teapot effect: sheetforming flows with deflection, wetting, and hysteresis, Journal of Fluid Mechanics 263, pp. 19–62, 1994. Cited on page 313.

- **256** J. WALKER, Boiling and the Leidenfrost effect, a chapter from DAVID HALLIDAY, ROBERT RESNICK & JEARL WALKER, Fundamentals of Physics, Wiley, 2007. The chapter can also be found on the internet as pdf file. Cited on page 313.
- **257** E. HOLLANDER, *Over trechters en zo ...*, Nederlands tijdschrift voor natuurkunde **68**, p. 303, 2002. Cited on page 314.
- **258** S. DORBOLO, H. CAPS & N. VANDEWALLE, Fluid instabilities in the birth and death of antibubbles, New Journal of Physics 5, p. 161, 2003. Cited on page 314.
- **259** T. T. Lim, *A note on the leapfrogging between two coaxial vortex rings at low Reynolds numbers*, Physics of Fluids 9, pp. 239–241, 1997. Cited on page 316.
- **260** P. Aussillous & D. Quéré, *Properties of liquid marbles*, Proc. Roy. Soc. London 462, pp. 973–999, 2006, and references therein. Cited on page 316.
- 261 Thermostatics and thermodynamics is difficult to learn also because it was not discovered in a systematic way. See C. TRUESDELL, *The Tragicomical History of Thermodynamics 1822–1854*, Springer Verlag, 1980. An excellent advanced textbook on thermostatics and thermodynamics is LINDA REICHL, *A Modern Course in Statistical Physics*, Wiley, 2nd edition, 1998. Cited on page 319.
- Gas expansion was the main method used for the definition of the official temperature scale. Only in 1990 were other methods introduced officially, such as total radiation thermometry (in the range 140 K to 373 K), noise thermometry (2 K to 4 K and 900 K to 1235 K), acoustical thermometry (around 303 K), magnetic thermometry (0.5 K to 2.6 K) and optical radiation thermometry (above 730 K). Radiation thermometry is still the central method in the range from about 3 K to about 1000 K. This is explained in detail in R. L. Rusby, R. P. Hudson, M. Durieux, J. F. Schooley, P. P. M. Steur & C. A. Swenson, *The basis of the ITS-90*, Metrologia 28, pp. 9–18, 1991. On the water boiling point see also Ref. 288. Cited on pages 320, 468, and 472.
- **263** Other methods to rig lottery draws made use of balls of different mass or of balls that are more polished. One example of such a scam was uncovered in 1999. Cited on page 319.
- **264** See for example the captivating text by GINO SEGRÈ, A Matter of Degrees: What Temperature Reveals About the Past and Future of Our Species, Planet and Universe, Viking, New York, 2002. Cited on page 321.
- 265 D. Karstädt, F. Pinno, K.-P. Möllmann & M. Vollmer, Anschauliche Wärmelehre im Unterricht: ein Beitrag zur Visualisierung thermischer Vorgänge, Praxis der Naturwissenschaften Physik 5-48, pp. 24–31, 1999, K.-P. Möllmann & M. Vollmer, Eine etwas andere, physikalische Sehweise Visualisierung von Energieumwandlungen und Strahlungsphysik für die (Hochschul-)lehre, Physikalische Bllätter 56, pp. 65–69, 2000, D. Karstädt, K.-P. Möllmann, F. Pinno & M. Vollmer, There is more to see than eyes can detect: visualization of energy transfer processes and the laws of radiation for physics education, The Physics Teacher 39, pp. 371–376, 2001, K.-P. Möllmann & M. Vollmer, Infrared thermal imaging as a tool in university physics education, European Journal of Physics 28, pp. S37–S50, 2007. Cited on page 321.
- **266** See for example the article by H. Preston-Thomas, *The international temperature scale of 1990 (ITS-90)*, Metrologia 27, pp. 3–10, 1990, and the errata H. Preston-Thomas, *The international temperature scale of 1990 (ITS-90)*, Metrologia 27, p. 107, 1990, Cited on page 326.
- **267** For an overview, see Christian Enss & Siegfried Hunklinger, *Low-Temperature Physics*, Springer, 2005. Cited on page 326.

- 268 The famous paper on Brownian motion which contributed so much to Einstein's fame is A. EINSTEIN, Über die von der molekularkinetischen Theorie der Wärme geforderte Bewegung von in ruhenden Flüssigkeiten suspendierten Teilchen, Annalen der Physik 17, pp. 549–560, 1905. In the following years, Einstein wrote a series of further papers elaborating on this topic. For example, he published his 1905 Ph.D. thesis as A. EINSTEIN, Eine neue Bestimmung der Moleküldimensionen, Annalen der Physik 19, pp. 289–306, 1906, and he corrected a small mistake in A. EINSTEIN, Berichtigung zu meiner Arbeit: 'Eine neue Bestimmung der Moleküldimensionen', Annalen der Physik 34, pp. 591–592, 1911, where, using new data, he found the value 6.6 · 10²³ for Avogadro's number. However, five years before Smoluchowski and Einstein, a much more practically-minded man had made the same calculations, but in a different domain: the mathematician Louis Bachelier did so in his PhD about stock options; this young financial analyst was thus smarter than Einstein. Cited on page 327.
- **269** The first experimental confirmation of the prediction was performed by J. Perrin, Comptes Rendus de l'Académie des Sciences 147, pp. 475–476, and pp. 530–532, 1908. He masterfully sums up the whole discussion in Jean Perrin, *Les atomes*, Librarie Félix Alcan, Paris, 1913. Cited on page 328.
- **270** PIERRE GASPARD & al., Experimental evidence for microscopic chaos, Nature 394, p. 865, 27 August 1998. Cited on page 329.
- **271** An excellent introduction into the physics of heat is the book by LINDA REICHL, *A Modern Course in Statistical Physics*, Wiley, 2nd edition, 1998. Cited on page 329.
- **272** F. HERRMANN, *Mengenartige Größen im Physikunterricht*, Physikalische Blätter 54, pp. 830–832, September 1998. See also his lecture notes on general introductory physics on the website www.physikdidaktik.uni-karlsruhe.de/skripten. Cited on pages 198, 318, and 330.
- **273** These points are made clearly and forcibly, as is his style, by VAN KAMPEN, *Entropie*, Nederlands tijdschrift voor natuurkunde 62, pp. 395–396, 3 December 1996. Cited on page 332
- 274 This is a disappointing result of all efforts so far, as Grégoire Nicolis always stresses in his university courses. Seth Lloyd has compiled a list of 31 proposed definitions of complexity, containing among others, fractal dimension, grammatical complexity, computational complexity, thermodynamic depth. See, for example, a short summary in Scientific American p. 77, June 1995. Cited on page 332.
- 275 Minimal entropy is discussed by L. Szilard, Über die Entropieverminderung in einem thermodynamischen System bei Eingriffen intelligenter Wesen, Zeitschrift für Physik 53, pp. 840–856, 1929. This classic paper can also be found in English translation in his collected works. Cited on page 333.
- **276** G. COHEN-TANNOUDJI, Les constantes universelles, Pluriel, Hachette, 1998. See also L. Brillouin, Science and Information Theory, Academic Press, 1962. Cited on page 333.
- 277 H. W. ZIMMERMANN, Particle entropies and entropy quanta IV: the ideal gas, the second law of thermodynamics, and the P-t uncertainty relation, Zeitschrift für physikalische Chemie 217, pp. 55–78, 2003, and H. W. ZIMMERMANN, Particle entropies and entropy quanta V: the P-t uncertainty relation, Zeitschrift für physikalische Chemie 217, pp. 1097–1108, 2003. Cited on pages 333 and 334.
- 278 See for example A. E. SHALYT-MARGOLIN & A. YA. TREGUBOVICH, Generalized uncertainty relation in thermodynamics, arxiv.org/abs/gr-qc/0307018, or J. Uffink & J. van Lith-van Dis, Thermodynamic uncertainty relations, Foundations of Physics 29, p. 655, 1999. Cited on page 333.

- **279** B. LAVENDA, *Statistical Physics: A Probabilistic Approach*, Wiley-Interscience, New York, 1991. Cited on pages 333 and 334.
- **280** The quote given is found in the introduction by George Wald to the text by LAWRENCE J. HENDERSON, *The Fitness of the Environment*, Macmillan, New York, 1913, reprinted 1958. Cited on page 334.
- **281** A fascinating introduction to chemistry is the text by John Emsley, *Molecules at an Exhibition*, Oxford University Press, 1998. Cited on page 334.
- **282** B. Polster, *What is the best way to lace your shoes?*, Nature 420, p. 476, 5 December 2002. Cited on page 336.
- 283 L. Boltzmann, Über die mechanische Bedeutung des zweiten Huaptsatzes der Wärmetheorie, Sitzungsberichte der königlichen Akademie der Wissenschaften in Wien 53, pp. 155–220, 1866. Cited on page 337.
- **284** H. DE LANG, *Moleculaire gastronomie*, Nederlands tijdschrift voor natuurkunde 74, pp. 431–433, 2008. Cited on page 340.
- **285** EMILE BOREL, *Introduction géométrique à la physique*, Gauthier-Villars, 1912. Cited on page 340.
- **286** See V. L. Telegdi, *Enrico Fermi in America*, Physics Today 55, pp. 38–43, June 2002. Cited on page 341.
- **287** K. Schmidt-Nielsen, Desert Animals: Physiological Problems of Heat and Water, Oxford University Press, 1964. Cited on page 341.
- **288** Following a private communication by Richard Rusby, this is the value of 1997, whereas it was estimated as 99.975°C in 1989, as reported by GARETH JONES & RICHARD RUSBY, Official: water boils at 99.975°C, Physics World 2, pp. 23–24, September 1989, and R. L. Rusby, Ironing out the standard scale, Nature 338, p. 1169, March 1989. For more details on temperature measurements, see Ref. 262. Cited on pages 341 and 466.
- **289** Why entropy is created when information is erased, but not when it is acquired, is explained in C. H. Bennett & R. Landauer, *Fundamental Limits of Computation*, Scientific American 253:1, pp. 48–56, 1985. The conclusion: we should pay to throw the newspaper away, not to buy it. Cited on page 343.
- **290** See, for example, G. Swift, *Thermoacoustic engines and refrigerators*, Physics Today 48, pp. 22–28, July 1995. Cited on page 345.
- **291** Quoted in D. Campbell, J. Crutchfield, J. Farmer & E. Jen, *Experimental mathematics: the role of computation in nonlinear science*, Communications of the Association of Computing Machinery **28**, pp. 374–384, 1985. Cited on page 348.
- **292** For more about the shapes of snowflakes, see the famous book by W. A. Bentley & W. J. Humphreys, *Snow Crystals*, Dover Publications, New York, 1962. This second printing of the original from 1931 shows a large part of the result of Bentley's lifelong passion, namely several thousand photographs of snowflakes. Cited on page 348.
- **293** K. Schwenk, *Why snakes have forked tongues*, Science **263**, pp. 1573–1577, 1994. Cited on page 351.
- E. Martínez, C. Pérez-Penichet, O. Sotolongo-Costa, O. Ramos, K.J. Måløy, S. Douady, E. Altshuler, *Uphill solitary waves in granular flows*, Physical Review 75, p. 031303, 2007, and E. Altshuler, O. Ramos, E. Martínez, A.J. Batista-Leyva, A. Rivera & K. E. Bassler, *Sandpile formation by revolving rivers*, Physical Review Letters 91, p. 014501, 2003. Cited on page 353.

P. B. UMBANHOWAR, F. MELO & H. L. SWINNEY, Localized excitations in a vertically vibrated granular layer, Nature 382, pp. 793–796, 29 August 1996. Cited on page 353.

- **296** D. K. Campbell, S. Flach & Y. S. Kivshar, Localizing energy through nonlinearity and discreteness, Physics Today 57, pp. 43–49, January 2004. Cited on page 354.
- **297** B. Andreotti, *The song of dunes as a wave-particle mode locking*, Physical Review Letters 92, p. 238001, 2004. Cited on page 354.
- 298 D. C. Mays & B. A. Faybishenko, Washboards in unpaved highways as a complex dynamic system, Complexity 5, pp. 51–60, 2000. See also N. Taberlet, S. W. Morris & J. N. McElwaine, Washboard road: the dynamics of granular ripples formed by rolling wheels, Physical Review Letters 99, p. 068003, 2007. Cited on pages 354 and 478.
- **299** K. KÖTTER, E. GOLES & M. MARKUS, Shell structures with 'magic numbers' of spheres in a swirled disk, Physical Review E 60, pp. 7182–7185, 1999. Cited on page 355.
- **300** A good introduction is the text by DANIEL WALGRAEF, Spatiotemporal Pattern Formation, With Examples in Physics, Chemistry and Materials Science, Springer 1996. Cited on page 355.
- **301** For an overview, see the Ph.D. thesis by Joceline Lega, *Défauts topologiques associés à la brisure de l'invariance de translation dans le temps*, Université de Nice, 1989. Cited on page 356.
- **302** An idea of the fascinating mechanisms at the basis of the heart beat is given by A. Babloyantz & A. Destexhe, *Is the normal heart a periodic oscillator?*, Biological Cybernetics **58**, pp. 203–211, 1989. Cited on page 357.
- **303** For a short, modern overview of turbulence, see L. P. KADANOFF, *A model of turbulence*, Physics Today **48**, pp. 11–13, September 1995. Cited on page 358.
- **304** For a clear introduction, see T. SCHMIDT & M. MAHRL, A simple mathematical model of a dripping tap, European Journal of Physics 18, pp. 377–383, 1997. Cited on page 359.
- An overview of science humour can be found in the famous anthology compiled by R. L. Weber, edited by E. Mendoza, *A Random Walk in Science*, Institute of Physics, 1973. It is also available in several expanded translations. Cited on page 359.
- **306** W. Dreybrodt, *Physik von Stalagmiten*, Physik in unserer Zeit pp. 25–30, Physik in unserer Zeit February 2009. Cited on page 359.
- **307** K. MERTENS, V. PUTKARADZE & P. VOROBIEFF, Braiding patterns on an inclined plane, Nature 430, p. 165, 2004. Cited on page 360.
- These beautifully simple experiments were published in G. MÜLLER, Starch columns: analog model for basalt columns, Journal of Geophysical Research 103, pp. 15239–15253, 1998, in G. MÜLLER, Experimental simulation of basalt columns, Journal of Volcanology and Geothermal Research 86, pp. 93–96, 1998, and in G. MÜLLER, Trocknungsrisse in Stärke, Physikalische Blätter 55, pp. 35–37, 1999. Cited on page 361.
- **309** To get a feeling for viscosity, see the fascinating text by STEVEN VOGEL, *Life in Moving Fluids: the Physical Biology of Flow*, Princeton University Press, 1994. Cited on page 361.
- 310 B. Hof, C. W. H. VAN DOORNE, J. WESTERWEEL, F. T. M. NIEUWSTADT, H. WEDIN, R. KERSWELL, F. WALEFFE, H. FAISST & B. ECKHARDT, Experimental observation of nonlinear traveling waves in turbulent pipe flow, Science 305, pp. 1594–1598, 2004. See also B. Hof & al., Finite lifetime of turbulence in shear flows, Nature 443, p. 59, 2006. Cited on page 362.
- **311** A fascinating book on the topic is Kenneth Laws & Martha Swope, *Physics and the Art of Dance: Understanding Movement*, Oxford University Press 2002. Cited on page 363.

- 312 The fascinating variation of snow crystals is presented in C. MAGONO & C. W. LEE, *Mete-orological classification of natural snow crystals*, Journal of the Faculty of Science, Hokkaido University Ser. VII, II, pp. 321–325, 1966, also online at the eprints.lib.hokudai.ac.jp website. Cited on page 364.
- **313** Josef H. Reichholf, *Eine kurze Naturgeschichte des letzten Jahrtausends*, S. Fischer Verlag, 2007. Cited on page 364.
- **314** See for example, E. F. Bunn, *Evolution and the second law of thermodynamics*, American Journal of Physics 77, pp. 922–925, 2009. Cited on page 364.
- **315** A good introduction of the physics of bird swarms is T. Feder, *Statistical physics is for the birds*, Physics Today **60**, pp. 28–30, October 2007. Cited on page 364.
- **316** J. J. LISSAUER, *Chaotic motion in the solar system*, Reviews of Modern Physics 71, pp. 835–845, 1999. Cited on page 366.
- **317** See JEAN-PAUL DUMONT, *Les écoles présocratiques*, Folio Essais, Gallimard, 1991, p. 426. Cited on page 367.
- 318 For information about the number π , and about some other mathematical constants, the website oldweb.cecm.sfu.ca/pi/pi.html provides the most extensive information and references. It also has a link to the many other sites on the topic, including the overview at mathworld.wolfram.com/Pi.html. Simple formulae for π are

$$\pi + 3 = \sum_{n=1}^{\infty} \frac{n \, 2^n}{\binom{2n}{n}} \tag{152}$$

or the beautiful formula discovered in 1996 by Bailey, Borwein and Plouffe

$$\pi = \sum_{n=0}^{\infty} \frac{1}{16^n} \left(\frac{4}{8n+1} - \frac{2}{8n+4} - \frac{1}{8n+5} - \frac{1}{8n+6} \right) . \tag{153}$$

The mentioned site also explains the newly discovered methods for calculating specific binary digits of π without having to calculate all the preceding ones. The known digits of π pass all tests of randomness, as the mathworld.wolfram.com/PiDigits.html website explains. However, this property, called *normality*, has never been proven; it is the biggest open question about π . It is possible that the theory of chaotic dynamics will lead to a solution of this puzzle in the coming years.

Another method to calculate π and other constants was discovered and published by D. V. Chudnovsky & G. V. Chudnovsky, *The computation of classical constants*, Proceedings of the National Academy of Sciences (USA) 86, pp. 8178–8182, 1989. The Chudnowsky brothers have built a supercomputer in Gregory's apartment for about 70 000 euros, and for many years held the record for calculating the largest number of digits of π . They have battled for decades with Kanada Yasumasa, who held the record in 2000, calculated on an industrial supercomputer. However, the record number of (consecutive) digits in 2010 was calculated in 123 days on a simple desktop PC by Fabrice Bellard, using a Chudnovsky formula. Bellard calculated over 2.7 million million digits, as told on bellard.org. New formulae to calculate π are still occasionally discovered.

For the calculation of Euler's constant γ see also D. W. DeTemple, A quicker convergence to Euler's constant, The Mathematical Intelligencer, pp. 468–470, May 1993. Cited on pages 368 and 397.

319 GEORGE BIRKBECK NORMAN HILL, *Johnsonian Miscellanies*, Clarendon Press, 1897, in Seward's Biographiana/ Cited on page 372.

320 The first written record of the letter U seems to be LEON BATTISTA ALBERTI, *Grammatica della lingua toscana*, 1442, the first grammar of a modern (non-latin) language, written by a genius that was intellectual, architect and the father of cryptology. The first written record of the letter J seems to be ANTONIO DE NEBRIJA, *Gramática castellana*, 1492. Before writing it, Nebrija lived for ten years in Italy, so that it is possible that the I/J distinction is of Italian origin as well. Nebrija was one of the most important Spanish scholars. Cited on page 373.

- **321** For more information about the letters thorn and eth, have a look at the extensive report to be found on the website www.everytype.com/standards/wynnyogh/thorn.html. Cited on page 373.
- **322** For a modern history of the English language, see DAVID CRYSTAL, *The Stories of English*, Allen Lane, 2004. Cited on page 373.
- **323** Hans Jensen, *Die Schrift*, Berlin, 1969, translated into English as *Sign, Symbol and Script:* an Account of Man's Efforts to Write, Putnam's Sons, 1970. Cited on page 374.
- **324** DAVID R. LIDE, editor, *CRC Handbook of Chemistry and Physics*, 78th edition, CRC Press, 1997. This classic reference work appears in a new edition every year. The full Hebrew alphabet is given on page 2-90. The list of abbreviations of physical quantities for use in formulae approved by ISO, IUPAP and IUPAC can also be found there.

However, the ISO 31 standard, which defines these abbreviations, costs around a thousand euro, is not available on the internet, and therefore can safely be ignored, like any standard that is supposed to be used in teaching but is kept inaccessible to teachers. Cited on pages 376 and 377.

- **325** See the mighty text by Peter T. Daniels & William Bright, *The World's Writing Systems*, Oxford University Press, 1996. Cited on page 376.
- 326 The story of the development of the numbers is told most interestingly by Georges Ifrah, *Histoire universelle des chiffres*, Seghers, 1981, which has been translated into several languages. He sums up the genealogy of the number signs in ten beautiful tables, one for each digit, at the end of the book. However, the book itself contains factual errors on every page, as explained for example in the review found at www.ams.org/notices/200201/rev-dauben.pdf and www.ams.org/notices/200202/rev-dauben.pdf. Cited on page 376.
- **327** See the for example the fascinating book by STEVEN B. SMITH, *The Great Mental Calculators The Psychology, Methods and Lives of the Calculating Prodigies*, Columbia University Press, 1983. The book also presents the techniques that they use, and that anybody else can use to emulate them. Cited on page 377.
- 328 See for example the article 'Mathematical notation' in the Encyclopedia of Mathematics, 10 volumes, Kluwer Academic Publishers, 1988–1993. But first all, have a look at the informative and beautiful jeff560.tripod.com/mathsym.html website. The main source for all these results is the classic and extensive research by Florian Cajori, A History of Mathematical Notations, 2 volumes, The Open Court Publishing Co., 1928–1929. The square root sign is used in Christoff Rudolff, Die Coss, Vuolfius Cephaleus Joanni Jung: Argentorati, 1525. (The full title was Behend vnnd Hubsch Rechnung durch die kunstreichen regeln Algebre so gemeinlicklich die Coss genent werden. Darinnen alles so treülich an tag gegeben, das auch allein auss vleissigem lesen on allen mündtliche vnterricht mag begriffen werden, etc.) Cited on page 377.
- **329** J. TSCHICHOLD, Formenwamdlungen der et-Zeichen, Stempel AG, 1953. Cited on page 379.

330 MALCOLM B. PARKES, *Pause and Effect: An Introduction to the History of Punctuation in the West*, University of California Press, 1993. Cited on page 379.

- **331** This is explained by Berthold Louis Ullman, *Ancient Writing and its Influence*, 1932. Cited on page 379.
- **332** PAUL LEHMANN, Erforschung des Mittelalters Ausgewählte Abhandlungen und Aufsätze, Anton Hiersemann, 1961, pp. 4–21. Cited on page 379.
- **333** Bernard Bischoff, *Paläographie des römischen Altertums und des abendländischen Mittelalters*, Erich Schmidt Verlag, 1979, pp. 215–219. Cited on page 379.
- **334** HUTTON WEBSTER, *Rest Days: A Study in Early Law and Morality*, MacMillan, 1916. The discovery of the unlucky day in Babylonia was made in 1869 by George Smith, who also rediscovered the famous *Epic of Gilgamesh*. Cited on page 380.
- 335 The connections between Greek roots and many French words and thus many English ones can be used to rapidly build up a vocabulary of ancient Greek without much study, as shown by the practical collection by J. Chaineux, Quelques racines grecques, Wetteren De Meester, 1929. See also Donald M. Ayers, English Words from Latin and Greek Elements, University of Arizona Press, 1986. Cited on page 381.
- **336** In order to write well, read WILLIAM STRUNK & E. B. WHITE, *The Elements of Style*, Macmillan, 1935, 1979, or WOLF SCHNEIDER, *Deutsch für Kenner Die neue Stilkunde*, Gruner und Jahr, 1987. Cited on page 382.
- Breteuil, Parc de Saint Cloud, 92310 Sèvres, France. All new developments concerning SI units are published in the journal *Metrologia*, edited by the same body. Showing the slow pace of an old institution, the BIPM launched a website only in 1998; it is now reachable at www.bipm.fr. See also the www.utc.fr/~tthomass/Themes/Unites/index.html website; this includes the biographies of people who gave their names to various units. The site of its British equivalent, www.npl.co.uk/npl/reference, is much better; it provides many details as well as the English-language version of the SI unit definitions. Cited on page 383.
- The bible in the field of time measurement is the two-volume work by J. Vanier & C. Audoin, *The Quantum Physics of Atomic Frequency Standards*, Adam Hilge, 1989. A popular account is Tony Jones, *Splitting the Second*, Institute of Physics Publishing, 2000.

The site opdafl.obspm.fr/www/lexique.html gives a glossary of terms used in the field. For precision *length* measurements, the tools of choice are special lasers, such as modelocked lasers and frequency combs. There is a huge literature on these topics. Equally large is the literature on precision *electric current* measurements; there is a race going on for the best way to do this: counting charges or measuring magnetic forces. The issue is still open. On *mass* and atomic mass measurements, see the volume on relativity. On high-precision *temperature* measurements, see Ref. 262. Cited on page 384.

Vol. II, page 65

- The unofficial prefixes were first proposed in the 1990s by Jeff K. Aronson of the University of Oxford, and might come into general usage in the future. Cited on page 385.
- 340 The most precise clock built in 2004, a caesium fountain clock, had a precision of one part in 10¹⁵. Higher precision has been predicted to be possible soon, among others by M. Такамото, F.-L. Hong, R. Higashi & H. Katori, *An optical lattice clock*, Nature 435, pp. 321–324, 2005. Cited on page 387.
- **341** J. Bergquist, ed., *Proceedings of the Fifth Symposium on Frequency Standards and Metrology*, World Scientific, 1997. Cited on page 387.
- **342** J. Short, *Newton's apples fall from grace*, New Scientist 2098, p. 5, 6 September 1997. More details can be found in R. G. Keesing, *The history of Newton's apple tree*, Contemporary

- Physics 39, pp. 377-391, 1998. Cited on page 388.
- 343 The various concepts are even the topic of a separate international standard, ISO 5725, with the title *Accuracy and precision of measurement methods and results*. A good introduction is JOHN R. TAYLOR, *An Introduction to Error Analysis: the Study of Uncertainties in Physical Measurements*, 2nd edition, University Science Books, Sausalito, 1997. Cited on page 389.
- 344 The most recent (2010) recommended values of the fundamental physical constants are found only on the website physics.nist.gov/cuu/Constants/index.html. This set of constants results from an international adjustment and is recommended for international use by the Committee on Data for Science and Technology (CODATA), a body in the International Council of Scientific Unions, which brings together the International Union of Pure and Applied Physics (IUPAP), the International Union of Pure and Applied Chemistry (IUPAC) and other organizations. The website of IUPAC is www.iupac.org. Cited on page 390.
- 345 Some of the stories can be found in the text by N. W. WISE, *The Values of Precision*, Princeton University Press, 1994. The field of high-precision measurements, from which the results on these pages stem, is a world on its own. A beautiful introduction to it is J. D. FAIRBANKS, B. S. DEAVER, C. W. EVERITT & P. F. MICHAELSON, eds., *Near Zero: Frontiers of Physics*, Freeman, 1988. Cited on page 390.
- **346** The details are given in the well-known astronomical reference, Kenneth Seidelmann, *Explanatory Supplement to the Astronomical Almanac*, 1992. Cited on page 395.





ACKNOWLEDGEMENTS

Many people who have kept their gift of curiosity alive have helped to make this project come true. Most of all, Saverio Pascazio has been – present or not – a constant reference for this project. Fernand Mayné, Anna Koolen, Ata Masafumi, Roberto Crespi, Serge Pahaut, Luca Bombelli, Herman Elswijk, Marcel Krijn, Marc de Jong, Martin van der Mark, Kim Jalink, my parents Peter and Isabella Schiller, Mike van Wijk, Renate Georgi, Paul Tegelaar, Barbara and Edgar Augel, M. Jamil, Ron Murdock, Carol Pritchard, Richard Hoffman, Stephan Schiller and, most of all, my wife Britta have all provided valuable advice and encouragement.

Many people have helped with the project and the collection of material. Most useful was the help of Mikael Johansson, Bruno Barberi Gnecco, Lothar Beyer, the numerous improvements by Bert Sierra, the detailed suggestions by Claudio Farinati, the many improvements by Eric Sheldon, the detailed suggestions by Andrew Young, the continuous help and advice of Jonatan Kelu, the corrections of Elmar Bartel, and in particular the extensive, passionate and conscientious help of Adrian Kubala.

Important material was provided by Bert Peeters, Anna Wierzbicka, William Beaty, Jim Carr, John Merrit, John Baez, Frank DiFilippo, Jonathan Scott, Jon Thaler, Luca Bombelli, Douglas Singleton, George McQuarry, Tilman Hausherr, Brian Oberquell, Peer Zalm, Martin van der Mark, Vladimir Surdin, Julia Simon, Antonio Fermani, Don Page, Stephen Haley, Peter Mayr, Allan Hayes, Norbert Dragon, Igor Ivanov, Doug Renselle, Wim de Muynck, Steve Carlip, Tom Bruce, Ryan Budney, Gary Ruben, Chris Hillman, Olivier Glassey, Jochen Greiner, squark, Martin Hardcastle, Mark Biggar, Pavel Kuzin, Douglas Brebner, Luciano Lombardi, Franco Bagnoli, Lukas Fabian Moser, Dejan Corovic, Paul Vannoni, John Haber, Saverio Pascazio, Klaus Finkenzeller, Leo Volin, Jeff Aronson, Roggie Boone, Lawrence Tuppen, Quentin David Jones, Arnaldo Uguzzoni, Frans van Nieuwpoort, Alan Mahoney, Britta Schiller, Petr Danecek, Ingo Thies, Vitaliy Solomatin, Carl Offner, Nuno Proença, Elena Colazingari, Paula Henderson, Daniel Darre, Wolfgang Rankl, John Heumann, Joseph Kiss, Martha Weiss, Antonio González, Antonio Martos, André Slabber, Ferdinand Bautista, Zoltán Gácsi, Pat Furrie, Michael Reppisch, Enrico Pasi, Thomas Köppe, Martin Rivas, Herman Beeksma, Tom Helmond, John Brandes, Vlad Tarko, Nadia Murillo, Ciprian Dobra, Romano Perini, Harald van Lintel, Andrea Conti, François Belfort, Dirk Van de Moortel, Heinrich Neumaier, Jarosław Królikowski, John Dahlman, Fathi Namouni, Paul Townsend, Sergei Emelin, Freeman Dyson, S.R. Madhu Rao, David Parks, Jürgen Janek, Daniel Huber, Alfons Buchmann, William Purves, Pietro Redondi, Damoon Saghian, Frank Sweetser, plus a number of people who wanted to remain unnamed.

The software tools were refined with extensive help on fonts and typesetting by Michael Zedler and Achim Blumensath and with the repeated and valuable support of Donald Arseneau; help came also from Ulrike Fischer, Piet van Oostrum, Gerben Wierda, Klaus Böhncke, Craig Upright, Herbert Voss, Andrew Trevorrow, Danie Els, Heiko Oberdiek, Sebastian Rahtz, Don Story, Vincent Darley, Johan Linde, Joseph Hertzlinger, Rick Zaccone, John Warkentin, Ulrich Diez, Uwe

Siart, Will Robertson, Joseph Wright, Enrico Gregorio, Rolf Niepraschk, Alexander Grahn and Werner Fabian.

The typesetting and book design is due to the professional consulting of Ulrich Dirr. The typography and the table on mathematical signs was greatly improved with the help of Johannes Küster. The design of the book and its website owe also much to the suggestions and support of my wife Britta.

I also thank the lawmakers and the taxpayers in Germany, who, in contrast to most other countries in the world, allow residents to use the local university libraries.

Since May 2007, the electronic edition and distribution of the Motion Mountain text is generously supported by the Klaus Tschira Foundation.

FILM CREDITS

The clear animation of a suspended spinning top, shown on page 134, was made for this text by Lucas Barbosa. The beautiful animation of the lunation on page 167 was calculated from actual astronomical data and is copyright and courtesy by Martin Elsässer. It can be found on his website www.mondatlas.de/lunation.html. The beautiful animation of the planets and planetoids on page 190 is copyright and courtesy by Hans-Christian Greier. It can be found on his wonderful website www.parallax.at. The film of an oscillating quartz on page 251 is copyright and courtesy of Micro Crystal, part of the Swatch Group, found at www.microcrystal.com. The animation illustrating group and wave velocity on page 257 is courtesy and copyright of the ISVR at the University of Southhampton. The animation illustrating the motion of molecules in air with sound, shown on page 265, is found on Wikimedia Commons. The films of solitons on page 270 and of dromions on page 280 are copyright and courtesy by Jarmo Hietarinta. They can be found on his website users.utu.fi/hietarin. The film of leapfrogging vortex rings on page 315 is copyright and courtesy by Lim Tee Tai. It can be found via his fluid dynamics website serve.me.nus.edu.sg. The film of the growing snowflake on page 355 is copyright and courtesy by Kenneth Libbrecht. It can be found on his website www.its.caltech.edu/~atomic/snowcrystals.

IMAGE CREDITS

The photograph of a peak in the Himalaya on the front cover is courtesy and copyright by Dave Thompson and used to be on his website www.daveontrek.co.uk. The lightning photograph on page 14 is courtesy and copyright by Harald Edens and found on the www.lightningsafety.noaa. gov/photos.htm and www.weather-photography.com websites. The motion illusion on page 18 is courtesy and copyright by Michael Bach and found on his website www.michaelbach.de/ot/ mot_rotsnake/index.html. It is a variation of the illusion by Kitaoka Akiyoshi found on www. ritsumei.ac.jp/~akitaoka and used here with his permission. The figures on pages 20, 55 and 179 were made especially for this text and are copyright by Luca Gastaldi. The high speed photograph of a bouncing tennis ball on page 20 is courtesy and copyright by the International Tennis Federation, and were provided by Janet Page. The figure of Etna on pages 22 and 137 is copyright and courtesy of Marco Fulle and taken from the wonderful website www.stromboli.net. The famous photograph of the Les Poulains and its lighthouse by Philip Plisson on page 23 is courtesy and copyright by Pechêurs d'Images; see the websites www.plisson.com and www. pecheurs-d-images.com. It is also found in Plisson's magnus opus La Mer, a stunning book of photographs of the sea. The picture on page 23 of Alexander Tsukanov jumping from one ultimate wheel to another is copyright and courtesy of the Moscow State Circus. The photograph of a deer on page 25 is copyright and courtesy of Tony Rodgers and taken from his website www. flickr.com/photos/moonm. The photographs of speed measurement devices on page 34 are courtesy and copyright of the Fachhochschule Koblenz, of Silva, of Tracer, and of Wikimedia. The

graph on page 37 is redrawn and translated from the wonderful book by Henk Tennekes, De wetten van de vliegkunst - Over stijgen, dalen, vliegen en zweven, Aramith Uitgevers, 1993. The photographs of the ping-pong ball on page 39 and of the dripping water tap on page 303 are copyright and courtesy of Andrew Davidhazy and found on his website www.rit.edu/~andpph. The photograph of the bouncing water droplet on page 39 are copyright and courtesy of Max Groenendijk and found on the website www.lightmotif.nl. The photograph of the precision sundial on page 43 is copyright and courtesy of Stefan Pietrzik and found at commons.wikimedia. org/wiki/Image:Präzissions-Sonnenuhr_mit_Sommerwalze.jpg The other clock photographs in the figure are from public domain sources as indicated. The graph on the scaling of biological rhythms on page 46 is drawn by the author using data from the European Molecular Biology Organisation found at www.nature.com/embor/journal/v6/nls/fig tab/7400425 f3.html and Enrique Morgado. The drawing of the human ear on page page 49 and on page 276 are copyright of Northwestern University and courtesy of Tim Hain; it is found on his website www.dizziness-and-balance.com/disorders/bppy/otoliths.html. The illustrations of the vernier caliper and the micrometer screw on page 52 and 62 are copyright of Medien Werkstatt, courtesy of Stephan Bogusch, and taken from their instruction course found on their website www. medien-werkstatt.de. The photo of the tiger on page 52 is copyright of Naples zoo (in Florida, not in Italy), and courtesy of Tim Tetzlaff; see their website at www.napleszoo.com. The other length measurement devices on page 52 are courtesy and copyright of Keyence and Leica Geosystems, found at www.leica-geosystems.com. The curvimeter photograph on page 54 is copyright and courtesy of Frank Müller and found on the www.wikimedia.org website. The crystal photograph on the left of page 57 is copyright and courtesy of Stephan Wolfsried and found on the www. mindat.org website. The crystal photograph on the right of page 57 is courtesy of Tullio Bernabei, copyright of Arch. Speleoresearch & Films/La Venta and found on the www.laventa.it and www.naica.com.mx websites. The hollow Earth figure on pages 59 and 263 is courtesy of Helmut Diel and was drawn by Isolde Diel. The wonderful photographs on page 67, page 135, page 153, page 169, page 185 and page 430 are courtesy and copyright by Anthony Ayiomamitis; the story of the photographs is told on his beautiful website at www.perseus.gr. The anticrepuscular photograph on page 68 is courtesy and copyright by Peggy Peterson. The rope images on page 69 are copyright and courtesy of Jakob Bohr. The image of the tight knot on page 70 is courtesy and copyright by Piotr Pieranski. The firing caterpillar figure of page 76 is courtesy and copyright of Stanley Caveney. The photograph of an airbag sensor on page 83 is courtesy and copyright of Bosch; the accelerometer picture is courtesy and copyright of Rieker Electronics; the three drawings of the human ear are copyright of Northwestern University and courtesy of Tim Hain and found on his website www.dizziness-and-balance.com/disorders/bppv/otoliths.html. The photograph of Orion on page 84 is courtesy and copyright by Matthew Spinelli. The photograph of Regulus and Mars on page 84 is courtesy and copyright of Jürgen Michelberger and found on www.jmichelberger.de. The photograph of N. decemspinosa on page 86 is courtesy and copyright of Robert Full, and found on his website rjf9.biol.berkeley.edu/twiki/bin/view/PolyPEDAL/ LabPhotographs. The photograph of P. ruralis on page 86 is courtesy and copyright of John Brackenbury, and part of his wonderful collection on the website www.sciencephoto.co.uk. The millipede photograph on page 86 is courtesy and copyright of David Parks, and found on his website www.mobot.org/mobot/madagascar/image.asp?relation=A71. The photograph of the Gekko climbing the bus window on page 86 is courtesy and copyright of Marcel Berendsen, and found on his website www.flickr.com/photos/berendm. The photograph of comet McNaught on page 88 is courtesy and copyright by its discoveror, Robert McNaught; it is taken from his website at www. mso.anu.edu.au/~rmn and is found also on antwrp.gsfc.nasa.gov/apod/ap070122.html. The sonoluminsecence picture on page 89 is courtesy and copyright of Detlev Lohse. The photograph of the standard kilogram on page 92 is courtesy and copyright by the Bureau International des Poids

et Mesures (BIPM). The photograph of Mendeleyev's balance on page 99 is copyright of Thinktank Trust and courtesy of Jack Kirby; it can be found on the www.birminghamstories.co.uk website. The photograph of the laboratory scale on page 99 is copyright and courtesy of Mettler-Toledo. The measured graph of the walking human on page 113 is courtesy and copyright of Ray McCoy. The photograph of the clock that needs no winding up is copyright Jaeger-LeCoultre and courtesy of Ralph Stieber. Its history and working are described in detail in a brochure available from the company. The company's website is www.Jaeger-LeCoultre.com. The photograph of the ship lift at Strépy-Thieux on page 120 is courtesy and copyright of Jean-Marie Hoornaert and found on Wikimedia Commons. The photograph of the Celtic wobble stone on page 121 is courtesy and copyright of Ed Keath and found on Wikimedia Commons. The photograph of Foucault's gyroscope on page 129 is courtesy and copyright of the museum of the CNAM, the Conservatoire National des Arts et Métiers in Paris, whose website is at www.arts-et-metiers.net. The photograph of the laser gyroscope on page 129 is courtesy and copyright of JAXA, the Japan Aerospace Exploration Agency, and found on their website at jda.jaxa.jp. The drawing of the precision laser gyroscope on page 130 is courtesy of Thomas Klügel and copyright of the Bundesamt für Kartographie und Geodäsie. The photograph of the instrument is courtesy and copyright of Carl Zeiss. The machine is located at the Fundamentalstation Wettzell, and its website is found at www.wettzell.ifag.de. The graph of the temperature record on page 139 is copyright and courtesy Jean Jouzel and Science/AAAS. Figure 104 on page 147 is courtesy and copyright of the international Gemini project (Gemini Observatory/Association of Universities for Research in Astronomy) at www.ausgo.unsw.edu.au and www.gemini.edu. The basilisk running over water, page 149 and on the back cover, is courtesy and copyright by the Belgian group TERRA vzw and found on their website www.terra.vzw.org. The water strider photograph on page 149 is courtesy and copyright by Charles Lewallen. The photograph of the water robot on page 149 is courtesy and copyright by the American Institute of Physics. The illustration of the motion of Mars on page 152 is courtesy and copyright of Tunc Tezel. The photograph of the precision pendulum clock on page 155 is copyright of Erwin Sattler OHG, Sattler OHG, Erwin and courtesy of Ms. Stephanie Sattler-Rick; it can be found at the www.erwinsattler.de website. The figure on the triangulation of the meridian of Paris on page 158 is copyright and courtesy of Ken Alder and found on his website www.kenalder.com. The photographs of the home version of the Cavendish experiment on page 159 are courtesy and copyright by John Walker and found on his website www.fourmilab. ch/gravitation/foobar. The photographs of the precision Cavendish experiment on page 160 are courtesy and copyright of the Eöt-Wash Group at the University of Washington and found at www.npl.washington.edu/eotwash. The geoid of page 162 is courtesy and copyright by the GeoForschungsZentrum Potsdam, found at www.gfz-potsdam.de. The moon maps on page 167 are courtesy of the USGS Astrogeology Research Program, astrogeology.usgs.gov, in particular Mark Rosek and Trent Hare. The graph of orbits on page 170 is courtesy and copyright of Geoffrey Marcy. The photograph of the tides on page 173 is copyright and courtesy of Gilles Régnier and found on his website www.gillesregnier.com; it also shows an animation of that tide over the whole day. The pictures of fast descents on snow on page 182 are copyright and courtesy of Simone Origone, www.simoneorigone.it, and of Éric Barone, www.ericbarone.com. The photograph of the Galilean satellites on page 183 is courtesy and copyright by Robin Scagell and taken from his website www.galaxypix.com. The pictures of solar eclipses on page 193 are courtesy and copyright by the Centre National d'Etudes Spatiales, at www.cnes.fr, and of Laurent Laveder, from his beautiful site at www.PixHeaven.net. The pictures of daisies on page 211 are copyright and courtesy of Giorgio Di Iorio, found on his website www.flickr.com/photos/gioischia, and of Thomas Lüthi, found on his website www.tiptom.ch/album/blumen/. The photograph of fireworks in Chantilly on page 214 is courtesy and copyright of Christophe Blanc and taken from his beautiful website at christopheblanc.free.fr. The figure of myosotis on page 230 is courtesy and copyright by Markku

Savela. The image of the wallpaper groups on page page 231 is copyright and courtesy of Dror Bar-Natan, and is taken from his fascinating website at www.math.toronto.edu/~drorbn/Gallery. The images of solid symmetries on page page 232 is copyright and courtesy of Jonathan Goss, and is taken from his website at www.phys.ncl.ac.uk/staff.njpg/symmetry. Also David Mermin and Neil Ashcroft have given their blessing to the use. The drawings of a ringing bell on page 252 are courtesy and copyright of H. Spiess. The interference figures on page 260 are copyright and courtesy of Rüdiger Paschotta and found on his free laser encyclopedia at www.rp-photonics.com. The image of the portable ultrasound machine on page 266 is courtesy and copyright General Electric. The ultrasound image on page 266 courtesy and copyright Wikimedia. The figure of the soliton in the water canal on page 269 is copyright and courtesy of Dugald Duncan and taken from his website on www.ma.hw.ac.uk/solitons/soliton1.html. The image of Rubik's cube on page 275 is courtesy of Wikimedia. The photographs of wakes on page 279 are courtesy Wikimedia and courtesy and copyright of Christopher Thorn. The fractal mountain on page 284 is courtesy and copyright by Paul Martz, who explains on his website www.gameprogrammer.com/fractal.html how to program such images. The soap bubble photograph on page page 291 is copyright and courtesy of LordV and found on his website www.flickr.com/photos/lordv. The photographs of silicon carbide on page 292 are copyright and courtesy of Dietmar Siche. The photograph of a single barium ion on page 293 is copyright and courtesy of Werner Neuhauser at the Universität Hamburg. The AFM image of silicon on page 293 is copyright of the Universität Augsburg and is used by kind permission of German Hammerl. The figure of helium atoms on metal on page 293 is copyright and courtesy of IBM. The photograph of an AFM on page 293 is copyright of Nanosurf (see www.nanosurf.ch) and used with kind permission of Robert Sum. The photograph of the tensegrity tower on page 296 is copyright and courtesy of Kenneth Snelson. The photograph of the Atomium on page 298 is courtesy and copyright by the Asbl Atomium Vzw and used with their permission, in cooperation with SABAM in Belgium. Both the picture and the Atomium itself are under copyright. The photographs of the granular jet on page 299 in sand are copyright and courtesy of Amy Shen, who discovered the phenomenon together with Sigurdur Thoroddsen. The photographs of the machines on page 299 are courtesy and copyright ASML and Voith. The photograph of the bucket-wheel excavator on page 300 is copyright and courtesy of RWE and can be found on their website www.rwe.com. The photographs of fluid motion on page 303 are copyright and courtesy of John Bush, Massachusetts Institute of Technology, and taken from his web site www-math.mit.edu/~bush. The photograph of the effects of air pressure on page 306 is courtesy and copyright of Kornelia Möller. The sailing boat on page 307 is courtesy and copyright of Bladerider International. The photograph of the smoke ring at Etna on page 315 is courtesy and copyright by Daniela Szczepanski and found at her extensive websites www.vulkanarchiv.de and www.vulkane.net. The thermographic images of a braking bicycle on page 320 are copyright Klaus-Peter Möllmann and Michael Vollmer, Fachhochschule Brandenburg/Germany, and courtesy of Michael Vollmer and Frank Pinno. The image of page 320 is courtesy and copyright of ISTA. The images of thermometers on page 323 are courtesy and copyright Wikimedia, Ron Marcus, Braun GmbH, Universum, Wikimedia and Thermodevices. The balloon photograph on page 325 is copyright Johan de Jong and courtesy of the Dutch Balloon Register found at www.dutchballoonregister.nl.ballonregister, nederlands The pollen image on page 327 is courtesy Wikimedia. The scanning tunnelling microscope picture of gold on page 335 is courtesy of Sylvie Rousset and copyright by CNRS in France. The photograph of the Ranque-Hilsch vortex tube is courtesy and copyright Coolquip. The photographs and figure on page 352 are copyright and courtesy of Ernesto Altshuler, Claro Noda and coworkers, and found on their website www.complexperiments.net. The road corrugation photo is courtesy of David Mays and taken from his paper Ref. 298. The oscillon picture on page 354 is courtesy and copyright by Paul Umbanhowar. The drawing of swirled spheres on page 354 is courtesy and copyright by Karsten

Kötter. The pendulum fractal on page 358 is courtesy and copyright by Paul Nylander and found on his website bugman123.com. The fluid flowing over an inclined plate on page 360 is courtesy and copyright by Vakhtang Putkaradze. The photograph of the Belousov-Zhabotinski reaction on page 361 is courtesy and copyright of Yamaguchi University and found on their picture gallery at www.sci.yamaguchi-u.ac.jp/sw/sw2006/. The photographs of starch columns on page 362 are copyright of Gerhard Müller (1940-2002), and are courtesy of Ingrid Hörnchen. The other photographs on the same page are courtesy and copyright of Raphael Kessler, from his websitewww. raphaelk.co.uk, of Bob Pohlad, from his websitewww.ferrum.edu/bpohlad, and of Cédric Hüsler. The photograph of a swarm of starlings on page 365 is copyright and courtesy of Andrea Cavagna and Physics Today. The photograph of sunbeams on page 407 is copyright and courtesy by Fritz Bieri and Heinz Rieder and found on their website www.beatenbergbilder.ch. The drawing on page 411 is courtesy and copyright of Daniel Hawkins. The photograph of a slide rule on page 415 is courtesy and copyright of Jörn Lütjens, and found on his website www.joernluetjens. de. The photograph of the bursting soap bubble on page 404 is copyright and courtesy by Peter Wienerroither and found on his website homepage.univie.ac.at/Peter.Wienerroither. The photograph on the back cover, of a basilisk running over water, is courtesy and copyright by the Belgian group TERRA vzw and found on their website www.terra.vzw.org. All drawings are copyright by Christoph Schiller. If you suspect that your copyright is not correctly given or obtained, this has not done on purpose; please contact me in this case.



$\frac{A}{A_{BE}}$

**
Aarts, D. 462
Abe, F. 449
Ackermann, Rudolph 412
Adachi, S. 462
Adelberger, E.G. 457
Adenauer, Konrad 179
Aetius 91, 452
Ahlgren, A. 448
Aigler, M. 56
AIP 149
Åkerman, N. 459
Alder, Ken 158, 457, 477
Alighieri, Dante 151
Alizad, A. 462
Alsdorf, D. 459
Altshuler, E. 468
Altshuler, Ernesto 352, 353,
478
Amundsen, Roald 151, 456
An, K. 464
Anders, S. 457
Anderson, R. 454
Andreotti, B. 469
Angell, C.A. 465
Aniçin, I.V. 458
Anonymous 43, 99, 211
Aquinas, Thomas 285, 446
Archimedes
life 92
Aristarchus of Samos 123, 135
453
Aristotle 47, 154, 450
life 39
Armstrong, J.T. 451
Aronson, Jeff K. 472, 474

Arseneau, Donald 474

Artigas, M. 463
Ashby, M.F. 464
Ashcroft, Neil 232, 478
Ashida, H. 445
ASML 299, 478
Asterix 295
Ata Masafumi 474
Atomium 298, 478
Au, W.W.L. 463
Audoin, C. 472
Augel, Barbara 474
Augel, Edgar 474
Ausloos, M. 463
Aussillous, P. 466
Avogadro, Amedeo
life 289
Axelrod, R. 445
Ayiomamitis, Anthony 67, 125,
135, 153, 169, 185, 430, 476
Azbel, M.Ya. 453
В
Babinet, Jacques
life 384
Babloyantz, A. 469
Bach, Michael 18, 475
Bachelier, Louis 467
Baez, John 474
Bagnoli, Franco 474
Bahr, D.B. 462
Baille, Jean-Baptistin 457
Balibar, S. 465
ballonregister, nederlands 478
Banach, Stefan 56, 284
life 55
Banavar, J.R. 461
Bandler, Richard 206, 446

Bar-Natan, Dror 231, 478
Barber, B.P. 452
Barberini, Francesco 285
Barberi Gnecco, Bruno 474
Barbosa, Lucas 134, 475
Barbour, Julian 449
Barone, Éric 182, 477
Bartel, Elmar 474
Bassler, K.E. 468
Batista-Leyva, A.J. 468
Batty, R.S. 463
Baudelaire, Charles
life 18
Bautista, Ferdinand 474
Beaty, William 474
Becker, J. 459
Beeksma, Herman 474
Behroozi, C.H. 448
Bekenstein, Jacob 346
Belfort, François 474
Bellard, Fabrice 470
Bellizzi, M.J. 456
Benka, S. 452
Bennet, C.L. 455
Bennet-Clark, H.C. 450
Bennett, C.H. 468
Bentley, W.A. 468
Benzenberg, Johann Friedrick
124
Berendsen, Marcel 86, 476
Bergquist, J. 472
Berman, D.H. 463
Bernabei, Tullio 476
Bernoulli, Daniel 325
life 304
Bernoulli, Johann 378, 433
Bessel 132

Bessel, Friedrich Wilhelm	Brennan, Richard 446	Cartesius
life 137	Brenner, M.P. 452	life 50
Bevis, M. 459	Bright, William 471	Casati, Roberto 452
Beyer, Lothar 474	Brillouin, L. 467	Cassius Dio 191
Bielefeldt, H. 464	Brillouin, Léon 333	Cauchy, Augustin-Louis 235
Bieri, Fritz 407, 479	Brooks, Mel 329	Cavagna, Andrea 365, 479
Biggar, Mark 474	Brouwer, Luitzen 49	Cavendish, H. 457
Bilger, H.R. 454	Brown, J.H. 461	Cavendish, Henry
BIPM 92	Brown, Robert 326	life 158
Bischoff, Bernard 472	Browning, R.C. 462	Caveney, S. 450
Bizouard, C. 455	Bruce, Tom 474	Caveney, Stanley 75, 76, 476
Bladerider International 307,	Brush, S. 455	Cayley, Arthur 235
478	Buchmann, Alfons 474	Celsius, Anders
Blagden, Charles 341	Bucka, H. 454	life 341
Blanc, Christophe 214, 477	Bucka, Hans 131	Chaineux, J. 472
Blumensath, Achim 474	Buckley, Michael 445	Chan, M.A. 454, 455
Bocquet, Lydéric 438, 465	Budney, Ryan 474	Chandler, Seth 133
Bode, Johann Elert 190	Bújovtsev, B.B. 445	Chao, B.F. 455
Boethius 402	Bundesamt für Kartographie	Charlemagne 379
Bogusch, Stephan 476	und Geodäsie 130, 477	Childs, J.J. 464
Bohr, J. 449	Bunn, E.F. 470	Chomsky, Noam 448
Bohr, Jakob 69, 476	Bunyan, Paul 464	Christlieb, N. 458
Bohr, Niels 333	Buridan, Jean 154, 457	Christou, A.A. 458
Boiti, M. 280, 463	Burša, Milan 450	Chudnovsky, D.V. 470
Bolt, Usain 75	Busche, Detlev 448	Chudnovsky, G.V. 470
Boltzmann, L. 468	Bush, J.W.M. 465	Clancy, Tom 462
Boltzmann, Ludwig 219, 337	Bush, John 303, 478	Clanet, Christophe 438
life 327	Böhncke, Klaus 474	Clarke, S. 447
Bombelli, Luca 474		Clausius, Rudolph 328
Boone, Roggie 474	С	life 330
Borel, Emile 340, 468	Caesar, Gaius Julius 313	Clavius, Christophonius
Borelli, G. 446	Caesar, Gaius Julius 379	life 62
Born, Max 286	Caizergues, Alex 35	Cleobulus
Bosch 83, 476	Cajori, Florian 471	life 51
Bourbaki, Nicolas 378	Caldwell, R. 451	CNAM 129, 477
Bowden, F.P. 460	Caldwell, R.L. 451	CNES 193
Bower, B. 461	Calzadilla, A. 460	CNRS 335, 478
Brackenbury, J. 451	Campbell, D. 468	Cohen-Tannoudji, G. 467
Brackenbury, John 86, 476	Campbell, D.K. 469	Cohen-Tannoudji, Gilles 333
Bradley, James 137	Caps, H. 466	Colazingari, Elena 474
life 137	Carl Zeiss 130, 477	Coleman-McGhee, Jerdome
Braginsky, V.B. 458	Carlip, Steve 474	465
Brahe, Tycho 152	Carlyle, Thomas	Colladon, Daniel 312
life 151	life 113	Collins, J.J. 446
Brahm, A. de	Carnot, Sadi	Compton, A.H. 454
life 379	life 104	Compton, Arthur 130
Brandes, John 474	Carr, Jim 474	Conkling, J.A. 450
	Carroll, Lewis, or Charles	Conservatoire National des
Brantjes, R. 429	Carron, Lewis, of Charles	Consei vatorie mational des

Lutwidge Dogson 337

BESSEL

Brebner, Douglas 474

Motion Mountain – The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christoph Schiller November 1997–March 2012

Arts et Métiers 477

Conti, Andrea 474	Diel, Helmut 476	Elswijk, H.B. 464
Coolquip 345, 478	Diel, Isolde 476	Elswijk, Herman B. 474
Cooper, Heather 456	Diez, Ulrich 474	Elsässer, Martin 167, 475
Cooper, John N. 445	DiFilippo, Frank 474	EMBO 46
Coriolis, Gustave-Gaspard	Dill, L.M. 463	Emelin, Sergei 474
124	Dio, Cassius 459	Emerson, Ralph Waldo 271
life 101	Dirac, Paul 247, 378	Emsley, John 468
Cornell, E.A. 448	Dirr, Ulrich 475	Engelmann, Wolfgang 448
Cornu, Marie-Alfred 457	Dixon, Bernard 449	Engels, Friedrich 204
Corovic, Dejan 474	Di Iorio, Giorgio 211, 477	Engemann, S. 464
Costabel, Pierre 285	Dobra, Ciprian 474	Enquist, B.J. 461
Coyne, G. 463	Doorne, C.W.H. van 469	Enss, Christian 466
Coyne, George 463	Dorbolo, S. 466	Eötvös, R. von 458
Craven, J.D. 458	Dorbolo, Stéphane 314	Erdős, Paul 56
Crespi, Roberto 474	Douady, S. 468	Eriksen, H.K. 450
Cross, R. 416	Dougherty, R. 449	Erwin Sattler OHG 155, 477
Crutchfield, J. 468	Doyle, Arthur Conan 283	Euclid, or Eukleides 34
Crystal, David 471	Dragon, Norbert 474	Euler, Leonhard 133, 192, 378,
, , , ,	Drake, Stillman 450	453
D	Dreybrodt, W. 469	life 196
Dahlman, John 474	Dubelaar, N. 429	European Molecular Biology
Danecek, Petr 474	Dubrulle, B. 459	Organisation 476
Dante Alighieri 151	Dumont, Jean-Paul 445, 452,	Evans, J. 459
Darius, J. 452	470	Everitt, C.W. 473
Darley, Vincent 474	Duncan, Dugald 269, 478	Eöt-Wash Group 160, 477
Darre, Daniel 474	Durben, D.J. 465	Eötvös, Roland von 128
Dasari, R.R. 464	Durieux, M. 466	
Davidhazy, Andrew 39, 278,	Dusenbery, David 451	F
303, 476	Dutton, Z. 448	Fabian, Werner 475
Davidovich, Lev 381	Dyson, Freeman 474	Fairbanks, J.D. 473
Davis, S.D. 464	7	Faisst, H. 469
Deaver, B.S. 473	E	Falk, G. 458
Deconinck, Bernard 281	Earls, K. 451	Falkovich, G. 465
Dehant, V. 455	Earman, John 449	Fantoli, Annibale 463
Dehmelt, H. 464	Eckhardt, B. 469	Farinati, Claudio 474
Dehmelt, Hans 464	Eddington, Arthur	Farmer, J. 468
Dehn, Max	life 98	Farrant, Penelope 452
life 56	Edelmann, H. 458	Fatemi, M. 462
Demko, T.M. 454	Edens, Harald 475	Fatio de Duillier, Nicolas 189
Democritus 288, 381	Edwards, B.F. 421	Fatio de Duillier, Nicolas 459
life 288	effect, Kaye 316	Faybishenko, B.A. 469
Descartes, René 378, 414	Einstein, A. 467	Fayeulle, Serge 460
life 50	Einstein, Albert 17, 172, 176,	Feder, T. 470
Destexhe, A. 469	177, 220, 239, 241, 247, 327	Fehr, C. 460
DeTemple, D.W. 470	Ekman, Walfrid	Fekete, E. 458
Deutschmann, Matthias 402	life 453	Feld, M.S. 464
Dewdney, Alexander 69	Ellerbreok, L. 463	Fermani, Antonio 474
Dicke, R.H. 458	Els, Danie 474	Fernandez-Nieves, A 456
	12.1	

Elsevier, Louis 406

Fibonacci, Leonardo 381

Conti

Diehl, Helmut 59

life 376	GeoForschungsZentrum	Hain, Tim 476
Figueroa, D. 460	Potsdam 162, 477	Halberg, F. 448
Filippov, T. 462	Georgi, Renate 474	Haley, Stephen 474
Fink, Hans-Werner 464	Ghahramani, Z. 460	Halley, Edmund 169
Finkenzeller, Klaus 474	Ghavimi-Alagha, B. 456	Halliday, David 466
Firpić, D.K. 458	Giessibl, F.J. 464	Halliwell, J.J. 448
Fischer, Ulrike 474	Gilligan, James 460	Hamilton, William 378
Flach, S. 469	Glassey, Olivier 474	Hammerl, German 478
Flachsel, Erwein 445	Gold, Tommy 261	Hancock, M.J. 465
Flindt, Rainer 448	Goldrich, P. 437	Hardcastle, Martin 474
Fokas, A.S. 280, 463	Goldstein, Herbert 460	Hardy, Godfrey H.
Foreman, M. 449	Goles, E. 469	life 166
Forsberg, B. 459	Golubitsky, M. 446	Hare, Trent 477
Fortes, L.P. 459	González, Antonio 474	Harriot, Thomas 378
Foster, James 460	Gooch, Van 45	Harris, S.E. 448
Foucault, Jean Bernard Léon	Gordillo, J.M. 465	Hartman, W.K. 457
life 127	Goriely, A. 465	Hasha, A.E. 465
Franklin, Benjamin 284	Goss, Jonathan 232, 478	Hausherr, Tilman 474
Frenzel, H. 89	Gracovetsky, Serge 112, 453	Hawkins, Daniel 63, 410, 411,
Fresnel, Augustin 261	Grahn, Alexander 475	479
Friedman, David 195	Graner, F. 459	Hayes, Allan 474
Fritsch, G. 463	Graw, KU. 436	Hays, J.D. 455
Frova, Andrea 450	Gray, C.G. 422, 460, 461	Heath, Thomas 453
Full, R. 451	Greenleaf, J.F. 462	Heber, U. 458
Full, Robert 86, 476	Greenside, Henry 400	Heckel, B.R. 457
Fulle, Marco 22, 475	Gregorio, Enrico 475	Heckman, G. 457
Fumagalli, Giuseppe 452, 460	Greier, Hans-Christian 190,	Hediger, Heini 450
Fundamentalstation Wettzell	475	Heisenberg, Werner 229, 305,
477	Greiner, Jochen 474	333, 461
Furrie, Pat 474	Grimaldi, Francesco 273	Helden, A. van 456
	Grinder, John 206, 446	Heller, Carlo 43
G	Groenendijk, Max 39, 476	Heller, E. 462
Galilei, Galileo 42, 66, 72, 73,	Gross, R.S. 455	Hellinger, Bert 460
124, 141, 157, 178, 181, 182,	Grossmann, A. 463	Helmholtz, Hermann von 104
287, 288, 406, 417	Grünbaum, Branko 461	life 321
life 33, 285	Guglielmini,	Helmond, Tom 474
Galileo 123, 381	Giovanni Battista 124	Helmont, Johan Baptista van
Galois, Evariste 235	Guillén, Antonio 86	325
Gardner, Martin 414	Gustav Jacobi, Carl 378	Hembacher, S. 464
Garrett, A. 452	Gutierrez, G. 460	Henbest, Nigel 456
Gaspard, Pierre 329, 467	Gácsi, Zoltán 474	Henderson, D.W. 460
Gastaldi, Luca 20, 55, 179, 475	Günther, B. 448	Henderson, Paula 474
Gauthier, N. 461		Henon, M. 459
Gekle, S. 465	Н	Henson, Matthew 426
Gelb, M. 446	Haandel, M. van 457	Hentig, Hartmut von 7
Gemini Observatory/AURA	Haber, John 474	Heraclitus
147	Hagen, J.G. 454	life 17
General Electric 266, 478	Hagen, John 128	Heraclitus of Ephesus 26, 230
Geng, Tao 227	Haigneré, Jean-Pierre 193	Herbert, Wally 456

FIGUEROA

Geng, Tao 227

Hermann, Jakob 427
Heron of Alexandria 311
Herrmann, F. 467
Herrmann, Friedrich 444
Herschel, William 140
Hertz, H.G. 452
Hertz, Heinrich 219
Hertz, Heinrich Rudolf
life 202
Hertzlinger, Joseph 474
Hestenes, D. 459
Heumann, John 474
Hewett, J.A. 457
Hietarinta, J. 463
Hietarinta, Jarmo 270, 280,
475
Higashi, R. 472
Hilbert, David 220, 242
Hilgenfeldt, S. 452
Hillman, Chris 474
Hipparchos 132
Hirano, M. 459
Hirota, R. 280, 463
Hodges, L. 458
Hodgkin, A.L. 268, 462
Hoeher, Sebastian 465
Hof, B. 469
Hohenstatt, M. 464
Holbrook, N.M. 464
Hollander, E. 466
Hong, FL. 472
Hooke, Robert 166, 248
life 152
Hoornaert, Jean-Marie 120,
477
Horace, in full Quintus
Horatius Flaccus 203
Hosoi, A.E. 465
Hoyle, R. 410
Huber, Daniel 474
Hudson, R.P. 466
Humphreys, R. 455
Humphreys, W.J. 468
Hunklinger, Siegfried 466
Hunter, D.J. 451
Huxley, A.F. 268, 462
Huygens, Christiaan 154, 155,
261

life 94

HERMANN

Hörnchen, Ingrid 479
Hüsler, Cédric 362, 479
IBM 293
frah, Georges 471
Illich, Ivan 399
life 398
Imae, Y. 451
Imbrie, J. 455
Ingenhousz, Jan 326
Ingram, Jay 463
INMS 43
Inquisition 285
International Tennis
Federation 20, 475
IPCC 439
ISTA 320, 478
SVR, University of
Southhampton 257, 475
Ivanov, Igor 474
Ivry, R.B. 448
I
acobi, Carl 219
aeger-LeCoultre 119, 477
ahn, O. 461
alink, Kim 474
ames, M.C. 458
famil, M. 474
ammer, M. 456
anek, Jürgen 474
apan Aerospace Exploration
Agency 477
AXA 129, 477
effreys, Harold 131
en, E. 468
ennings, G.M. 464
ensen, Hans 471
esus 313
ob, Georg 444
ohansson, Mikael 474
ohn Paul II 286
ohnson, Ben 450
ohnson, Michael 75
ohnson, Samuel 372, 398
ohnston, K.J. 451
ones, Gareth 468
ones, Quentin David 474

Jones, 10ny 472
Jones, William 378
Jong, Johan de 325, 478
Jong, Marc de 474
Joule, James 321
Joule, James P. 104
Joule, James Prescott
life 321
Joutel, F. 458
Jouzel, J. 455
Jouzel, Jean 139, 477
Jürgens, H. 357
Jürgens, Hartmut 449
K

, 0
K
Köppe, Thomas 474
Kadanoff, L.P. 469
Kamiya, K. 433
Kanada Yasumasa 470
Kanecko, R. 459
Kant, Immanuel 177
Kantor, Yacov 400
Karl, G. 422, 461
Karstädt, D. 466
Katori, H. 472
Kawagishi, I. 451
Keath, Ed 121, 477
Keesing, R.G. 447, 472
Kelu, Jonatan 474
Kemp, David 261
Kempf, Jürgen 448
Kenderdine, M.A. 452
Kendrick, E. 459
Kepler, Johannes 164, 166
life 151
Kerswell, R. 469
Kessler, Raphael 362, 479
Keyence 52, 476
Khallikan, Ibn
life 405
Kirby, Jack 477
Kirchhoff, Gustav 262
Kiss, Joseph 474
Kistler 313
Kistler, S.F. 465
Kitaoka Akiyoshi 16, 18, 445
475
Kitaoka, A. 445
Kitaoka, Akiyoshi 445

Kivshar, Y.S. 469	Lang, H. de 468	Liu, C. 448
Klaus Tschira Foundation 475	Lang, K.R. 84	Llobera, M. 453
Kluegel, T. 454	Langangen, Ø. 450	Lloyd, Seth 467
Klügel, T. 454	Laplace, Pierre Simon	Lockwood, E.H. 461
Klügel, Thomas 477	life 171	Lodge, Oliver
Koblenz, Fachhochschule 34,	Larsen, J. 455	life 130
475	Laskar, J. 457, 459	Lohse, D. 452, 465
Koch, G.W. 464	Laskar, Jacques 190	Lohse, Detlev 89, 476
Kooijman, J.D.G. 418	Laveder, Laurent 193, 477	Lombardi, Luciano 474
Koolen, Anna 474	Lavenda, B. 468	LordV 291, 478
Koomans, A.A. 464	Lavoisier, Antoine-Laurent	Loschmidt, Joseph
Korteweg, Diederik 270	life 93	life 290
Kramp, Christian 378	Laws, Kenneth 469	Lucretius 288
Krampf, Robert 400	Lecar, M. 459	life 39
Krehl, P. 464	Lee, C.W. 470	Lucretius Carus, Titus 39, 288
Krehl, Peter 297	Lee, S.J. 456	Luke, Lucky 295
Krijn, Marcel 474	Lega, Joceline 469	Lévy-Leblond, Jean-Marc 111,
Kristiansen, J.R. 450	Legendre, Adrien-Marie 378	189, 444
Krívchenkov, V.D. 445	Lehmann, Paul 379, 472	Lüders, Klaus 444
Krotkow, R. 458	Leibniz, Gottfried Wilhelm	Lüthi, Thomas 211, 477
Kruskal, M.D. 462	101, 166, 213, 217, 220, 378,	Lütjens, Jörn 415, 479
Kruskal, Martin 271	447	240,0110, 70111 425, 475
Królikowski, Jarosław 474	life 79	M
Kubala, Adrian 474	Leica Geosystems 52, 476	MacDonald, G.J.F. 454
Kudo, S. 451	Leidenfrost, Johann Gottlieb	MacDougall, Duncan 453
Kumar, K.V. 450	313	MacDougalls, Duncan 114
Kurths, Jürgen 457	Lekkerkerker, H. 462	Mach, Ernst
Kuzin, Pavel 474	Lennard, J. 379	life 95
Kvale, E.P. 454	Lennerz, C. 422	Macmillan, R.H. 461
Kötter, K. 469	Leon, JP. 280, 463	Maekawa, Y. 451
Kötter, Karsten 354, 355, 478	Leonardo of Pisa	Magariyama, Y. 451
Küster, Johannes 475	life 376	Magono, C. 470
Küstner, Friedrich 455	Lesage, GL. 459	Mahajan, S. 437
life 132	Lesage, Georges-Louis 189	Mahoney, Alan 474
ine 132	Leucippus of Elea	Mahrl, M. 469
L	life 288	Maiorca, Enzo 465
La Caille 153	Leutwyler, K. 462	Malin, D. 84
Lagrange, Joseph Louis 225	Lewallen, Charles 149, 477	Mallat, Stéphane 463
Lagrange, Joseph Louis 225 Lagrange, Joseph Louis 216	Le Verrier, Urbain 172	Malley, R. 459
Lagrangia, Giseppe 378	Libbrecht, Kenneth 355, 363,	Mandelbrot, Benoît 54, 283
Lagrangia, disceppe 3/6	364, 475	Mandelstam, Stanley 460
Giuseppe Lodovico	Lichtenberg, Georg Christoph	Mannhart, J. 464
life 216	life 33	Manogg, P. 463
Lalande 153	Lim Tee Tai 315, 316, 475	
		Marcus, Richard 446
Lambeck, K. 454	Lim, T.T. 466	Marcus, Ron 323, 478
Lambert, S.B. 455	Linde, Johan 474	Marcy, Geoffrey 170, 477
Lancaster, George 460	Lintel, Harald van 474	Maris, H. 465
Landau, Lev 381	Lissauer, J.J. 470	Mark, Martin van der 342, 474
Landauer, R. 468	Lith-van Dis, J. van 467	Markus, M. 469

Motion Mountain - The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright © Christoph Schiller November 1997–March 2012

Kivshar

Marsaglia, G. 460
Martin, A. 461
Martina, L. 280, 463
Martos, Antonio 474
Martz, Paul 284, 478
Martínez, E. 468
Marx, Groucho 373
Masudi, Ibn 405
Matthews, R. 458
Maupertuis, Pierre Louis
Moreau de
life 123
Mayer, Julius Robert 104
life 321
Maynard, Finian 35
Mayné, Fernand 474
Mayr, Peter 474
Mays, D.C. 469
Mays, David 353, 478
McCoy, Ray 113, 477
McElwaine, J.N. 469
McLaughlin, William 445
McLean, H. 450
McMahob, T.A. 464
McMillen, T. 465
McNaught, Robert 88, 476
McQuarry, George 474
Medenbach, Olaf 449
Medien Werkstatt 476
Meijaard, J.P. 418
Melo, F. 469
Mendeleyev,
Dmitriy Ivanovich 99
Mendoza, E. 469
Merckx, Eddy 105
Mermin, David 232, 478
Merrit, John 474
Mertens, K. 469
Mettler-Toledo 99, 477
Mettrie, J. Offrey de la 359
Miákishev, G.Ya. 445
Michaelson, P.F. 473
Michel, Stanislav 101
Michelangelo 381
Michelberger, Jürgen 84, 476
Michell, John
life 158
Micro Crystal 475

Microcrystal 251

Mikkola, Seppo 458
Milankovitch, Milutin
life 137
Mineyev, A. 464
Minnaert, Marcel G.J. 90
Mirabel, I.F. 448
Mirsky, S. 416
Mohazzabi, P. 458
Mole, R. 462
Monitz, E.J. 452
Moore, J.A. 461
Moortel, Dirk Van de 474
Moreau de Maupertuis,
Pierre Louis
life 123
Moreira, N. 461
Morgado, E. 448
Morgado, Enrique 46, 476
Morlet, J. 463
Morris, S.W. 469
Moscow State Circus 23, 475
Moser, Lukas Fabian 474
Mozurkewich, D. 451
Muller, R.A. 462
Mulligan, J.F. 452
Munk, W.H. 454
Muramoto, K. 451
Murata, Y. 459
Murdock, Ron 474
Mureika, J.R. 460
Murillo, Nadia 474
Murphy, Robert 378
Murray, C.D. 458
Muskens, O. 462
Muynck, Wim de 474
Måløy, K.J. 468
Möller, Kornelia 306, 478
Möllmann, KP. 466
Möllmann, Klaus-Peter 320,
478
17
Müller, Frank 54, 476
Müller, G. 469
Müller, Gerhard 361, 362
life 479
N
Namba, K. 451
Namouni, F. 458

Namouni, Fathi 474

Nanosurf 293
Nansen, Fridtjof 453
Napiwotzki, R. 458
Naples Zoo 52
Naples zoo 476
Napoleon 172
NASA 43, 99, 439
Nassau, K. 90
Nauenberg, M. 456
Navier, Claude
life 306
Needham, Joseph 460
Nelson, A.E. 457
Neuhauser, W. 464
Neuhauser, Werner 293, 478
Neumaier, Heinrich 474
Neumann, Dietrich 168, 457
Newcomb, Simon 172, 458
Newton, Isaac 47, 124, 153, 160
388, 447
life 33
Nicolis, Grégoire 467
Niederer, U. 241, 461
Niepraschk, Rolf 475
Nieuwpoort, Frans van 474
Nieuwstadt, F.T.M. 469
Nieves, F.J. de las 456
Nightingale, J.D. 460
Nitsch, Herbert 308
Nobel, P. 464
Noda, Claro 478
Noether, Emmy
•
life 241 Nolte, John 446
Nonnius, Peter
life 62
Norfleet, W.T. 450
Northwestern University 49,
83, 276, 476
Novikov, V.A. 422, 461
Nuñes, Pedro
life 62
Nylander, Paul 358, 479
O'Keefe R 456
LIBERIE K AFD

Oberdiek, Heiko 474 Oberquell, Brian 474 Offner, Carl 474

Marsaglia

Offrey de la Mettrie, J. 359	Piccard, Auguste	R
Ogburn, P.L. 462	life 312	Rahtz, Sebastian 474
Olsen, K. 449	Pieranski, P. 450	Ramanujan, Srinivasa 166
Oostrum, Piet van 474	Pieranski, Piotr 70, 476	Ramos, O. 468
Oppenheimer, Robert 222	Pietrzik, Stefan 430, 476	Randi, James 22
Origone, Simone 182, 477	Pikler, Emmi 446	Rankl, Wolfgang 474
Oughtred, William 378	Pikovsky, Arkady 457	Rawlins, D. 456
3,1	Pinker, Steven 448	Raymond, David 401
P	Pinno, F. 466	Rechenberg, Ingo 451
Page, Don 474	Pinno, Frank 478	Recorde, Robert 378
Page, Janet 475	PixHeaven.net 193	life 377
Pahaut, Serge 474	Planck, Max 327, 332	Redondi, Pietro 33, 285, 474
Pais, A. 461	Plato 366	Rees, W.G. 444
Panov, V.I. 458	life 288	Reichl, Linda 466, 467
Papadopoulos, J.M. 418	Plinius, in full Gaius Plinius	Reimers, D. 458
Pappus 92	Secundus 444	Renselle, Doug 474
Park, David 444	Plisson, Philip 23, 475	Reppisch, Michael 474
Parks, David 86, 474, 476	Plutarchus 208	Resnick, Robert 466
Parlett, Beresford 24	Pohl, Robert 444	Richardson, Lewis Fray
Parmenides of Elea 17, 80	Pohl, Robert O. 444	life 53
Pascazio, Saverio 474	Pohlad, Bob 362, 479	Rickwood, P.C. 449
Paschotta, Rüdiger 260, 478	Poincaré, Henri 140	Rieder, Heinz 407, 479
Pasi, Enrico 474	Poinsot, Louis 128	Rieflin, E. 451
Paul, T. 463	Poisson, Siméon-Denis	Rieker Electronics 83, 476
Pauli, Wolfgang 427	life 162	Rinaldo, A. 461
Peano, Giuseppe 378	Polster, B. 468	Rindt, Jochen
Peary, Robert 151, 426, 456	Pompeius, Gnaeus 208	life 34
Pěč, Karel 450	Preston-Thomas, H. 466	Rivas, Martin 474
Pedley, T.J. 450	Price, R.H. 98	Rivera, A. 468
Peeters, Bert 474	Prigogine, Ilya 319, 361	Robertson, Will 475
Peitgen, HO. 357	Pritchard, Carol 474	Robutel, P. 458
Peitgen, Heinz-Otto 449	Proença, Nuno 474	Rodgers, Tony 25, 475
Pekár, V. 458	Protagoras 387	Rodin, Auguste 154
Pelizzari, Umberto 465	Przybyl, S. 450	Rodríguez, L.F. 448
Pempinelli, F. 280, 463	Ptolemy 66	Rohrbach, Klaus 455
Perc, M. 453	Purves, William 474	Roll, P.G. 458
Perelman, Yakov 275, 444	Putkaradze, V. 469	Romer, R.H. 431
Perini, Romano 474	Putkaradze, Vakhtang 360,	Rosek, Mark 477
Perrin, J. 467	479	Rosenblum, Michael 457
Perrin, Jean 440, 467	Putterman, S.J. 452	Rothacher, M. 454
life 328	Pythagoras 271	Rousset, Sylvie 478
Peters, I.R. 465	Päsler, Max 460	Ruben, Gary 474
Peterson, Peggy 68, 476	Pérez-Mercader, J. 448	Rudolff, Christoff 378, 471
Petit, Jean-Pierre 402	Pérez-Penichet, C. 468	Ruga, Spurius Carvilius 373
Pfeffer, W.T. 462		Ruina, A. 418
Phillips, R.J. 457	Q	Ruppel, W. 458
Phinney, S. 437	Quinlan, G.D. 458	Rusby, R.L. 466, 468

Quéré, D. 466

Rusby, Richard 468

Russel, J.S. 462

Piaget, Jean 446

Physics Today 365, 479

Russel, Mark 193	Schröder, Ernst 378	Sitti, M. 456
Russell, Bertrand 218	Schultes, H. 89	Sitti, Metin 149
Russell, John Scott	Schwab, A.L. 418	Slabber, André 474
life 268	Schwab, Arend 418	Sluckin, T.J. 453
Russo, Lucio 444	Schwartz, Richard 446	Smith, George 472
Rutherford, Ernest 24	Schwenk, K. 468	Smoluchowski, Marian von
RWE 300, 478	Schwenkel, D. 464	327
Régnier, Gilles 173, 477	Schönenberger, C. 464	Snelson, Kenneth 296, 478
	Schörner, E. 422	Socrates 366
S	Science 139	Sokal, A.D. 421
S.R. Madhu Rao 474	Science Photo Library 86	Soldner, J. 458
SABAM 298, 478	Science/AAAS 477	Soldner, Johann 177
Sade, Donatien de 202	Scime, E.E. 421	Solomatin, Vitaliy 474
Saghian, Damoon 474	Scott, Jonathan 474	Sonett, C.P. 454
Sagnac, Georges	Scriven 313	Song, Y.S. 456
life 130	Scriven, L.E. 465	Sotolongo-Costa, O. 468
Sanctorius	Segrè, Gino 466	Speleoresearch & Films/La
life 94	Settles, Gary 278	Venta 57, 476
Sands, Matthew 444	Sexl, Roman	Spence, J.C.H. 464
Santini, P.M. 280, 463		Spencer, R. 448
Santorio Santorio	life 58	-
	Shackleton, N.J. 455	Spiderman 295
life 94	Shakespeare, William 22	Spiess, H. 252, 478
Saráeva, I.M. 445	Shalyt-Margolin, A.E. 467	Spinelli, Matthew 84, 476
Sattler OHG, Erwin 477	Shapiro, A.H. 454	Spiropulu, M. 457
Sattler-Rick, Stephanie 477	Shapiro, Asher 127	Sreedhar, V.V. 461
Sauer, J. 422	Sheldon, Eric 474	Sreenivasan, K.P. 465
Sauerbruch, Ferdinand 438	Shen, A.Q. 465	Stalla, Wolfgang 294
Saupe, D. 357	Shen, Amy 298, 299, 478	Stasiak, A. 450
Saupe, Dietmar 449	Sheng, Z. 465	Stedman, G.E. 454
Savela, Markku 230, 477	Shephard, G.C. 461	Steiner, Kurt 465
Scagell, Robin 183, 477	Shinjo, K. 459	Stengel, Ingrid 448
Schiller, Britta 474, 475	Shirham, King 30	Stephenson, Richard 455
Schiller, C. 464	Short, J. 472	Sternlight, D.B. 456
Schiller, Christoph 236, 479	Siart, Uwe 474	Steur, P.P.M. 466
Schiller, Isabella 474	Siche, Dietmar 292, 478	Stewart, I. 446
Schiller, Peter 474	Sierra, Bert 474	Stewart, Ian 56
Schiller, Stephan 474	Sigwarth, J.B. 458	Stieber, Ralph 477
Schlichting, HJ. 425	Sillett, S.C. 464	Stokes, Georges Gabriel
Schlüter, W. 454	Silva 34, 475	life 306
Schmidt, Klaus T. 451	Silverman, M. 451	Stong, C.L. 451
Schmidt, M. 462	Simmons, J.A. 463	Story, Don 474
Schmidt, T. 469	Simon, Julia 474	Stroock, A.D. 464
Schmidt-Nielsen, K. 468	Simon, M.I. 451	Strunk, William 472
Schneider, Jean 461	Simonson, A.J. 431	Su, Y. 458
Schneider, M. 454	Simoson, A.J. 459	Suchocki, John 444
Schneider, Wolf 472	Simplicius 367	Sugiyama, S. 451
Schooley, J.F. 466	Singh, C. 452	Suhr, S.H. 456
Schreiber, K.U. 454	Singleton, Douglas 474	Sum, Robert 478

Sissa ben Dahir 30

Russel

Schreiber, U. 454

Surdin, Vladimir 474

Surry, D. 450	Tonzig, Giovanni 445	Volin, Leo 474
Sussman, G.J. 459	Topper, D. 457	Vollmer, M. 466
Swackhamer, G. 459	Torge, Wolfgang 450	Vollmer, Michael 320, 478
Swatch Group 475	Toschek, P.E. 464	Voltaire 192, 388
Sweetser, Frank 474	Townsend, Paul 474	life 220
Swenson, C.A. 466	Tracer 34, 475	von Baer, Karl Ernst 143
Swift, G. 468	Trefethen, L.M. 454	Vorobieff, P. 469
Swinney, H.L. 469	Tregubovich, A.Ya. 467	Voss, Herbert 474
Swope, Martha 469	Trevorrow, Andrew 474	Vries, Gustav de 270
Szczepanski, Daniela 315, 478	Truesdell, C. 466	
Szilard, L. 467	Truesdell, Clifford 33	W
Szilard, Leo 333	Tsang, W.W. 460	Wagon, Stan 56, 449
	Tschichold, J. 471	Wah 170
Т	Tschira, Klaus 475	Wald, George 334, 468
Taberlet, N. 469	Tsuboi, Chuji 450	Waleffe, F. 469
Tabor, D. 460	Tsukanov, Alexander 23, 475	Walgraef, Daniel 469
Taimina, D. 460	Tucholsky, Kurt 233	Walker, J. 466
Tait, Peter 378	Tuijn, C. 416	Walker, Jearl 313, 445, 466
Takamoto, M. 472	Tuppen, Lawrence 474	Walker, John 159, 477
Talleyrand 157	Turner, M.S. 455	Wallis, John 378
Tamman, Gustav 294	Twain, Mark 112	Walter, Henrik 460
Tarko, Vlad 474		Ward, R.S. 462
Tarski, Alfred 284	U	Warkentin, John 474
life 56	Ucke, C. 425	Weber, R.L. 469
Tartaglia, Niccolò 381	Uffink, J. 467	Webster, Hutton 472
life 31	Uguzzoni, Arnaldo 474	Wedin, H. 469
Taylor, E.F. 460	Ulam, Stanislaw 348	Wegener, Alfred 133, 455
Taylor, G.J. 457	Umbanhowar, P.B. 469	Wehner, R. 456
Tegelaar, Paul 474	Umbanhowar, Paul 353, 354,	Wehus, I.K. 450
Telegdi, V.L. 468	478	Weierstall, U. 464
Tennekes, Henk 37, 476	Universität Augsburg 293	Weierstrass, Karl 378
TERRA 149	Upright, Craig 474	Weil, André 378
Tetzlaff, Tim 476		Weiss, M. 450
Tezel, Tunc 152, 477	V	Weiss, Martha 474
Thaler, Jon 474	Vandewalle, N. 466	Wells, M. 459
Thies, Ingo 474	Vanier, J. 472	Weltner, K. 465
Thinktank Trust 99, 477	Vannoni, Paul 474	Weninger, K.R. 452
Thomas Aquinas 285, 446	Vareschi, G. 433	West, G.B. 461
Thompson, Dave 475	vector, Runge-Lenz 427	Westerweel, J. 469
Thomson, W. 465	Velikoseltsev, A. 454	Weyand, P.G. 456
Thomson-Kelvin 104	Verne, Jules 431	Weyl, Hermann
Thomson-Kelvin, W. 298	Vernier, Pierre	life 48
Thomson-Kelvin, William	life 62	Wheeler, John 403
life 321	Vincent, D.E. 457	Wheeler, T.D. 464
Thorn, Christopher 279, 478	Virgo, S.E. 464	White, E.B. 472
Thoroddsen, S.T. 465	Vitali, Giuseppe 54	White, M. 455
Thoroddsen, Sigurdur 298,	Viviani, Vincenzo 128	Whitney, C.A. 84
478	Vogel, Steven 469	Widmann, Johannes 378
Titius, Johann Daniel 190	Voith 299, 478	Widmer-Schnidrig, R. 461

	Wierzbicka, Anna 448, 474
	Wijk, Mike van 474
	Wikimedia 34, 43, 118, 252,
	266, 275, 279, 327, 475, 478
	Wilder, J.W. 421
	Wiley, Jack 465
	Wilk, Harry 449
	Wilkie, Bernard 452
7	Williams, G.E. 455
	Wilson, B. 463
	Wiltshire, D.L. 454
_	Wisdom, J. 459
NERROITHER	Wise, N.W. 473
	Wittgenstein, Ludwig 16, 24,
	26, 47, 82, 142, 204, 336,
	369
	Wittlinger, M. 456

Wienerroither, Peter 404, 479

Wierda, Gerben 474

Wittlinger, Matthias 150
Wolf, G.H. 465
Wolf, H. 456
Wolfsried, Stephan 57, 476
Wolpert, Lewis 446
Wong, M. 451
Wright, Joseph 475
Wright, S. 456
Wulfila 375
Υ
Y Yamafuji, K. 465
•
Yamafuji, K. 465
Yamafuji, K. 465 Yamaguchi University 361,
Yamafuji, K. 465 Yamaguchi University 361, 479
Yamafuji, K. 465 Yamaguchi University 361, 479 Yatsenko, Dimitri 410
Yamafuji, K. 465 Yamaguchi University 361, 479 Yatsenko, Dimitri 410 Young, Andrew 474
Yamafuji, K. 465 Yamaguchi University 361, 479 Yatsenko, Dimitri 410 Young, Andrew 474 Young, Thomas 102

4
Zabusky, N.J. 462
Zabusky, Norman 271
Zaccone, Rick 474
Zakharian, A. 454
Zalm, Peer 474
Zanker, J. 447
Zedler, Michael 474
Zenkert, Arnold 448
Zeno of Elea 15, 17, 63, 80
Zheng, Q. 465
Ziegler, G.M. 56
Zimmermann, H.W. 467
Zimmermann, Herbert 333
Zweck, Josef 453
Zwieniecki, M.A. 464
Zürn, W. 461



abjad 376



Symbols	abugida 376	microscope 294
(378	Academia del Cimento 74	aggregate
) 378	acausality 205	overview 223
+ 378	acceleration 80	aggregates
- 378	acceleration, centrifugal 144	in nature 223
• 378	acceleration, centripetal 144,	aggregates of matte
× 378	153	air 438
: 378	acceleration, tidal 174	composition tab
< 378	accelerations, highest 88	air pressure 101
= 378	accelerometer 405	air resistance 77
> 378	accents, in Greek language 375	Airbus 406
[], measurement unit 378	accumulation 317	airflow instrument
Ø, empty set 378	accuracy 368, 388	alchemy 33
≠ 378	limits to 390	Aldebaran 83, 225
√ 378	Acetabularia 45	aleph 375
@ (at sign) 379	Acinonyx jubatus 35	algebraic surfaces 4
Δ, Laplace operator 378	Ackermann steering 412	Alice in Wonderlan
∩, set intersection 378	acoustical thermometry 466	allometric scaling 3
∪, set union 378	action 216, 238	Alpha Centauri 225
€, element of 378	as integral over time 216	alphabet 374
∞ 378	measured values 215	phonemic 376
⟨ \$, brastatevector378	action principle 208	alphabet, Greek 374
∇, 'nabla' 378	action, physical 213	alphabet, Hebrew 3
⊗, dyadic product 378	action, principle of least 163	alphabet, Latin 372
	action, quantum of 333	alphabet, story of 3
⊃, superset of/contains 378	action, quantum of, \hbar	alphabets
x , absolute value 378	physics and 8	syllabic 376
, 'similar to' 378	actuators 199	alphasyllabaries 370
	addition 235	Alps 163, 189
A	additivity 36, 42, 50, 98	Altair 225
<i>a</i> ⁿ 378	additivity of area and volume	Amazon 134
$a_n 378$	55	Amazon River 272,
a (year) 40	adenosine triphosphate 452	Amoeba proteus 86
abacus 377	aeroplane toilet 316	ampere
Abelian, or commutative 235	aeroplanes 95	definition 383
aberration 123, 137	aerostat 312	amplitude 249

AFM, atomic force

aggregate
overview 223
aggregates
in nature 223
aggregates of matter 222
air 438
composition table 439
air pressure 101
air resistance 77
Airbus 406
airflow instruments 263
alchemy 33
Aldebaran 83, 225
aleph 375
algebraic surfaces 401
Alice in Wonderland 337
allometric scaling 36
Alpha Centauri 225
alphabet 374
phonemic 376
alphabet, Greek 374
alphabet, Hebrew 376
alphabet, Latin 372
alphabet, story of 373
alphabets
syllabic 376
alphasyllabaries 376
Alps 163, 189
Altair 225
Amazon 134
Amazon River 272, 315
Amoeba proteus 86
ampere
definition 383
amplitude 249
anagyre 121

analemma 185, 430	Armillaria ostoyae 51, 97
angels 21, 91	arrow of time 47
angle 110	artefact 87, 384
in the night sky 414	ash 373
angle, plane 65	associativity 235
angle, solid 65	asteroid 184
angular acceleration 110	asteroid, Trojan 171
angular momentum 107, 110,	asteroids 179, 225
=	Astrid, an atom 464
144	
angular momentum	astrology 166, 185, 186
conservation 163	astronaut see cosmonaut
angular momentum, extrinsic	astronomers, smallest known
108, 110	168
angular momentum, intrinsic	astronomical unit 190
108	astronomy picture of the day
angular velocity 107, 110	401
anharmonicity 268	at-sign 379
anholonomic constraints 219	athletics 150
Antares 83, 225	atmosphere 143
anti-bubbles 314	angular momentum 109
anti-Hermitean 237	composition 438
anti-unitary 237	atmosphere of the Moon 344
antigravity 171, 180	atmospheric pressure 394
antigravity device 180	Atomic Age 321
antimatter 98	atomic clock 43
Antiqua 375	atomic force microscope 293,
antisymmetry 237	294, 421
apex angle 414	atomic force microscopes 202
aphelion 394	atomic mass unit 391
Aphistogoniulus	Atomium 298
erythrocephalus 86	atoms
Apis mellifera 97	arranging helium 293
apnoea 308	explain dislocations 292
apogee 394	explain round crystal
Apollo 428	reflection 292
apple trees 388	explain steps 292
apple, standard 388	Galileo and 285-287
apples 91	Greeks thinkers and 288
Aquarius 186	image of silicon 293
aqueducts 307	in ferritic steel 298
Arabic numbers 376	Lego and 288
Arabidopsis 45	photo of levitated 293
Arcturus 225	atoms are not indivisible 340
argon 438	atoms, manipulating single
Aries 186	292
Aristarchus of Samos 453	ATP 100, 452
Aristotelian view 197	atto 385
Aristotle 426	Atwood machine 426
arithmetic sequence 405	augmented fourth 263
•	

ANALEMMA

auricola 276
austenitic steels 297
average 220
Avogadro's number 290, 313,
392
axioms 34
axis of the Earth, motion of
132
axis, Earth's 137
axle, impossibility in living
beings 417
В
Babylonia 380
Babylonians 191
background 26, 27, 48
bacteria 87
bacterium 105
badminton smash, record 35
badminton smashes 409
Balaenoptera musculus 97, 272
balloon
inverted, puzzle 325
puzzle 324
balloons 325
baloon rope puzzle 312
Banach measure 55
Banach-Tarski paradox or
theorem 56, 284
banana catching 111
barycentre 187
baryon number density 396
basal metabolic rate 105
base units 383
Basiliscus basiliscus 148, 149
basilisk 148
bath-tub vortex 454
bathroom scales 143, 295
bathtub vortex 127

BCE 380

bear puzzle 60

Beaufort 35

becquerel 385 beer 314

beer mat 61

beetle, click 148

bear, colour of 60

beauty 18, 348, 356

origin of 348

before the Common Era 380	human 296	C
behaviour 20	bones 287	c. 380
belief systems 91	book and physics paradox 368	caesium 40
beliefs 22	books 372	calculating prodigies 377
bell	books, information and	calculus of variations 218
vibration patterns 252	entropy in 332	calendar 379, 380
Belousov-Zhabotinski	boom	calendar, Gregorian 380
reaction 361	sonic, due to supersonic	calendar, Julian 379
Bernoulli equation 304, 314	278	calendar, modern 380
Bernoulli's equation 304	boost 243	caliper 52, 62
Bessel functions 137	bottle 100, 117	Callisto 183
Betelgeuse 83, 84, 225	bottle, full 210	calorie 387
beth 375	boundaries 90	Calpodes ethlius 75
bets, how to win 128	boundary layer 432	camera 64
bicycle 418	brachistochrone 209, 433	Cancer 186
stability 418	braid pattern 360	candela
bicycle riding 91	brain 343	definition 384
bicycle weight 326	brain and physics paradox 368	candle 30, 345
bifurcation 356	brain stem 25	in space 441
billiards 95	brass instruments 263	canoeing 149
bimorphs 199	Braun GmbH 323, 478	Canopus 225
biographies of	bread 285, 407	cans of peas 115
mathematicians 401	breathings 375	cans of ravioli 115
biological evolution 21, 351	Bronze Age 321	capacity 318
biology 199	brooms 180, 433	Capella 225
BIPM 383, 384	brown dwarfs 225	capillary wave
bird speed 35	Brownian motion 326, 329	in water 273
bird, singing 280	typical path 328	Capricornus 186
bismuth 40	browser 399	capture, in universal gravity
bits to entropy conversion 393	bubble	186
black holes 205	soap 404	car
blasphemy 121	soap and molecular size	wheel angular momentum
block and tackle 30	291	109
blood pressure 309	bubbles 313	car engine 324
blood supply 85	bucket 19	car parking 63
board divers 111	bucket experiment, Newton's	car weight 326
boat 100	142	carbon dioxide 326
sailing 307	bullet	Carparachne 86
Bode's rule 190	speed measurement 408	carraige, south-pointing 433
bodies, rigid 209	bullet speed 35	carriage, south-pointing 209,
body 26	bullet speed measurement 58	460
body fluids 309	Bunsen burner 322	cars 368
Bohr magneton 392	Bureau International des	Cartesian 50
Bohr radius 392	Poids et Mesures 383	cartoon physics, 'laws' of 90
Boltzmann constant 325, 327,	bushbabies 148	cat 100
329, 333, 391	butterfly effect 357	Cataglyphis fortis 150
Boltzmann constant <i>k</i>	button, future 337	catenary 414
physics and 8	5 dition, facule 33/	caterpillars 75
bone		catholicism 285
~~~~		

B

BEFORE

causai 205	314
causality of motion 205	circalunar 168
cavitation 89, 267	circle packing 414
cavity resonance 281	circular definition in physics
CD	369
angular momentum 109	classical electron radius 392
Cebrennus 86, 451	classical mechanics 195
celestrocentric system 408	Clay Mathematics Institute
Celsius temperature scale 321	306
Celtic wobble stone 121	click beetles 148
cement kiln 417	clock 41, 44, 101, 157
cementite 297	clock puzzles 61, 445
centi 385	clock, air pressure powered
centre of gravity 185	101
centre of mass 108, 185	clock, exchange of hands 61
centrifugal acceleration 144,	clocks 41, 369
155	clockwise rotation 47
centrifugal force 163	closed system 337, 339
centripetal acceleration 144	Clunio 168
cerebrospinal fluid 309	coastline length 53
Ceres 186	CODATA 473
cerussite 57	coffee machines 336
Cetus 186	coin
chain 66	puzzle 316
challenge classification 9	collisions 101, 199, 367
challenges 25	colour symmetries 229
chandelier 157	comet 169
chaos 357	comet, Halley's 169
and initial conditions 358	comets 308
in magnetic pendulum 358	comic books 295
chapter sign 379	Commission Internationale
charge 242	des Poids et Mesures 383
elementary <i>e</i> , physics and	compass 209
8	completeness 42, 50, 98
positron or electron, value	complexity 332
of 391	Compton tube 130
chaturanga 30	Compton tube 130  Compton wavelength 392
check 234	Compton wavelength 392 Compton wheel 130
cheetah 35, 81	•
chemistry 367	computers 343 concatenation 235
chess 30	
	concave Earth theory 408 concepts 368
chest operations 438 child's mass 114	conditions for motion's
childhood 35, 48	existence 17
Chimborazo, Mount 163	conductance quantum 392
chocolate 310	Conférence Générale des
does not last forever 283	Poids et Mesures 383, 388
chocolate bars 283	configuration 26
Chomolungma, Mount 163,	configuration space 74

CAUSAL

connected bodies 85
conservation 98, 247
conservation of momentum
99
conservation principles 122
conservative 200
conservative systems 219
conserved quantities 17
constants
table of astronomical 393
table of astrophysical 395
table of basic physical 391
table of derived physical
392
constellations 83, 185
constraints 219
contact 367
container 48
containers 52
continental motion 133
continuity 42, 48, 50, 98, 239
continuity 42, 46, 56, 98, 239
continuity, limits of 71
continuum 36
continuum approximation 85
continuum mechanics 210
continuum physics 316
convection 339
Convention du Mètre 383
conventions 372
cooking 367
cooperative structures 355
coordinates 50, 72
coordinates, generalized 219
Copernicus 453
Corallus caninus 323
Coriolis acceleration 124, 126,
422, 453
Coriolis effect 124, 422, 453 Coriolis force 254
cork 100, 117
corn starch 361, 363
**
lower left film 25, 26
corner figures 25
corner film

Conférence Générale des Poids et Mesures 384

conic sections 169

lower left 25, 26	length of 131	direction 36
corner patterns 25	day length, past 40	disappearance of motion 99
corpuscle 83	day, length of 175	discrete 234
corrugations, road 353, 354	day, mean solar 136	dispersion 259, 260
cortex 25	day, sidereal 136, 393	dissection of volumes 56
cosmic evolution 21	day, time unit 385	dissipative 200
cosmic mirror 401	de Broglie wavelength 384	dissipative systems 202, 218,
cosmological constant 395	dead water 254	355
cosmonauts 428, 429	death 241, 369	distance 53
cosmos 27	and energy consumption	distinguish 24
coulomb 385	114	distinguishability 36, 42, 50,
countertenors 281	conservation and 101	98
crackle 417	energy and 113	distribution 262
crackpots 402	mass change with 114	Gaussian 389
creation 17	origin 15	normal 389
creation of motion 99	rotation and 107	distribution, Gaussian normal
crest of a wave 260	death sentence 285	328
cricket bowl 409	deca 385	divergence 162
crooks 364	decay 343	diving 308
cross product 104	deci 385	DNA 51, 388
crystal	deer 25	DNA (human) 224
•		
symmetry table 232	degree Celsius 385 degree, angle unit 385	DNA, ripping apart 196 Dolittle 220
crystal classes 232	delta function 262	
crystal groups 232		donate
crystallographic point groups	denseness 42, 50	for this free pdf 9
232	derivative 79	doublets 238
crystals 57	derivative at a point 79	Dove prisms 451
cumulonimbus 274	description 72	drag 198, 201
curiosities 29	descriptions, accuracy of 368	drag coefficient 201, 368
curiosity 19, 207	Desmodium gyrans 45	drifts 387
curvature 62	details and pleasure 19	dromion 280
curves of constant width 32	determinism 205, 206, 358	film of motion 280
curvimeter 54	devil 24	drop 201
cycle 268	Devonian 175	Drosophila melanogaster 45
cycloid 227, 433	diamond, breaking 295	duck
cyclotron frequency 392	Diet Coca Cola 314	wake behind 278
Cyrillic alphabet 375	differential 79	ducks, swimming 274
	differential equation 409	Duration 40
D	diffraction 259	Dutch Balloon Register 478
ð 378	diffusion 343, 344	duvet 338
dx 378	digamma 374	dwarf planets 186
daleth 375	digits, history of 376	dwarfs 225
damping 198, 259	dihedral angles 56	dx 378
dancer 75	dilations 241	dyadic product 378
angular momentum 109	dimensionality 36, 50	dynabee 146
dark stars 225	dimensionless 392	dynamic friction 200
Davey-Stewartson equation	dimensions 49	dynamics 195
280	diminished fifth 263	

dinosaurs 179

CORNER

day

measured values 331

exclusion principle 222

effort, everyday 197

Ε

effort 196

e, natural exponential 378	egg cooking 340	random motion and 330
e.g. 380	eigenvalues 237	specific, measured values
ear 49, 51, 261, 289, 462	Ekman layer 453	331
as atomic force	elasticity 199	entropy flow 338
microscope 294	elders 399	entropy, quantum of 333
illustration 276	Elea 288	entropy, smallest in nature 333
ear problems 276	Electric Age 321	entropy, state of highest 346
ear, human 276	electric effects 199	environment 26
Earth 123, 339	electrodynamics 316	Epargyreus clarus 75, 450
age 394	electromagnetism 160	ephemeris time 41
angular momentum 109	electron see also positron	eponyms 381
average density 394	g-factor 393	EPR 381
equatorial radius 394	electron charge see also	equal intonation 265
flattening 394	positron charge	equilibrium 336
gravitational length 394	electron magnetic moment	equilibrium, thermal 336
mass 393	392	Eris 186
normal gravity 394	electron mass 391	eros 159
stops rotating 163	electron speed 35	error
Earth dissection 56	electron volt	relative 389
Earth from space 401	value 393	systematic 389
Earth rotation	element of set 36, 42, 50, 98	total 389
speed of 131	elementary particles 300	errors
Earth speed 35	elephants 274, 348	example values 367
Earth's age 40	ellipse 168	in measurements 388
Earth's axis 137	as orbit 155	ESA 401
Earth's mass, time variation	email 399	escape velocity 184
179	emergence 358	Escherichia coli 87
Earth's radius 394	emergent properties 358	et al. 380
Earth's rotation 386	emit waves 261	eth 373, 471
Earth's rotation change of 131	empty space 162, 260	ethel 373
Earth's shadow 147	EMS98 145	ethics 207
Earth's speed through the	energy 101, 102, 198, 225, 239	Eucalyptus regnans 287
universe 140	conservation 161	Eucharist 285
Earth, flat 416	energy conservation 104, 163,	Euclidean space 49, 50
Earth, flattened 123, 143	242	Euclidean vector space 34, 36
Earth, hollow 57	energy consumption in First	Europa 183
Earth, mass of 158	World countries 104	evaporation 343
Earth, shape of 162	energy flows 99	evenings, lack of quietness of
earthquake 145, 163	energy of a wave 258	274
earthquake, starting 295	energy, observer	event 40
earthquake, triggered by	independence 239	Everest, Mount 163, 314
humans 295	energy, scalar generalization	everlasting life 101
earthquakes 145, 168, 455	of 239	everything flows 17
eccentricity 171	engine	evolution 26
eccentricity of Earth's axis 138	car 324	evolution equations 204
echo 261	enlightenment 177	Exa 385
ecliptic 164	entropy 318, 369	exclamation mark 379

definition 330, 332

avnancian as a	fifth perfect acc	Equalities nondulum wah
expansion 320	fifth, perfect 263	Foucault's pendulum, web
expansion of the universe 140	figures in corners 25	cam 128
expansions 241	filament 322	fountain, Heron's 311
Experience Island 16, 21	fine structure constant 391	Fourier analysis 252
exponential notation 63	finite 329	Fourier decomposition 252
extended bodies, non-rigid	fire 322	Fourier transformation 277
210	fire pump 330	fourth, perfect 263
extension 300	firework 65	fractal
and waves 251	fish's eyes 314	landscapes 284
extensive quantities 318	flagella 87	fractals 54, 80, 283
extensive quantity 99	flagellar motor 105	frames of reference 229
extrasolar planets 225	flame 30, 144, 341	frass 75
eye 25	flatness 57	free fall, speed of 35
eye motion 49	flattening	free will 206
eye, blinking after guillotine	of the Earth 123, 143	frequency 253
94	fleas 148	measured values 250
eyelid 62	flies 25	friction 100, 101, 198, 200, 201,
eyes of fish 314	flip film 25, 26	336, 338
,	explanation of 72	picture of heat 320
F	flow 317	Friction between planets and
$\varphi x$ 378	minimum, measured	the Sun 101
f(x) 378	values 335	friction produced by tides 175
f'(x) 378	flow of time 47	friction, importance of 336
F. spectabilis 45	fluid	froghopper 81
F. suspensa 45	and self-organization 351	Froude number, critical 273
F. viridissima 45	fluid mechanics 210	fuel consumption 368
Falco peregrinus 35	fluid motion	full width at half maximum
fall 154	example pictures 303	389
fall and flight are independent	ink puzzle 346	funnel 314
	know-how 307	
fall is not wantical to 4		puzzle 313
fall is not vertical 124	fluids, body 309	Futhers 373
fall is parabolic 74	flute 248	Futhorc 373
fall of Moon 154	fly, common 159	futhorc 375
false quint 263	flying saucers 274	future button 337
familiarity 24	focal point 416	future, fixed 205
family names 381	focus 416	future, remembering 337
fantasy and physics 370	foetus 266	fx 378
farad 385	force 96, 110	
Faraday's constant 392	definition of 196	G
farting for communication	force, central 96	Gaia 357
274	force, physical 196	gait 112
fear of formulae 31	force, use of 196	galaxy centre 140
feet 309	Ford and precision 368	galaxy cluster 223
femto 385	forest 25	galaxy group 223
Fermi coupling constant 391	forget-me-not 230	galaxy supercluster 223
Fermi problems 341	formulae, ISO 471	Galilean physics 28, 33, 195,
Fermilab 63	formulae, liking them 31	366
ferritic steels 297	formulae, mathematical 401	Galilean satellites 183

Forsythia europaea 45

Galilean space 49

Fiat Cinquecento 312

EXPANSION

Galilean space-time 71
Galilean time 42
Galilean transformations 240
Galilean velocity 34
galvanometer 199
Ganymede 183
gas 325
gas constant, universal 392
gas planets 225
gases
as particle collections 339
gauge change 234
gauge symmetries 241
gauge theory 48
Gaussian distribution 328, 389
Gemini 186
generalized coordinates 220
generators 242
Geocentric gravitational
constant 394
geocentric system 188
geodesics 202
geoid 162
geological maps 402
geometric sequence 405
geometric symmetries 229
geostationary satellites 171
Gerridae 149
ghosts 30, 91
Giant's Causeway 57, 361
giants 287
Giga 385
gimel 375
girl-watching 147
glass 294
Global infrasound Network
277
global warming 275, 339
gnomonics 41
gods 33, 121, 142, 149, 172, 445,
447
gold 407
surface atoms 335
golf balls 409
Gondwanaland 133
gorilla test for random
numbers 205
Gosper's formula 343
Gothic alphabet 375
· · · · · · · · · · · · · · · · · · ·

GALILEAN

Gothic letters 375
GPS 41
grace 22, 112, 363, 446
gradient 160
granular jet 298
gravitation see also universal
gravitation, 151, 199
essence of 189
gravitation and
measurements 370
gravitation, universal see
universal gravitation
gravitational acceleration,
standard 81
gravitational constant 152, 39
gravitational constant G
physics and 8
gravitational coupling
constant 391
gravitational field 162
gravitational mass 177
gravity see gravitation
gravity inside matter shells
189
gravity waves 254
gravity, centre of 185
gravity, sideways action of 157
gray 385
Greek alphabet 374
Greek number system 377
greenhouse effect 339
Greenland 40
Gregorian calendar 380
Gregorian calendar reform 6
group, mathematical 235
growth 21, 211
as self-organization 351
growth of deep sea
manganese crust 35
GRW approach 381
Gulf Stream 453
guns and the Coriolis effect
127
gymnasts 111
gynaecologist 266
gyroscope 111, 128
gyroscope, laser 130
<i>57</i> 1 - 7 3 -

hafnium carbide 322
Hagedorn temperature 322
hair clip 200
hair growth 35
hair, diameter 51
halfpipe 227
Hamilton's principle 213
hammer drills 116
hands of clock 61
hard discs, friction in 202
harmonic motion 248
heartbeat 157
heat
in everyday life 330
in physics 330
heat & pressure 199
heat engines 324
Hebrew alphabet 376
hecto 385
helicity of stairs 47
helicopter 314
helicopters 402
heliocentric system 188
helium 223, 322, 343
helium, superfluid 312
henry 385
heresy 285
Hermitean 237
Heron's fountain 311
herring, farting 274
hertz 385
hiccup 364
Himalaya age 40
hips 112
hodograph 74, 153, 182
hole
through the Earth 192
holes 27
hollow Earth theory 57
Hollywood 220
Hollywood films 213, 283, 451
holonomic 219
holonomic systems 219
holonomic-rheonomic 219
holonomic-scleronomic 219
Homo sapiens 45
homogeneity 42, 50
homomorphism 237

honey bees 97	thermodynamics 333	Alphabet 372
Hooke 164	index of refraction 434	internet 398, 399
Hopi 448	Indian digits 376	list of interesting websites
horse 105	Indian number system 377	399
horse power 105	Indian numbers 376, 377	interval, musical 263
horses, speed of 75	Indian numbers and digits	invariance 98, 235
hour 385	376	invariant property 229
hourglasses 59	individuality 27, 203	inverse element 235
Hubble parameter 395	inductions 243-245	invisibility of objects 85
Hubble space telescope 401	inertia 91	invisible loudspeaker 342
Hudson Bay 40	inertial mass 177	Io 175, 183
human growth 35	inf. 380	ionosphere, shadow of 147
humour 359	infinite coastlines 54	IPA 372
humour, physical 402	infinite number of SI prefixes	Iron Age 321
Huygens' principle 261	388	irreducible 237
illustration of 262	infinity 17, 36, 42, 50, 98	irreversibility of motion 205
illustration of consequence	infinity in physics 80	irreversible 336
262	information 332	Island, Experience 16
hydrofoils 307	information, erasing of 343	isolated system 339
Hydroptère 35	information, quantum of 333	isomorphism 237
hyperbola 168	infrasound 277	isotomeograph 128
	initial condition	Istiophorus platypterus 35, 309
I	unfortunate term 206	Isua Belt 40
i, imaginary unit 378	initial conditions 203	italic typeface 372
	injective 237	IUPAC 473
i.e. 380	111/001110 25/	101110 4/3
ibid. 380	inner world theory 408	IUPAP 473
ibid. 380 IBM 478 ice ages 138, 455	inner world theory 408	
ibid. 380 IBM 478	inner world theory 408 Inquisition 285	IUPAP 473
ibid. 380 IBM 478 ice ages 138, 455	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40	IUPAP 473
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17	J jerk 203, 417 Jesus 149 joints
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40	J jerk 203, 417 Jesus 149 joints cracking 281
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17	J jerk 203, 417 Jesus 149 joints cracking 281 Josephson effect 384
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331 ideal 51 ideal gas 320, 325 ideal gas constant 320, 440	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17 insulation power 338 integral 216 integration 55, 216	J jerk 203, 417 Jesus 149 joints cracking 281 Josephson effect 384 Josephson frequency ratio 392
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331 ideal 51 ideal gas 320, 325 ideal gas constant 320, 440 idem 380	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17 insulation power 338 integral 216	J jerk 203, 417 Jesus 149 joints cracking 281 Josephson effect 384 Josephson frequency ratio 392 joule 385
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331 ideal 51 ideal gas 320, 325 ideal gas constant 320, 440 idem 380 Illacme plenipes 89	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17 insulation power 338 integral 216 integration 55, 216 integration, symbolic 401 interaction 27, 196	J jerk 203, 417 Jesus 149 joints cracking 281 Josephson effect 384 Josephson frequency ratio 392 joule 385 juggling 76
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331 ideal 51 ideal gas 320, 325 ideal gas constant 320, 440 idem 380	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17 insulation power 338 integral 216 integration 55, 216 integration, symbolic 401	J jerk 203, 417 Jesus 149 joints cracking 281 Josephson effect 384 Josephson frequency ratio 392 joule 385 juggling 76 juggling robot 76
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331 ideal 51 ideal gas 320, 325 ideal gas constant 320, 440 idem 380 Illacme plenipes 89	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17 insulation power 338 integral 216 integration 55, 216 integration, symbolic 401 interaction 27, 196	J jerk 203, 417 Jesus 149 joints cracking 281 Josephson effect 384 Josephson frequency ratio 392 joule 385 juggling 76
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331 ideal 51 ideal gas 320, 325 ideal gas constant 320, 440 idem 380 Illacme plenipes 89 illness 309	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17 insulation power 338 integral 216 integration 55, 216 integration, symbolic 401 interaction 27, 196 interface waves on an 254 interference 259	J jerk 203, 417 Jesus 149 joints cracking 281 Josephson effect 384 Josephson frequency ratio 392 joule 385 juggling 76 juggling robot 76
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331 ideal 51 ideal gas 320, 325 ideal gas constant 320, 440 idem 380 Illacme plenipes 89 illness 309 illusions of motion 16	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17 insulation power 338 integral 216 integration 55, 216 integration, symbolic 401 interaction 27, 196 interface waves on an 254	J jerk 203, 417 Jesus 149 joints cracking 281 Josephson effect 384 Josephson frequency ratio 392 joule 385 juggling 76 juggling robot 76 Julian calendar 379 jump 148 jump, long 450
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331 ideal 51 ideal gas 320, 325 ideal gas constant 320, 440 idem 380 Illacme plenipes 89 illness 309 illusions of motion 16 illusions, optical 402 image 90 images 27	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17 insulation power 338 integral 216 integration 55, 216 integration, symbolic 401 interaction 27, 196 interface waves on an 254 interference 259	J jerk 203, 417 Jesus 149 joints cracking 281 Josephson effect 384 Josephson frequency ratio 392 joule 385 juggling 76 juggling robot 76 Julian calendar 379 jump 148
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331 ideal 51 ideal gas 320, 325 ideal gas constant 320, 440 idem 380 Illacme plenipes 89 illness 309 illusions of motion 16 illusions, optical 402 image 90 images 27 imagine 234	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17 insulation power 338 integral 216 integration 55, 216 integration, symbolic 401 interaction 27, 196 interface    waves on an 254 interference 259    illustration 260 interferometer 95, 130 internal 234	J jerk 203, 417 Jesus 149 joints cracking 281 Josephson effect 384 Josephson frequency ratio 392 joule 385 juggling 76 juggling robot 76 Julian calendar 379 jump 148 jump, long 450 jumping height of animals 76 Jupiter 188, 225
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331 ideal 51 ideal gas 320, 325 ideal gas constant 320, 440 idem 380 Illacme plenipes 89 illness 309 illusions of motion 16 illusions, optical 402 image 90 images 27 imagine 234 immovable property 92	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17 insulation power 338 integral 216 integration 55, 216 integration, symbolic 401 interaction 27, 196 interface    waves on an 254 interference 259    illustration 260 interferometer 95, 130 internal 234 International Earth Rotation	J jerk 203, 417 Jesus 149 joints cracking 281 Josephson effect 384 Josephson frequency ratio 392 joule 385 juggling 76 juggling robot 76 Julian calendar 379 jump 148 jump, long 450 jumping height of animals 76 Jupiter 188, 225 angular momentum 109
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331 ideal 51 ideal gas 320, 325 ideal gas constant 320, 440 idem 380 Illacme plenipes 89 illness 309 illusions of motion 16 illusions, optical 402 image 90 images 27 imagine 234 immovable property 92 impenetrability 98	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17 insulation power 338 integral 216 integration 55, 216 integration, symbolic 401 interaction 27, 196 interface    waves on an 254 interference 259    illustration 260 interferometer 95, 130 internal 234 International Earth Rotation    Service 386, 455, 457	J jerk 203, 417 Jesus 149 joints cracking 281 Josephson effect 384 Josephson frequency ratio 392 joule 385 juggling 76 juggling robot 76 Julian calendar 379 jump 148 jump, long 450 jumping height of animals 76 Jupiter 188, 225
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331 ideal 51 ideal gas 320, 325 ideal gas constant 320, 440 idem 380 Illacme plenipes 89 illness 309 illusions of motion 16 illusions, optical 402 image 90 images 27 imagine 234 immovable property 92 impenetrability 98 impenetrability of matter 33	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17 insulation power 338 integral 216 integration 55, 216 integration, symbolic 401 interaction 27, 196 interface    waves on an 254 interference 259    illustration 260 interferometer 95, 130 internal 234 International Earth Rotation	J  jerk 203, 417  Jesus 149  joints  cracking 281  Josephson effect 384  Josephson frequency ratio 392  joule 385  juggling 76  juggling robot 76  Julian calendar 379  jump 148  jump, long 450  jumping height of animals 76  Jupiter 188, 225  angular momentum 109  Jupiter's atmospheric pressure 394
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331 ideal 51 ideal gas 320, 325 ideal gas constant 320, 440 idem 380 Illacme plenipes 89 illness 309 illusions of motion 16 illusions, optical 402 image 90 images 27 imagine 234 immovable property 92 impenetrability 98 impenetrability of matter 33 inclination of Earth's axis 138	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17 insulation power 338 integral 216 integration 55, 216 integration, symbolic 401 interaction 27, 196 interface waves on an 254 interference 259 illustration 260 interferometer 95, 130 internal 234 International Earth Rotation Service 386, 455, 457 International Geodesic Union 395	J jerk 203, 417 Jesus 149 joints cracking 281 Josephson effect 384 Josephson frequency ratio 392 joule 385 juggling 76 juggling robot 76 Julian calendar 379 jump 148 jump, long 450 jumping height of animals 76 Jupiter 188, 225 angular momentum 109 Jupiter's atmospheric pressure 394 Jupiter's mass 394
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331 ideal 51 ideal gas 320, 325 ideal gas constant 320, 440 idem 380 Illacme plenipes 89 illness 309 illusions of motion 16 illusions, optical 402 image 90 images 27 imagine 234 immovable property 92 impenetrability 98 impenetrability of matter 33	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17 insulation power 338 integral 216 integration 55, 216 integration, symbolic 401 interaction 27, 196 interface    waves on an 254 interference 259    illustration 260 interferometer 95, 130 internal 234 International Earth Rotation    Service 386, 455, 457 International Geodesic Union	J jerk 203, 417 Jesus 149 joints cracking 281 Josephson effect 384 Josephson frequency ratio 392 joule 385 juggling 76 juggling robot 76 Julian calendar 379 jump 148 jump, long 450 jumping height of animals 76 Jupiter 188, 225 angular momentum 109 Jupiter's atmospheric pressure 394 Jupiter's mass 394 Jupiter's moons 191
ibid. 380 IBM 478 ice ages 138, 455 iceberg 97 icicles 295 icosane 331 ideal 51 ideal gas 320, 325 ideal gas constant 320, 440 idem 380 Illacme plenipes 89 illness 309 illusions of motion 16 illusions, optical 402 image 90 images 27 imagine 234 immovable property 92 impenetrability 98 impenetrability of matter 33 inclination of Earth's axis 138	inner world theory 408 Inquisition 285 insects 168, 344 instant 33, 40 instant, human 40 instant, single 17 insulation power 338 integral 216 integration 55, 216 integration, symbolic 401 interaction 27, 196 interface waves on an 254 interference 259 illustration 260 interferometer 95, 130 internal 234 International Earth Rotation Service 386, 455, 457 International Geodesic Union 395	J jerk 203, 417 Jesus 149 joints cracking 281 Josephson effect 384 Josephson frequency ratio 392 joule 385 juggling 76 juggling robot 76 Julian calendar 379 jump 148 jump, long 450 jumping height of animals 76 Jupiter 188, 225 angular momentum 109 Jupiter's atmospheric pressure 394 Jupiter's mass 394



K	lawyers 92
k-calculus 406	laziness, cosmic, principle of
Kadomtsev-Petviashvili	209, 218
equation 281	lead 312
kefir 224	leaf, falling 314
kelvin	leaning tower in Pisa 73
definition 383	leap day 379
Kepler's laws 164	leapfrogging of vortex rings
ketchup motion 35	316
kilo 385	learning 24
kilogram	without markers 8
definition 383	learning mechanics 197
kilotonne 103	learning, best method for 8
kinematics 72, 195	Lebesgue measure 56
kinetic energy 101	lecture scripts 401
Klitzing, von – constant 392	leg 157
knot 70	leg performance 148
knowledge, humorous	Lego 289, 294
definition of 432	legs 29
knuckle angles 65	advantages 148
koppa 374	vs wheels 148
Korteweg-de Vries equation	legs in nature 85
270, 281	Leidenfrost effect 313
Kuiper belt 186, 224, 308	length 49, 53, 78, 370
Kurdish 373	length scale 344
	Leo 186
L	letters 373
ladder 117	Leucanthemum vulgare 211
sliding 117	levitation 171
ladder puzzles 67	Libra 186
Lagrangain	librations 167
points in astronomy 428	lichen growth 35
Lagrange's equations of	life
motion 219	everlasting 101
Lagrangian	life, shortest 40
examples of 221	lifespan
Lagrangian (function) 216	animal 113
Lagrangian is not unique 219	lift
Lagrangian libration points	for ships 119
171	lift into space 294
Lake Nyos 326	light 199
laminarity 302	light deflection near masses
languages on Earth 376	177
large 249	light mill 199
laser gyroscopes 130	light speed measurement 58
laser loudspeaker 342	light tower 35
lateral force microscopes 202	light year 393
Latin 381	light, slow group velocity 448
Latin alphabet 372	lightbulb temperature 322

mach 95
Mach number 143
Mach's principle 96
machine 87
moving giant 300
machines
examples of high-tech 299
magentization of rocks 135
magic 402
Magna Graecia 288
magnetic effects 199

K-CALCULUS

magnetic flux quantum 392	matter 82, 91
magnetic thermometry 466	matter shell, gi
magnetism 96	matter, impend
magneton, nuclear 393	meaning 19
magnitude 78	measurability
magnitudes 239	measure 370
main result of modern	measurement
science 334	comparisor
male sopranos 281	definition 3
man, wise old 220	errors, exar
manakin, club winged 250	irreversibili
many-body problem 366	meaning 38
marble, oil covered 100	process 386
marker, bad for learning 8	measurement
Mars 386	definition 3
martensitic steels 297	measurements
mass 93, 97, 198, 370	gravitation
concept of 93	mechanics 195
mass conservation implies	mechanics, cla
momentum conservation	mechanics, qu
96	Medicean sate
mass is conserved 93	medicines 29
mass of children 114	Mega 385
mass of Earth, time variation	megatonne 103
179	memorize 24
mass point 83	memory 30, 33
mass ratio	menstrual cycl
neutron-electron 393	Mentos 314
neutron-proton 393	Mercalli scale
proton-electron 393	Mercury 172
mass, centre of 185	mercury 291
mass, Galilean 97	meridian 157
mass, gravitational 177	Mesopotamia
mass, identity of gravitational	-
and inertial 178	metabolic scor
mass, inertial 91, 177	metallurgy 367
mass, negative 98	meteorites 179
math forum 401	meteroids 225
mathematicians 401	metre
mathematics 34	definition 3
maths problem of the week	metre sticks 36
_	
401	metricity 36, 4
matrix, adjoint 237	micro 385
matrix, antisymmetric 237	microscope
matrix, complex conjugate	atomic forc
237	microscope, at
matrix, orthogonal 237	421
matrix, symmetric 237	microwave bac

matrix, transposed 237

ter 82, 91	mile 386
ter shell, gravity inside 189	Milky Way
ter, impenetrability of 33	angular momentum 109
ning 19	Milky Way's age 395
surability 36, 42, 50, 98	Milky Way's mass 395
sure 370	Milky Way's size 395
surement 36	milli 385
comparison 386	mind change 21
lefinition 383, 386	minerals 402
errors, example values 367	mini comets 186
rreversibility 386	minimum
neaning 386	of curve, definition 218
process 386	minor planets 225
surement error	minute 385, 395
lefinition 388	Mir 193
surements and	mirror invariance 28
gravitation 370	of everyday motion 246
hanics 195, 209	moguls, ski 354
hanics, classical 195	mol 290
hanics, quantum 195	molar volume 392
licean satellites 183	mole
licines 29	definition 383
ga 385	particles in a 290
gatonne 103	molecules 289
norize 24	moment 41
nory 30, 336	moment of inertia 107, 110,
nstrual cycle 168, 457	240, 245
ntos 314	moment of inertia, extrinsic
calli scale 145	108
cury 172	moment of inertia, intrinsic
cury 291	108
idian 157	momentum 94, 98, 102, 198,
opotamia 372	225
abolic rate 114	as a liquid 196
abolic scope, maximal 453	change 196
allurgy 367	flow 196
eorites 179, 277	momentum conservation
eroids 225	follows from mass
re	conservation 96
lefinition 383	momentum flows 99
re sticks 369	momentum of a wave 258
ricity 36, 42, 50, 98	momentum, angular 108
ro 385	money, humorous definition
roscope	of 432
tomic force 293	month 379
roscope, atomic force 294,	Moon 154
21	angular momentum 109
rowave background	angular size 394
emperature 396	density 394

distance 394
mass 394
radius 394
Moon calculation 154
Moon density and tides 175
Moon illusion 66
Moon path around Sun 184
•
Moon phase 188
Moon size illusion 66
Moon size, angular 66
Moon size, apparent 154
Moon's atmosphere 344
Moon's mean distance 394
Moon's surface gravity 394
Moon, dangers of 175
Moon, fall of 154
moons 224
Moons's atmospheric pressure
394
moped 121
morals 207
mornings, quietness of 274
moth 307
motion 27, 369
does not exist 16
in configuration space 357
is fundamental 384
is predictable 366
mirror invariance 246
parity invariance 246
reversibility 246
motion and measurement
units 384
motion as an illusion 17
motion as illusion 17
motion detectors 30
motion illusions, figures
showing 16
motion invariance 28
motion is based on friction
200
motion is change of position
with time 72
motion is due to particles 23
motion is relative 26
motion is the change of state
of objects 28
Motion Mountain 16

aims of book series 7

Moon

supporting the project 9
motion of continents 35
motion, conditions for its
existence 17
motion, drift-balanced 447
motion, faster than light 338
motion, harmonic 248
motion, infinite 370
motion, manifestations 16
motion, non-Fourier 447
motion, passive 22, 200
motion, simplest 24
motion, unlimited 370
motion, volitional 22
motion, voluntary 200
motor bike 92
motor, electrostatic 199
motor, linear 199
motors 199
moustache 52
movable property 92
movement 22
multiplet 235, 236
multiplication 235
muon
g-factor 393
muon magnetic moment 393
muon mass 391
Musca domestica 45, 81
music
notes and frequencies 265
Myosotis 230
myosotis 229
mystery of motion 15
injutery of motion 19
N
nabla 378
names of people 381
Nannosquilla decemspinosa 86
nano 385
Nanoarchaeum equitans 224
NASA 386
NASA 401
natural 51
natural unit 392
nature 27
Navier-Stokes equations 306
needle on water 315
negative vector 77

neocortex 446

observables, discrete 239	π and gravity 157	performers 399
ocean	π, circle number 378	perigee 394
origin of 308	$\pi = 3.141592470$	perihelion 172, 394
oceanography 277	paerlite 297	perihelion shift 138, 171
octave 263	painting puzzle 193	period 249
octet 238	paper aeroplanes 402	permanence 25
odometer 53	paper boat contest 312	permanence of nature 17
ohm 385	paper cup	permeability, vacuum 391
oil 100, 313	puzzle 341	permittivity, vacuum 391
oil film experiment 284	parabola 66, 74, 155, 168	permutation symmetry 234
Olympic year counting 380	parachutes 200	perpetual motion machine
Oort cloud 224	paradox	101
op. cit. 380	hydrodynamic 305	perpetuum mobile 98
operator 162	hydrostatic 305	perpetuum mobile, first and
Ophiuchus 186	paradox about physics books	second kind 101
optical radiation	368	Peta 385
thermometry 466	parallax 123, 137	phase 249, 253
orbit, elliptical 165	parallelepiped 106	phase space 27, 203
order 42, 50, 98, 348	parenthesis 379	phase space diagram 74
table of observed	parity	phase velocity 253
phenomena 348	inversion 246	PhD, enjoying it 363
order appearance	parity invariance 28	Philaenus spumarius 81
examples 348, 351	parking	photoacoustic effect 342, 343
mathematics of 356	mathematics 410	photon
order parameter 355	parking challenge 63	number density 396
orgasms 364	parsec 393	physical observations 233
orientation change needs no	particle 83	physical system 27
background 111	particle data group 400	physicists 400
origin	parts 27, 82, 335	physics 15
human 15	pascal 385	map of 8
Orion 84	passim 380	physics problems 400
ornament	passive motion 204	physics, circular definition in
Hispano-Arabic 236	past	369
orthogonality 78	of a system 203	physics, everyday 28
oscillation 248	path 72	physics, outdated definition
damped 249	Paul trap 293	22
definition 248	pea dissection 56	pico 385
harmonic and anharmonic	pearls 294	piezoelectricity 199
249	peas 115	pigeons 112
oscillation, harmonic or	Peaucellier-Lipkin linkage	ping-pong ball 144
linear 248	209	pinna 276
oscillons 354	pee research 360	Pioneer satellites 81
osmosis 199, 343	peers 399	Pisces 186
otoacoustic emissions 261	Peirce's puzzle 61	Planck constant
outer product 378	pencil, invention of 33	value of 391
oxygen 311	pencils 60	planet 101
	pendulum 157	planet-Sun friction 101
P	penguins 112	planetoids 175, 224, 225
$\varphi x$ 378	people names 381	planets

OBSERVABLES

and universal gravitation	precession, equinoctial 132
164	precision 19, 30, 34, 72, 368,
plants 29	388
plate tectonics 135	limits to 390
play 22	predictability of motion 366
Pleiades star cluster 152	prefixes 385, 472
pleural cavity 304, 438	prefixes, SI 385
Pleurotya ruralis 86	preprints 399, 400
plumb-line 57	pressure 197, 289, 325
Pluto 186	definition 302, 305
pneumothorax 438	measured values 304
point mass 83	principle of gauge invariance
point particle 83	234
point, mathematical 49	principle of least action 123,
point-like for the naked eye 85	163, 218
points 33, 85	principle of relativity 234, 40
poisons 29	principle of the straightest
Poisson equation 162	path 202
polar motion 132	principle of thermodynamic
Polaris 225	second 337
polarization 259	principle, extremal 213
pollen 327	principle, variational 213
pollutants 438	prism 451
pollution 438	prize, one million dollar 306
pool filled with corn starch	problems, physical 400
and water 363	process
pop 417	change in 217
Pororoca 272	process, sudden 207
Porpoise Cove 40	processes 336
positional system 377	Procyon 225
positions 77	prodigies, calculating 377
positivity 98	product, dyadic 378
positron charge	product, outer 378
value of 391	pronunciation, Erasmian 375
positron charge, specific 392	proof 34
possibility of knots 50	propagation velocity 253
postcard 59	propeller 417
stepping through 408	propellers in nature 85
potassium 268, 344	proper time 41
potential energy 161	proper velocity 35
potential, gravitational 160	properties, emergent 358
power, humorous definition	properties, intrinsic 27
of 432	properties, permanent 27
power, physical 104, 197	protestantism 285
ppm 439	proton  Compton wavelength 202
praying effects 225	Compton wavelength 393
praying effects 225	gyromagnetic ratio 393
precession 138	magnetic moment 393

specific charge 393

precession of a pendulum 128

proton age 40
proton mass 391
proton-muon mass ratio 393
pseudovector 144
PSR 1257+12 225
PSR 1913+16 40
psychokinesis 22
pulley 199
pulsar period 40
pulse 261
pyramid 106
Pythagoras' theorem 64
-
Q
O factor a co

#### Q-factor 249 qoppa 374 quanta smallest 288 quanti, piccolissimi 285, 286 quantities 239 conserved 17 quantity extensive types of 317 quantity of matter 94, 98 quantity, extensive 99 Quantum Age 321 quantum mechanics 195 quantum of circulation 392 quantum of entropy 333 quantum of information 333 quantum theory 26, 286 quartets 238 quartz 251 oscillator film 251 quasar 223 quietness of mornings 274 quotations, mathematical 402

# R R-complex 446 radian 65, 384 radiation 91, 339 radiation thermometry 466 radio speed 35 rain drops 76, 201 rain speed 35 rainbow 239

random 205

PLANTS

random errors 388	reversible 336
random pattern 25	Reynolds number 361
randomness 205	Rhine 315
rank 240	rhythmites, tidal 455
Ranque-Hilsch vortex tube	Richter magnitude 145
345	riddle 16
ratio between the electron	Rigel 225
magnetic moment and the	right-hand rule 104
Bohr magneton 367	ring laser 454
rattleback 121	rings, astronomical, and tides
ravioli 115	174
ray form 188	road, corrguated or
real numbers 42, 49	washboard 354
recognition 25	Roadrunner 295
recognize 25	robot 460
reducible 237	walking on water 149
reed instruments 263	robot juggling 76
reel 19	robot, walking 227
reflection of waves 259	Roche limit 432
refraction 227, 259	rock
and minimum time 226	self-organization 362
illustration 273	rock magnetization 135
refractive index 227	rocket launch sites 143
Regulus 225	rocket motor 199
relations 27	roller coasters 193
relativity 209	rolling 111
relativity, Galileo's principle	rolling puzzle 32
of 141	rolling wheels 111
reluctance to turn 107	Roman number system 377
representation 237	rope
representation of the	around Earth 409
observables 240	motion of hanging 363
representations 236	ropes
reproducibility 233	geometry of 69
reproduction 15	Rosby length 143
research 202	Rostock, University of 312
research in classical	Roswell incident 274
mechanics 366	rotation
	absolute or relative 142
resolution, of measurements	
28	as vector 110
resonance	rotation and arms 142
definition 251	rotation change of Earth 131
resources 399	rotation of the Earth 386
rest 17	rotation rate 107
rest, Galilean 78	rotation sense in the athletic
Reuleaux curves 405	stadium 47
reversibility	rotation speed 107
of everyday motion 246	rotational energy 108
reversibility of motion 205	roulette and Galilean

RANDOM

mechanics 115
Rubik's cube 276
rugby 102
Runic script 373
running 113
running backwards 340
running on water 456
running reduces weight 182
running speed record, human
75
Rydberg constant 367, 392
7 8
S
Sagarmatha, Mount 163, 314
Sagittarius 140, 186
Sagnac effect 130, 423
sailfish 35
sailing 100
sails 306
Salmonella 87
sampi 374
san 374
sand 352
granular jets in 299
self-organization 352
singing 353
table of patterns in 353 sand, Cuban 353
Saraeva 445
Sarcoman robot 76
satellite 84
satellite, artificial 84
satellites 145, 186
satellites, observable 401
Saturn 174, 186
scalar 78, 230, 239
scalar product 50, 78
scale invariance 287
scale symmetry 229
scales
bathroom 295
scales, bathroom 143
scaling law 44
Scorpius 186
screw 200
scripts, complex 376
sea 255

sea wave energy 277 sea waves, highest 272

season 164	shore birds 456	snowflake speed 35
second 385, 395	short pendulum 422	snowflakes 468
definition 383	shortest measured time 40	soap bubbles 291, 313
Sedna 186	shot noise 289	sodium 268, 322, 344, 448
selenite 57	shot, small 312	solar data 401
self-adjoint 237	shoulders 112	Solar system
self-organization 211	shutter 63	angular momentum 109
self-similarity 54	shutter time 64	solar system 224
sequence 39, 42, 50, 98	SI prefixes 388	solar system formation 132
Sequoiadendron giganteum 97	SI units 383, 390	solar system simulator 401
serpent bearer 186	definition 383	solar system, future of 190
Serpentarius 186	prefixes 385	solitary wave
sets, connected 79	supplementary 384	in sand 353
sexism in physics 288	siemens 385	solitary waves 270
sextant 456	sievert 385	soliton 271
sha 374	signal	film 270
shadow of ionosphere 147	decomposition in	solitons 270
shadow of the Earth 147	harmonic components 252	sonar 280
shadow of the Earth during a	list of types 264	sonic boom 278
lunar eclipse 154	similar 27	sonoluminescence 89
shadows 90, 452	Simon, Julia 451	soul 114
and attraction of bodies	sine curve 248	sound
189	singing 263	film of molecular motion
shadows of sundials 47	singlets 238	in 265
shampoo, jumping 316	singular 237	intensity table 266
shape 50, 82, 300	sink vortex 454	sound channel 274
shape deformation and	siren 272	sound speed 35
motion 87	Sirius 84, 225	sound threshold 266
shape of the Earth 185	situation 26	source term 162
sharp s 373	size 48	sources 162
sheep, Greek 375	skateboarding 227	South Pole 162
Shell study 452	skew-symmetric 237	south-pointing carriage 209
shell, gravity inside matter 189	ski moguls 271, 353, 354	space 369
ship 97	skin 85	absolute 128
critical speed 273	skipper 75	space is necessary 50
leaving river 310	sky, moving 132	space lift 294
lift 119	slide rule 414, 415	space points 85
mass of 119	slingshot effect 182	space sickness 429
pulling riddle 312	Sloan Digital Sky Survey 401	space travel 143
speed limit 273	small solar system bodies 225	space, absoluteness of 50
wake behind 278	smallest experimentally	space, absoluteness of 30 space, physical 48
ships and relativity 141	probed distance 63	space, physical 48 space, relative or absolute 52
shit 75, 450	smiley 379	space, white 379
shock wave	smoke 328	space, write 3/9 space-time 27
	_	
due to supersonic motion	ring picture 315	space-time diagram 74
278	snap 417	space-time, Galilean 71
shock waves 277	snorkeling 438	space-time, relative or
shoe size 387	snow flake	absolute 52

self-organization film 355

spatial inversion 241

shoelaces 49, 336

special relativity before the	star age 40	subscripts 375
age of four 234	star classes 225	Sun 97, 101, 188, 225
specific mechanical energy	starch	angular momentum 109
416	self-organization 362	will rise tomorrow 163
spectrum, photoacoustic 343	stargazers 401	Sun density and tides 175
speed 79	stars 83, 225	Sun size, angular 66
lowest 30	state 27	Sun's age 394
of light <i>c</i>	of a mass point, complete	Sun's lower photospheric
physics and 8	203	pressure 395
speed of light	of a system 203	Sun's luminosity 394
value of 391	state allows one to distinguish	Sun's mass 394
speed of light inside the Sun	27	Sun's surface gravity 395
35	state of motion 26	Sun, stopping 295
speed of sound 95, 131	state space diagram 75	Sun-planet friction 101
speed, highest 35	state, initial 203	sunbeams 406
speed, infinite 370	state, physical 26	sundial 43, 430, 448
sperm motion 35	states 27, 369	sundials 41, 47
Spica 225	static friction 200, 202	superboom 278
spin groups 229	stationary 217	supercavitation 309
spine spine	stationary fluid motion 302	superfluidity 312
	statistical mechanics 210	
human 296	statistical physics	supergiants 225 superlubrication 200
spinning top	definition 319	
angular momentum 109	T - T	supermarket 47
spinors 239	steel 296	supernovae 88
spirituality 207	types 297	superposition 259
Spirograph Nebula 459	steel, stainless 297	support
split personality 206	Stefan-Boltzmann constant	this free pdf 9
sponsor	393	surface 49
this free pdf 9	Steiner's parallel axis theorem	surface tension 201
spoon 121	108	surface tension waves 254
spring 248	steradian 65, 384	surfing 272
spring constant 248	sticking friction 200	surjective 237
spring tides 175	stigma 374	surprises 207, 242
sprinting 150	Stirling's formula 343	swan
squark 474	stone	wake behind 278
St. Louis arch 414	falling into water 258	swarms 364
stadia 47	skipping 311	self-organization in
stainless steel 297	stones 24, 39, 73, 77, 83, 100,	starling 365
staircase formula 150	121, 124, 152, 153, 155, 180,	swell 257
stairs 47	182, 192, 261, 275, 336	swimming speed underwater
stalactite 295	straight line, drawing a 209	309
stalagmites 35	straight lines in nature 57	swimming, olympic 273
stalagtites 359	straightness 48, 57, 62	Swiss cheese 283
standard apple 388	streets 425	switching off the lights 263
standard clock 183	strong coupling constant 391	syllabary 376
standard deviation 368, 388	structure, highest	symbols 377
standard kilogram 93, 121	human-built 51	symbols, mathematical 377
standard pitch 250	subgroup 235	symmetry 235

submarines 274

standing wave 261

Motion Mountain – The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright @ Christoph Schiller November 1997–March 2012

classification 230

SPECIAL

comparison 238	tensegrity structures 296	deduction from clocks 41
crystal, full list 232	tensor 240	flow of 47
discrete 246	tensor order 240	time intervals 42
parity 246	tensor product 378	time is necessary 42
types in nature 243	Tera 385	time measurement, ideal 41
wallpaper, full list 231	tesla 385	time translation 241
symmetry of the whole	testicle 451	time travel 141
Lagrangian 240	testimony 451	time, absolute 42
symmetry operations 235	tetrahedron 63, 106	time, absoluteness of 42
symmetry, external 234	themodynamics 319	time, arrow of 47
symmetry, low 229	thermal energy 321	time, definition of 242
symmetry, summary on 246	thermoacoustic engines 345	time, relative or absolute 52
Système International	Thermodevices 323, 478	time-bandwidth product 267
d'Unités (SI) 383	thermodynamic degree of	Titius's rule 190
system	freedom 329	TNT energy content 393
extended 251	thermodynamics 316, 324	Tocharian 451
system, geocentric 188	first law 104	tog 338
system, heliocentric 188	second law 104	toilet research 360
system, isolated 339	second principle 337	toilet, aeroplane 316
systems, conservative 161	third principle of 334	tokamak 322
systems, dissipative 202	thermodynamics, first	tonne, or ton 385
systems, simple 359	principle 323	topology 50
	thermodynamics,	torque 110
Т	indeterminacy relation of	total momentum 94
Talitrus saltator 45	333	touch 90
tantalum 40	thermodynamics, second	toys, physical 402
Taurus 186	principle of 343	train 97
tax collection 383	thermodynamics, third	trajectory 72
teaching, best method for 9	principle 321	transformation 20
teapot 313	thermodynamics, zeroth	transformations 235
technology 377	principle 320	transforms 100
tectonic activity 168	thermometer 323	translation invariance 42, 50
tectonics 455	thermometry 466	transport 20
teeth growth 351	thermostatics 324	transubstantiation 285
telekinesis 225	third, major 263	tree 388
telephone speed 35	thorn 373, 471	family 27
teleportation 100	thriller 220	tree growth 35
telescope 273	throw 24	tree height, limits to 287
television 24	throw, record 51	tree leaves and Earth rotation
temperature 318-320	throwing speed, record 35, 61	131
introduction 319	throwing, importance of 24	trees 287
measured values 322	thumb 66	trees and pumping of water
temperature scale 466	tidal acceleration 174	464
temperature, absolute 320	tides 127, 168, 172, 454	triboscopes 202
temperature, lowest in	tides and friction 175	tripod 199
universe 322	tides, once or twice per day	tritone 263
temperature, negative 322, 346	192	Trojan asteroids 171
		tropical year 393

deduction 40

truth 285

SYMMETRY

example of 296

tsunamis 257	387	Vendekta 385
tuft 465	universality of gravity 171	Venturi gauge 305
tumbleweed 86	Universe	vernier 62
tunnel	angular momentum 109	video recorder 337
through the Earth 192	universe 27, 204	Vietnam 192
turbulence 305, 358, 367	universe, description by	viewpoint-independence 233
turbulence in pipes 361	universal gravitation 183	viewpoints 229
Turkish 373	universe, two-dimensional 69	Virgo 186
two-dimensional universe 69	Universum 323, 478	vis viva 101
Tycho Brahe 164	unpredictability, practical 205	viscosity
Tyrannosaurus rex 148	URL 399	dynamic 290, 361
,	usenet 399	kinematic 361, 440
U	UTC 41	viscous 201
U(1) 435	•	vocal cords 263
udeko 385	V	voice
Udekta 385	vacuum permeability 391	human 263
UFOs 274	vacuum permittivity 391	void 288
ultimate wheel 475	vacuum wave resistance 392	volcanoes 455
ultrasound	vacuum, human exposure to	volt 385
imaging 266	144	volume 56
ultrasound imaging 266	variability 25	voluntary motion 200, 204
ultrasound motor 199	variable 26	vomit comet 429
umbrella 75	variance 389	vortex
unboundedness 36, 42, 50, 239	variation 26	ring film 315
unboundedness, openness 98	variational principle 218	vortex in bath tub or sink 454
uncertainty	variational principles 208	vortex rings 316
relative 389	Varuna 224	vortex tube 345
total 389	vector 77, 239	vortices 307
uncertainty relation of	vector product 104	Vulcan 172
thermodynamics 333	vector product 104 vector space 77	Vulcanoids 225
underwater swimming 309	vector space, Euclidean 78	vuicanoius 225
underwater, speed records	Vega 225	w
308	Vega at the North pole 132	wafer, consecrated 294
Unicode 376	velars 374	wafer, silicon 294
	velocimeters 405	wake 255, 257
unicycle 91, 298 uniqueness 42, 50	velocity 34, 78, 370	angle 278
unit 338	first cosmic 184	behind ship or swan 279
astronomical 393	•	behind swans and ships
	in space 36	
unitary 237	phase 253	278
units 49, 383	propagation 253	wake angle 279
non-SI 386	second cosmic 184	walking 112, 336
provincial 386	third cosmic 184	walking man
units, SI	wave 253	angular momentum 109
definition 383	velocity as derivative 79	walking on water 258
universal 'law' of gravitation	velocity is not Galilean 36	walking robots 227
152	velocity of birds 36	walking speed 157
universal gravitation 152	velocity, escape 184	walking speed, maximum
origin of name 154	velocity, Galilean 34	animal 427
universal time coordinate 41,	vendeko 385	wallpaper groups 231

TSUNAMIS

Motion Mountain - The Adventure of Physics pdf file available free of charge at www.motionmountain.net Copyright @ Christoph Schiller November 1997–March 2012

washboard roads 354
water 464
origin of oceans 308
patterns in 360
water strider 149
water taps 359
water wave
formation of deep 254
photographs of types 255
solitary 269
thermal capillary 273
types and properties 256
water waves 254, 280
water waves, group velocity of
274
water, speed records under
308
water, walking on 258
water-walking robot 149
watt 385
wave 251
deep water 255
gravity water 254
group 261
long water 255
longitudinal 253
shallow 257
shallow water 255
short water 255
solitary 269
speed
measured values 253
surface tension 254
transverse 254
velocity 253
water 254
wave dispersion
films of types 257
wave motion
six main properties and
effects 259
wave reflection 259

wave, harmonic 253

wave, linear 253 wavelength 253, 260

WASHBOARD

wavelet transformation 277
waves
solitary, in sand 353
waves, in deep water 278
waves, water 280
waw 374
weak mixing angle 391
weather 143, 364
weather, unpredictability of
205
web
list of interesting sites 399
web cam, Foucault's
pendulum 128
web, largest spider 51
weber 385
week 380
week days, order of 191
Weib 451
weight 177, 197
weight of the Moon 189
weight of the Woon 109 weightlessness, feeling of 429
weko 385
Wekta 385
well-tempered intonation 265
whale, blue 272
whales 186, 274, 276
wheel axle, effect of 417
wheel ultimate 23
wheels
vs legs 148
wheels in living beings 85
wheels in nature 87
whip cracking 297
whip, speed of 35
whirl 140
white colour does not last 337
Wien's displacement constant
393
Wilrimodia 222, 478
Wikimedia 323, 478
wind generator
angular momentum 109
wind resistance 201
wine 285

wine arcs 360
wine bottle 100, 117
wings 306
Wirbelrohr 345
wise old man 220
WMAP 428
women 168, 451
women, dangers of looking
after 148
wonder
impossibility of 324
work 101, 102, 218
work, physical 197
world 15, 27
World Geodetic System 395,
457
world question center 402
world-wide web 399
writing 372
wrong 368
wyn 373
X
X xenno 385
<del></del>
xenno 385 Xenta 385
xenno 385 Xenta 385
xenno 385 Xenta 385 Y year, number of days 175
xenno 385 Xenta 385 Y year, number of days 175 yo-yo 156
xenno 385 Xenta 385 Y year, number of days 175 yo-yo 156 yocto 385
xenno 385 Xenta 385 Y year, number of days 175 yo-yo 156 yocto 385 yogh 373
xenno 385 Xenta 385 Y year, number of days 175 yo-yo 156 yocto 385 yogh 373 yot 374
xenno 385 Xenta 385 Y year, number of days 175 yo-yo 156 yocto 385 yogh 373
xenno 385 Xenta 385 Y year, number of days 175 yo-yo 156 yocto 385 yogh 373 yot 374 Yotta 385
xenno 385 Xenta 385 Y year, number of days 175 yo-yo 156 yocto 385 yogh 373 yot 374 Yotta 385
xenno 385 Xenta 385 Y year, number of days 175 yo-yo 156 yocto 385 yogh 373 yot 374 Yotta 385  Z zenith 145
xenno 385 Xenta 385 Y year, number of days 175 yo-yo 156 yocto 385 yogh 373 yot 374 Yotta 385  Z zenith 145 zenith angle 154
xenno 385 Xenta 385 Y year, number of days 175 yo-yo 156 yocto 385 yogh 373 yot 374 Yotta 385  Z zenith 145 zenith angle 154 zepto 385
xenno 385 Xenta 385 Y year, number of days 175 yo-yo 156 yocto 385 yogh 373 yot 374 Yotta 385  Z zenith 145 zenith angle 154 zepto 385 zero 377
xenno 385 Xenta 385 Y year, number of days 175 yo-yo 156 yocto 385 yogh 373 yot 374 Yotta 385  Z zenith 145 zenith angle 154 zepto 385 zero 377 zero gravity 429
xenno 385 Xenta 385  Y year, number of days 175 yo-yo 156 yocto 385 yogh 373 yot 374 Yotta 385  Z zenith 145 zenith angle 154 zepto 385 zero 377 zero gravity 429 Zetta 385
xenno 385 Xenta 385 Y year, number of days 175 yo-yo 156 yocto 385 yogh 373 yot 374 Yotta 385  Z zenith 145 zenith angle 154 zepto 385 zero 377 zero gravity 429

ZODIAC 511



## **MOTION MOUNTAIN**

## The Adventure of Physics - Vol. I

### Fall, Flow and Heat

Why do change and motion exist?

How does a rainbow form?

What is the most fantastic voyage possible?

Is 'empty space' really empty?

How can one levitate things?

At what distance between two points does it become impossible to find room for a third one in between?

What does 'quantum' mean?

Which problems in physics are unsolved?



Answering these and other questions on motion, this series gives an entertaining and mind-twisting introduction into modern physics – one that is surprising and challenging on every page.

Starting from everyday life, the adventure provides an overview of the recent results in mechanics, thermodynamics, electrodynamics, relativity, quantum theory, quantum gravity and unification. It is written for undergraduate students and for anybody interested in physics.

Christoph Schiller, PhD Université Libre de Bruxelles, is a physicist with more than 25 years of experience in the presentation of physical topics.

Pdf file available free of charge at www.motionmountain.net